

University of New Hampshire

University of New Hampshire Scholars' Repository

Space Science Center

Institute for the Study of Earth, Oceans, and
Space (EOS)

2009

GRB Polarimetry with POET

Mark L. McConnell

University of New Hampshire - Main Campus, mark.mcconnell@unh.edu

L Angelini

NASA Goddard Space Flight Ctr.

Matthew Baring

Rice University

Scott Barthelmy

NASA Goddard Space Flight Ctr.

J Kevin Black

Rock Creek Scientific

See next page for additional authors

Follow this and additional works at: <https://scholars.unh.edu/ssc>



Part of the [Astrophysics and Astronomy Commons](#)

Recommended Citation

GRB Polarimetry with POET McConnell, M. L. and Angelini, L. and Baring, M. G. and Barthelmy, S. and Black, J. K. and Bloser, P. F. and Dennis, B. and Emslie, A. G. and Greiner, J. and Hajdas, W. and Harding, A. K. and Hartmann, D. H. and Hill, J. E. and Ioka, K. and Kaaret, P. and Kanbach, G. and Kniffen, D. and Legere, J. S. and Macri, J. R. and Morris, R. and Nakamura, T. and Produit, N. and Ryan, J. M. and Sakamoto, T. and Toma, K. and Wu, X. and Yamazaki, R. and Zhang, B., AIP Conference Proceedings, 1133, 64-66 (2009), DOI:<http://dx.doi.org/10.1063/1.3155969>

This Conference Proceeding is brought to you for free and open access by the Institute for the Study of Earth, Oceans, and Space (EOS) at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Space Science Center by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

Authors

Mark L. McConnell, L Angelini, Matthew Baring, Scott Barthelmy, J Kevin Black, Peter F. Bloser, B Dennis, A G. Emslie, J Greiner, W Hajdas, A K. Harding, H Hartmann, J E. Hill, Kunihito Ioka, Philip Kaaret, G Kanbach, D Kniffen, Jason S. Legere, John R. Macri, R Morris, Takashi Nakamura, N Produit, James M. Ryan, Takanori Sakamoto, Kenji Toma, X Wu, Ryo Yamazaki, and Bing Zhang



GRB Polarimetry with POET

M. L. McConnell, L. Angelini, M. G. Baring, S. Barthelmy, J. K. Black, P. F. Bloser, B. Dennis, A. G. Emslie, J. Greiner, W. Hajdas, A. K. Harding, D. H. Hartmann, J. E. Hill, K. Ioka, P. Kaaret, G. Kanbach, D. Kniffen, J. S. Legere, J. R. Macri, R. Morris, T. Nakamura, N. Produit, J. M. Ryan, T. Sakamoto, K. Toma, X. Wu, R. Yamazaki, and B. Zhang

Citation: [AIP Conference Proceedings](#) **1133**, 64 (2009); doi: 10.1063/1.3155969

View online: <http://dx.doi.org/10.1063/1.3155969>

View Table of Contents: <http://scitation.aip.org/content/aip/proceeding/aipcp/1133?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[Development of the GRB Monitor for the CALET Experiment](#)

AIP Conf. Proc. **1133**, 88 (2009); 10.1063/1.3155979

[Simulation Study of Expected Performance of MAXI for GRB observations](#)

AIP Conf. Proc. **1133**, 82 (2009); 10.1063/1.3155976

[GRB 080514B: the first highenergy AGILE burst with optical/NIR afterglow](#)

AIP Conf. Proc. **1133**, 58 (2009); 10.1063/1.3155967

[The Correlation of Spectral Lag Evolution with Prompt Optical Emission in GRB 080319B](#)

AIP Conf. Proc. **1133**, 356 (2009); 10.1063/1.3155918

[The GRB FollowUp Project at the Xinglong Observatory: 20062008](#)

AIP Conf. Proc. **1065**, 107 (2008); 10.1063/1.3027892

GRB Polarimetry with POET

M. L. McConnell^a, L. Angelini^b, M. G. Baring^c, S. Barthelmy^b, J. K. Black^d, P. F. Bloser^a, B. Dennis^b, A. G. Emslie^e, J. Greiner^f, W. Hajdas^g, A. K. Harding^b, D. H. Hartmann^h, J. E. Hill^{i,b}, K. Ioka^j, P. Kaaret^k, G. Kanbach^f, D. Kniffen^l, J. S. Legere^a, J. R. Macri^a, R. Morris^l, T. Nakamura^j, N. Produit^m, J. M. Ryanⁿ, T. Sakamoto^b, K. Toma^o, X. Wu^p, R. Yamazaki^q and B. Zhang^r

^aSpace Science Center, University of New Hampshire, Durham, NH 03824

^bNASA Goddard Space Flight Center, Greenbelt, Md 20771, USA

^cDepartment of Physics and Astronomy, Rice University, Houston, TX 77251, USA

^dRock Creek Scientific, Silver Spring, MD 20910, USA

^eDepartment of Physics, Oklahoma State University, Stillwater, OK 74075, USA

^fMax Planck Institute for Extraterrestrial Physics, 85748 Garching, Germany

^gPaul Scherrer Institute, 5232 Viligen PSI, Switzerland

^hDepartment of Physics and Astronomy, Clemson University, Clemson, SC 29634, USA

ⁱUniversities Space Research Association, CRESST

^jTheoretical Astrophysics Group, Department of Physics, Kyoto University, Sakyo-ku, Kyoto 606-8502, Japan

^kDepartment of Physics and Astronomy, University of Iowa, Iowa City, IA 52242, USA

^lUniversities Space Research Association, Columbia, MD 21044, USA

^mINTEGRAL Science Data Center, 1290 Versoix, Switzerland

ⁿSpace Science Center, University of New Hampshire, Durham, NH 03824, USA

^oDepartment of Astronomy and Astrophysics, Pennsylvania State University, University Park, PA 16802, USA

^pPurple Mountain Observatory, Chinese Academy of Sciences, Nanjing 210008, China

^qDepartment of Physical Science, Hiroshima University, Higashi-Hiroshima 739-8526, Japan

^rDepartment of Physics, University of Nevada, Las Vegas, NV 89154, USA

Abstract. POET (Polarimeters for Energetic Transients) represents a concept for a Small Explorer (SMEX) satellite mission, whose principal scientific goal is to understand the structure of GRB sources through sensitive X-ray and γ -ray polarization measurements. The payload consists of two wide field-of-view (FoV) instruments: a Low Energy Polarimeter (LEP) capable of polarization measurements in the energy range from 2-15 keV and a high energy polarimeter (Gamma-Ray Polarimeter Experiment or GRAPE) that would measure polarization in the 60-500 keV energy range. The POET spacecraft provides a zenith-pointed platform for maximizing the exposure to deep space. Spacecraft rotation provides a means of effectively dealing with any residual systematic effects in the polarization response. POET provides sufficient sensitivity and sky coverage to measure statistically significant polarization (for polarization levels in excess of 20%) for ~ 80 GRBs in a two-year mission. High energy polarization data would also be obtained for SGRs, solar flares, pulsars and other sources of astronomical interest.

Keywords: Gamma-ray Bursts, Polarimetry, X-ray, Gamma-ray

PACS: 95.55.Ka

INTRODUCTION

Gamma-ray bursts (GRBs) are the most energetic events in the universe, and have stimulated intense observational and theoretical research. Theoretical models indicate that a refined understanding of the inner structure of GRBs, including the geometry and physical processes close to the central engine, requires the exploitation of high energy X-ray and γ -ray polarimetry. To date, observations have been of limited sensitivity and subject to poorly understood systematics. Polarimeters for Energetic Transients (POET) is a SMEX mission concept that is capable of measuring the high energy polarization of GRBs and other sources of astronomical interest [1]. POET obtains measurements with two different polarimeters (both with wide fields of view) to provide observations over a broad energy range: LEP (Low Energy Polarimeter) covers 2–15 keV and GRAPE (Gamma-RAY Polarimeter Experiment) covers 60–500 keV. In the context of studying GRBs, POET would address the following questions : What is the magnetic structure of the jets? What is the geometric structure of the jets? What is the prompt emission mechanism? Where does the emission originate? For




FIGURE 1. The LEP TPC polarimeter uses a simple strip readout and time of arrival to form a pixelized image of a photoelectron track. The TPC polarimeter forms an image by digitizing the signal on each readout strip. The signal from a 6 keV X-ray shows the interaction point, emission angle and end of the photoelectron track. The size of each circle is proportional to the deposited charge in each virtual ($132 \mu\text{m}$) pixel.




FIGURE 2. A histogram of reconstructed polarized 6.4 keV photoelectron tracks demonstrates a modulation of 45%.

example, by measuring a distribution of GRB polarizations, POET could distinguish between jet models with ordered magnetic fields and those with random magnetic fields, each of which predicts different distributions of polarization values [2, 3].

LOW ENERGY POLARIMETER (LEP)

LEP measures the direction of photoelectrons that are ejected after the photoelectric absorption of an incident photon. Photoelectrons tend to be ejected in a direction parallel to the electric field vector of the incident photon. The photoelectron tracks are measured with the innovative operation of a Time Projection Chamber (TPC), a proven technology used in high-energy particle physics. The design provides unmatched polarization sensitivity over the 2–15 keV band-pass [4, 5]. The LEP polarimeter enclosure consists of four dual-readout detector modules each with an isolated gas volume contained by a Be X-ray window. Each detector module contains two $6 \times 12 \times 24 \text{ cm}^3$ TPCs. Each TPC is comprised of a micropattern proportional counter, consisting of a shared drift electrode and a high-field gas electron multiplier (GEM) positioned 1 mm from a strip readout plane. When an X-ray is absorbed in the gas between the drift electrode and the GEM, a photoelectron is ejected in a preferential direction with a $\cos^2 \phi$ distribution, where ϕ is the azimuthal angle measured from the X-ray polarization vector. As the photoelectron travels through the gas it creates a path of ionization that drifts in a moderate, uniform field to the GEM where an avalanche occurs. Figure 1 illustrates how a track image projected onto the x-y plane is formed by digitizing the charge pulse waveforms. The coordinates are defined by strip location in one dimension, and arrival time multiplied by the drift velocity in the orthogonal dimension. The strips are smaller than the mean free path of the photoelectron and therefore an image of the track can be reconstructed and the initial direction of the photoelectron determined. The magnitude and orientation of the source polarization can be determined from a histogram of the emission angles (Figure 2), where the polarization angle corresponds to the maximum in the modulation pattern. LEP is sensitive to GRB emissions at off-axis angles up to at least 45° .

GAMMA-RAY POLARIMETER EXPERIMENT (GRAPE)

GRAPE [6, 7, 8, 9] is designed to measure polarization from 60-500 keV and to provide spectroscopy over a broad energy range from 15 keV to 1 MeV. It relies on the fact that a Compton scattered photon tends to scatter at right angles to the electric field vector of the incident photon. The GRAPE instrument is composed of 64 independent detector modules arranged in two identical assemblies that provide the associated electronics and the required mechanical and thermal support. Each polarimeter module incorporates an array of optically independent $5 \times 5 \times 50 \text{ mm}^3$ scintillator elements aligned with and optically coupled to the 8×8 scintillation light sensors of a 64-channel MAPMT. Two types of scintillators are employed. Low-Z plastic scintillator is used as an effective medium for Compton scattering. High-Z inorganic scintillator (Bismuth Germanate, BGO) is used as a calorimeter, for absorbing the full energy of the




FIGURE 3. GRAPE measures photons that scatter from a plastic element to a BGO element.




FIGURE 4. A histogram of azimuthal scatter angles at a incident photon energy of 288 keV, as measured in the lab. The modulation provides a measure of both the polarization fraction and the polarization angle of the incident radiation.




FIGURE 5. The effective area of the POET instruments for a GRB that is 30° off-axis, illustrating the wide FoV. For GRAPE, the sensitivity to both polarization and spectroscopy are shown.

scattered photon. Valid polarimeter events are those in which a photon Compton scatters in one of the plastic elements and is subsequently absorbed in one of the BGO elements (Figure 3). The azimuthal scatter angle is determined for each valid event by the relative locations of hit scintillator elements. To facilitate spectral measurements over a broader energy range (15 keV–1 MeV), GRAPE also includes 2 independent NaI(Tl) spectrometer modules. The magnitude and orientation of the source polarization can be determined from a histogram of the emission angles (Figure 4), where the polarization angle corresponds to the minimum in the modulation pattern. GRAPE will be sensitive to GRB emissions at off-axis angles up to at least 60°.

THE POET MISSION

The mission concept involves a standard SMEX launch into a circular 600 km orbit. A zenith-pointed spacecraft provides continuous exposure to deep space. Coupled with the wide FoV of both instruments (Figure 5), this maximizes the probability of prompt GRB detection. Spacecraft rotation mitigates the effects of any residual systematic effects in the polarization response. The combined capabilities of LEP and GRAPE will allow us to measure the polarization over a broad energy range (2–500 keV) and also to measure E_p for a large number of GRBs. Simulations based on realistic GRB distributions indicate that both LEP and GRAPE will both be capable of detecting about 40 GRBs per year with a polarization sensitivity of about 20% [1] in their respective energy bands. During a nominal two-year mission, POET would provide statistically significant polarization measurements for ~100 GRBs, providing diagnostic information on the GRB emission mechanism that cannot be obtained from currently-available data [3].

REFERENCES

1. J. E. Hill, M. L. McConnell, P. Bloser, J. Legere, J. Macri, J. Ryan, S. Barthelmy, L. Angelini, T. Sakamoto, J. K. Black, D. H. Hartmann, P. Kaaret, B. Zhang, K. Ioka, T. Nakamura, K. Toma, R. Yamazaki and X. Wu, *AIP Conf. Proc.*, **1065**, pp. 331-337 (2008).
2. E. Waxman, *Nature* **423**, 388-389 (2003).
3. K. Toma, T. Sakamoto, B. Zhang, J. E. Hill, M. L. McConnell, P. F. Bloser, R. Yamazaki, K. Ioka and T. Nakamura, *ApJ*, submitted (2009), arXiv:0812.2483 [astro-ph].
4. J. K. Black, R. G. Baker, P. Deines-Jones, J. E. Hill and K. Jahoda, *NIM A*, **581**, 755-760 (2007).
5. J. E. Hill, S. Barthelmy, J. K. Black, P. Deines-Jones, K. Jahoda, T. Sakamoto, P. Kaaret, M. L. McConnell, P. F. Bloser, J. R. Macri, J. S. Legere, J. M. Ryan, B. R. Smith and B. Zhang, *Proc. SPIE* **6686**, 66860Y (2007).
6. M. L. McConnell, J. Ledoux, J. Macri and J. Ryan, *Proc SPIE* **5165**, 334-345 (2004).
7. J. Legere, P. Bloser, J. R. Macri, M. L. McConnell, T. Narita, J. M. Ryan, *Proc. SPIE* **5898**, 413 (2005).
8. P. F. Bloser, J. S. Legere, M. L. McConnell, J. R. Macri, C. M. Bancroft, T. P. Connor and J. M. Ryan, *NIM A*, in press (2008), doi:10.1016/j.nima.2008.11.118.
9. M. L. McConnell, C. M. Bancroft, P. F. Bloser, T. P. Connor, J. S. Legere, J. R. Macri and J. M. Ryan, these proceedings (2009).