

University of New Hampshire

University of New Hampshire Scholars' Repository

Physics Scholarship

Physics

4-1-2013

New Insight into the Inner Magnetosphere Plasma Regimes with the van Allen Probes (RBSP)

Joerg-Micha Jahn

Richard E. Denton

Herbert O. Funsten

Los Alamos National Laboratory

Geoff Reeves

Harlan E. Spence

University of New Hampshire, harlan.spence@unh.edu

Follow this and additional works at: https://scholars.unh.edu/physics_facpub



Part of the [Physics Commons](#)

Recommended Citation

Jahn, Joerg-Micha; Denton, Richard E.; Funsten, Herbert O.; Reeves, Geoff; and Spence, Harlan E., "New Insight into the Inner Magnetosphere Plasma Regimes with the van Allen Probes (RBSP)" (2013). *Geophysical Research Abstracts*. 161.

https://scholars.unh.edu/physics_facpub/161

This Article is brought to you for free and open access by the Physics at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Physics Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.



New Insight into the Inner Magnetosphere Plasma Regimes with the van Allen Probes (RBSP)

Joerg-Micha Jahn (1), Richard E Denton (2), Herbert O Funsten (3), Geoff Reeves (3), and Harlan E Spence (4)

(1) Southwest Research Institute, Space Science Department, San Antonio, TX, United States (jjahn@swri.edu, +1- 210-520-9935), (2) Dartmouth College, Department of Physics and Astronomy, Hanover, NH, United States, (3) ISR Division, Los Alamos National Laboratory, Los Alamos, NM 87545, United States, (4) Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, NH, 03824, United States

The recent successful launch of the twin van Allen spacecraft (formerly known as RBSP) provides a new and unprecedented window into the structure and dynamics of inner magnetospheric plasma content and dynamics. The equatorially orbiting van Allen spacecraft are returning clean high resolution, very low background ion composition and electron plasma data throughout the radiation belt and ring current region inside geosynchronous orbit. Since both van Allen spacecraft are positioned in near-identical chase orbits, lapping each other continuously throughout the mission, we are able to study both spatial and temporal variability in the inner magnetosphere with unprecedented resolution on a range of time and length scales. In this paper we are presenting initial results from plasma composition measurements in the nightside of Earth's magnetosphere, focussing on plasma fractional plasma composition of H⁺, He⁺, and O⁺ in the plasmasphere through lower ring current energies (< 50 keV). Early results do not only indicate a remarkable spatial and temporal variability in plasma ion composition in the inner magnetosphere, they also show frequent occurrences of multiple peak energy distributions in this energy range. Multi-peaked energy distributions with several peaks occurring in ring current, plasmasphere and (less often) plasma sheet are frequently observed, with distinct differences between the three ion species. Energy spectra with 5-6 peaks for a single species have been observed repeatedly.