

Spring 2015

Effects of Proline and Glycine on the Cnidocyte Discharge of *Hydra magnipapillata*

Janine R. Appleton

University of New Hampshire, janineappleton23@gmail.com

Follow this and additional works at: <http://scholars.unh.edu/honors>

 Part of the [Behavior and Ethology Commons](#), [Biodiversity Commons](#), [Cell Biology Commons](#), [Developmental Biology Commons](#), [Evolution Commons](#), [Integrative Biology Commons](#), and the [Molecular Genetics Commons](#)

Recommended Citation

Appleton, Janine R., "Effects of Proline and Glycine on the Cnidocyte Discharge of *Hydra magnipapillata*" (2015). *Honors Theses and Capstones*. 256.

<http://scholars.unh.edu/honors/256>

This Senior Honors Thesis is brought to you for free and open access by the Student Scholarship at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Honors Theses and Capstones by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact nicole.hentz@unh.edu.

Effects of Proline and Glycine on the Cnidocyte Discharge of *Hydra magnipapillata*

Abstract

The sense of taste enables animals to utilize environmental cues to detect favorable foods. Through specialized sensory receptors, Cnidarians employ stinging cells called cnidocytes to perform a variety of activities such as locomotion, capturing prey, inducing of feeding responses, and defense. Their discharge is highly regulated by mechanical and chemical signals that are mediated by a complex system including the opsin and taste pathways. Taste 1 Receptors (T1R) have previously been isolated in vertebrates but only until recently, have been noted in invertebrates. Receptors specific to L- amino acids corresponding to the taste sensation of umami, were studied to determine if the pathways of *Hydra magnipapillata* used for feeding were similar to the systems utilized in vertebrates. Amino acids, Proline and Glycine, were experimented using cnidocyte assays to induce feeding and capture cnidocytes. An optimal concentration of 10mM of Proline and Glycine was tested and found to be significant by eliciting greater cnidocyte discharge as compared to a Control of gelatin with P- Values of 0.003 and 0.0011 respectively. This indicates that amino acids, which have similar receptors in vertebrates, are capable of inducing feeding responses in invertebrates implying that T1Rs operate in similar mechanisms thus predating the current notion of the evolutions and diversification of such genes by around 400 million years.

Keywords

Cnidocytes; Opsin; Amino Acid; Taste Receptors; Umami; Invertebrate

Subject Categories

Behavior and Ethology | Biodiversity | Cell Biology | Developmental Biology | Evolution | Integrative Biology | Molecular Genetics

Effects of Proline and Glycine on the Cnidocyte Discharge of
Hydra magnipapillata

Janine Appleton

Honors Senior Thesis

Spring 2015

Thesis Advisor: Dr. David Plachetzki

University of New Hampshire

Table of Contents

Abstract	3
Introduction	4
Materials and Methods	10
Results	13
Discussion	21
References	23
Appendix	26
URC Poster	30

Abstract

The sense of taste enables animals to utilize environmental cues to detect favorable foods. Through specialized sensory receptors, Cnidarians employ stinging cells called cnidocytes to perform a variety of activities such as locomotion, capturing prey, inducing of feeding responses, and defense. Their discharge is highly regulated by mechanical and chemical signals that are mediated by a complex system including the opsin and taste pathways. Taste 1 Receptors (T1R) have previously been isolated in vertebrates but only until recently, have been noted in invertebrates. Receptors specific to L- amino acids corresponding to the taste sensation of umami, were studied to determine if the pathways of *Hydra magnipapillata* used for feeding were similar to the systems utilized in vertebrates. Amino acids, Proline and Glycine, were experimented using cnidocyte assays to induce feeding and capture cnidocytes. An optimal concentration of 10mM of Proline and Glycine was tested and found to be significant by eliciting greater cnidocyte discharge as compared to a Control of gelatin with P- Values of 0.003 and 0.0011 respectively. This indicates that amino acids, which have similar receptors in vertebrates, are capable of inducing feeding responses in invertebrates implying that T1Rs operate in similar mechanisms thus predating the current notion of the evolutions and diversification of such genes by around 400 million years.

Introduction

Taste 1 Receptors

The taste 1 receptor (T1R) family of G-protein-coupled receptors (GPCR) consists of three different subunits, T1R1, T1R2 and T1R3 all functioning to detect a range of sweetness. These taste receptors operate as chemoreceptors interacting with ligands or other taste stimuli to produce a taste perception in the brain. The T1Rs bind to G proteins, usually gustuctin Ga but also Gao and Gai, which activate and inhibit adenylyl cyclases and thus regulate cyclic adenosine monophosphate and cyclic guanosine monophosphates (cGMP) levels². These subparts combine into heterodimers as they are not functional independently. The heterodimer T1R2-T1R3 works in conjunction with the G-protein gustducin, to act as a sweet sense while the heterodimer T1R1- T1R3 senses more a natural sweetness and savor expressing the umami sensations of monosodium glutamate (MSG)¹⁸.

Taste cells in vertebrates are often small bipolar cells that have a relatively short lifetime of 10 days⁸. The cells are packed densely together in groups of 50-150 including precursor cells, support cells, and taste receptor cells²¹. The cells are generally located on the tongue within papillae, the palate, and the pharynx with distinction per type of receptor. The T1R1+3 receptors are found specifically in the fungiform papillae on the tongue and the palate on the roof of the mouth. The T1R2+3 receptors on the other hand, are found in the circumvallate papillae and foliate papillae on the back of the tongue and on the palate on the roof of the mouth². When activated, the T1R1+3 taste receptors synapse on the chorda tympani nerves sending signals to the brain while the T1R2+3 also acts on the glossopharyngeal nerves².

Physiology of Taste T1Rs

Taste receptor cells are innervated by nerves that transmit information through the synapse to the brain in order to develop of sort of taste perception²¹. The signal passes through the basolateral side of taste cells via axonal membranes consisting of voltage gated Na⁺, K⁺, and Ca²⁺ channels¹⁷. The ligands bind to surface chemoreceptors to activate the GTP- binding protein which causes either G α to induce the cAMP production via adenylate cyclase activation further activating protein kinase A to initiate actin polymerization and elongation of sensory cells or the activate G $\beta\gamma$ to inhibit the cAMP pathway by decreasing adenylate cyclase activity^{1,5}. After this chemical stimulation and GPCR activation, the channels become depolarized raising Ca²⁺ levels allowing for action potentials to release through the afferent axons and transmission to occur¹⁷. The linking of sensory outputs has been recently under investigation. The visual perception of animals is mediated by an opsin-mediated phototransduction cascade. In this process, GPCR signaling pathways are regulated by the depolarization or polarization of ion channels¹¹.

Umami as a Sense

Umami is related to MSG signifying savor or tastiness represented by the heterodimer of T1R1+3¹⁰. As it interacts with L- amino acids, it becomes very sensitive to glutamate, inosine monophosphate (IMP), and guanosine monophosphate (GMP) which act to enhance many of the amino acids⁸. The umami sensation is unique from other taste senses as it contains a truncated form of brain mGluR4 which is a metabotropic GPCR originating from the central nervous system. The receptor became adaptive to the high glutamate concentrations from food linking its association to the ribonucleotides¹¹. L-glutamate helps to facilitate the intake of peptides, specifically CCKs, and proteins further heightened by the presence of purines⁸.

Animals and Taste

The sense of taste is a necessity for animals as it allows them to systematically pick out food that yields nutritional benefits to them. Most animals respond to food via chemoreception with a lot of focus spent on the effects of amino acids on the stimulation of feeding in aquatic animals. This was first introduced by Nagle in 1802 who proved that dead objects were rejected by sea anemones but objects saturated with food were willingly ingested¹⁶. Furthermore, a study conducted by Pantin in 1942 tested a variety of food extracts, amino acids, proteins, carbohydrates, and fatty acids on their ability to sensitize the discharges of cnidae. Pantin found that minor discharge rates occurred with just mechanical stimulation, and in order to achieve optimal discharge rates, both chemical and mechanical stimulation was required¹⁹. The concept that objects needed to be infused with food in order to elicit a response indicated that the stimulus was more sensory related than strictly impulse. This was further studied in a groundbreaking study performed by Loomis, who analyzed the feeding responses of *Hydra littoralis* to glutathione. He found that interaction with glutathione resulted in contraction and writhing of the tentacles and as the concentration of glutathione present increased, the probability that the hydra's tentacles would retract pulling food in toward its mouth also increased⁴.

It has been noted in studies that there is a relationship with different chemoreceptors systems in one organism where one amino acid may control the movement of the tentacles while another may control the ingestion of food²⁰. The synaptic input to the cnidocytes following stimulus of food implies that sensory nerve cells may serve to detect such stimulus and sensitize the cnidocyte prior to mechanical stimulus and that chemical and mechanical cues from prey help to regulate feeding responses^{16, 20}.

The Phylogeny of T1Rs in Cnidaria

The presence of T1R in Cnidaria was previously thought to be non-existent due to the absence of the receptors during the branching of chordates. Recent phylogenomic analysis, however, has since determined that T1Rs are in fact present in the cnidarian genomes¹⁴. Plachetzki's ability to determine that differing light levels affect the discharge of cnidocytes links the opsin pathways to cnidocytes and thus mechanical and chemical cues¹⁴. Plachetzki was able to determine that the opsin in the sensory neurons connects to the cnidocytes as well as being complemented by the presence of the cyclic nucleotide-gated ion channels which are required for signal transmission¹²³. The discovery that cyclic nucleotide-gated channels are the ancestral state of the opsin-mediated photosensitivity helps to prove that hydra possess functional opsins despite not having eyes¹². This suggests that the T1Rs were present prior to the Cambrian Explosion and lost independently thereafter during the lineage diversification of protostomes and tunicates¹².

Cnidocyte Release in Hydra

Hydra are Cnidaria that have been evolving for over 600 million years, producing asexually through budding and acting as clones to one another. Hydra are the most primitive organism to have a nervous system with neurons dispersed within the epithelial layers to form networks and bidirectional synapses with non-nervous cells^{6, 15}. Organized by radial symmetry, hydra have a mouth encircled by tentacles lined with poison-filled cnidocytes. Cnidocytes are found exclusively in Cnidaria and represent one of the most complex cell types known in animals¹⁹. When stimulated, the cnidocytes expel an energetically expensive secretion product called cnidae

in as fast as three milliseconds that are used for multiple operations like feeding, locomotion, and defense^{1,3}.

The cnidae discharge is under the direct control of the cnidocyte acting as a specialized secretory doubling as a sensory cell with elaborate communication with the nervous system¹⁹. There are five steps to Cnidarian feeding responses which include: contact of prey to a tentacle, cnidocyte release, contraction of tentacles to the mouth, the opening of the mouth, and finally, ingestion of the prey⁹. The bodily fluids released from the prey after an attack consist of certain chemicals that match receptors on the predator to attract or detract them and elicit feeding reflexes⁹.

The cnidae can be discharged due to a variety of reasons such as responding to mechanical stimuli with or without chemosensitization and also due to vibrational frequencies³. The discharge is a multi-cellular occurrence requiring both the cnidocyte and surrounding cells, together called the cnidocyte supporting cell complex. This episode is mainly driven by the high pressurized system within the capsule showing the mechanisms acts a sensory neuron^{3,5}. Stimulation of the tentacles causes the tertiary structure of the inactive protein on the surface of cells to alter, activating the protein and thus eliciting a depolarizing effect as the surplus of calcium ions stored in the capsule get released through the opening of the ion channels^{5,7}. The concentration gradient yields an influx of water and thus pushes out the cnidae onto the prey.

Plachetzki Laboratory

Research in this laboratory is centered on determining the evolution of sensory genes in Cnidarians, specifically *Hydra magnipapillata*. Dr. Plachetzki's previous work determined that the role of photosensitivity in hydra and more specifically, that hydra, an eyeless organism, have

the ability to use visionary cues through the use of the cyclic nucleotide gated channels. This proved that the opsin genes used for vision in humans may also be attributed to phototransduction cascades in invertebrates. Other members of this lab work to analyze and isolate the opsin pathways within cnidocytes and surrounding cells to determine if taste and vision follow the same transduction cascades. Through double in situ hybridization and cloning of specific sensory genes, the laboratory will be about to use fluorescent markers to localize and show overlap of cell types within the hydra. This research helps to determine if the opsin pathways mediate cnidocyte discharge and thus how cnidarians are able to sense their environments.

Experiment and Expectations

In this study, it was hypothesized that amino acids, specifically Proline and Glycine, were capable of inducing feeding responses as dictated by the T1Rs in Hydra. Both chemical and mechanical stimulation were utilized to determine reflexes with an analysis based quantitatively on the cnidocyte assay and number of probes associated with the desired responses. Compared to a control group, the amino acid coated probes were expected to elicit greater cnidocyte discharge rates. Since vertebrates have a receptor for these amino acids, a behavioral response from the Hydra to Proline and Glycine would indicate that the feeding mechanisms in invertebrates operate similarly to vertebrates undertaking comparable pathways of T1Rs. This would construct a hypothesis on the evolution of T1Rs in invertebrates predating the current understandings of the evolutions and diversification of such genes by around 400 million years.

Materials and Methods

Handling and Preparation of the Experimental Animals

Specimens of *Hydra magnipapillata* in this laboratory are asexual and reproduce via budding. *H. magnipapillata* were kept in glass bowls containing hydra medium (1M CaCl₂, 80mM MgSO₄, 100mM MgCl₂, 1.5M NaHCO₃, 30mM KNO₃) and were washed and fed every day. The *H. magnipapillata* were fed Artemia, a salt water- grown marine invertebrate, following standard protocols post washing with new hydra medium. The placement of the Hydra in the laboratory was exposed to the natural daylight cycle while the temperature was regulated to maintain 23°C. In preparation for experimentation, 6-9 plastic FisherBrand polystyrene 100mm X 15mm Petri dishes were obtained and filled halfway with hydra medium. About 9-12 *H. magnipapillata* were removed from the holding dishes and placed in the new Petri dishes via a plastic pipette. The dishes were covered and the *H. magnipapillata* were left untouched and unfed for 24-30 hours. After the allotted time, the Petri dishes were moved to the dark room to be left untouched for 45 minutes.

Preparation of the Solution Probes

Knox unflavored gelatin was obtained as well as Craftsmart Plastalina Modeling Clay. To act as a control for the experiment, 20mg of gelatin was measured with the Sartorius weigh scale and added to a USA Scientific 15mL conical screw cap centrifuge tube to be mixed with 1000mL of hydra medium. L- Proline and Glycine were acquired from Sigma Aldrich and Fisher Scientific respectively. Each amino acid was mixed with milliQ water to produce a stock solution following standard procedures. For preparation of the experiment, variations of milliliters of Proline were used, specifically 1µl, 10µl and 100µl of the stock solution, and were

mixed with 999uL, 990uL and 900uL respectively of hydra medium with Eppendorf pipets and pipet tips as well as with 20mg of gelatin to produce a 1mM, 10mM, and 100mM solution in a USA scientific 15mL centrifuge tube. The second experiment utilized 10µl Glycine mixed with 990µl hydra medium and 20mg of gelatin producing a 10mM solution.

Preparation of the Probes

The Eppendorf Thermomixer was turned on and heated to 36°C. Both the control and experimental tubes were then placed in holders of the Thermomixer and left to liquefy with periodic mixing for 15 minutes. The probes were made from LebcO Omniflex Monofilament fishing line which was 0.008 inches in diameter and a 4lb test weight. A range of 36- 72 pieces of fishing line was cut at 2-3 inches per piece. Depending on the number of hydra being tested, half to one third of the probes were dipped in the into the 36°C control of gelatin and the other half and thirds were dipped into the 36°C amino acid containing tubes. The procedure for dipping consisted of raising and lowering the probes in the solutions three times slowly to minimize the occurrence of air bubbles. Large Petri dishes with a long rod of rolled clay centered in the middle held the dipped probes elevated to prevent drip and surface adhesion. Each tube was maintained at the 36°C temperature range throughout the process. After one round of dipping, the Thermomixer temperature was lowered to 32°C and left for 15 minutes in order for the solutions cool. The control and amino acid based probes were then re-dipped in the respective tubes with the same procedure as before and placed in the dishes and left to dry with the tops on for two hours.

The Cnidocyte Assay and Probing Mechanisms

Three trials were performed with the gelatin- coated control probes and the varying concentrations of solubilized amino acids in gelatin- coated probes. Each trial consisted of six- eight hydra individuals that were probed to trigger cnidocyte discharge. Two sets of 3-4 probes were wet mounted on a FisherBrand Microscope Slide coated with three drops of Amresco Glycerol and topped with a Corning Cover Glass to ease readability and lessen overlap. The probing mechanism consisted of identifying individual hydra that had lengthened and extended tentacles to ease the entry and exit points prior and post contraction of the probes thus limiting attachment of the hydra to the probe. Each individual's distal tentacle was grazed with the distal end of the gelatin coated probe for a count of one. The probe would only be mounted if a) the hydra contracted immediately after probing b) multiple tentacles were not touched in passing and c) the hydra did not attached to the probe. Rotations of trial one would begin with the control and followed by the amino acid dipped probes each utilizing a different Petri dish of hydra. Upon completion, commencement of the second and third trials would occur in the same fashion. Preparation of the experiments was performed in the mornings while counting procedures were performed in the afternoon to evening settings of each day however, it was verified that the light settings remained consistent throughout the research.

Microscopy

The wet mounts and resulting discharged cnidocytes were analyzed under the Differential Interference Contrast (DIC) microscopy. The probes were counted with a LEICA DM 2500 microscope at 10X with follow-ups done at 40X. The analysis accounted for the full length and width of the probe requiring appropriate manual focusing to get a full view of cnidocytes present. Probes were recorded even if no cnidocytes were present however, statistical data does exclude probes with zero values.

Results

Recording of Data

The discharge of cnidocytes in *Hydra* are dictated by both chemical and mechanical cues from the environment ranging from responses due to phototaxis, light exposure, feeding mechanisms and roles of defense as driven by the opsin- mediated phototransduction cascade. A contraction of the body of the *Hydra* into a ball can indicate both adverse and favorable discharge trends. In order to test hypothesis of the optimal concentration of amino acid that would elicit a response to feeding, a series of cnidocyte assays were conducted that differed with amino acids and concentration.

The cnidocyte assay and statistical data accounted for a) probes that had at least one discharged cnidocyte attached b) probes that did not result in hydra attachment post probing c) probes that resulted in hydra contraction post touching the tentacles and d) probes that only touched one tentacle in the process of probing in both the control, Proline, and Glycine solutions. A Fisher's Test, a two tailed T- Test, a Mann- Whitney test, and a Chi- squared test were run for each group, the control and amino acid as well as a comparison of differing amino acid concentrations, to determine if a difference in interaction amounted. A P- Value of less than 0.05 for all tests was considered significant.

Data Collection of Proline

Of the 630 probes made, 458 probes were mounted and 294 probes were counted representing about a 46% utilization rate which is further shown in Table 1. The preliminary results suggested that the optimal concentration of Proline to encourage cnidocyte release was

10mM while 1mM and 100mM matched values of the control test indicating a potential to act as an adverse reactant. The average cnidocyte discharge was highest among amino acids with concentrations of 10mM as a compared to the control group. This was later concurred with statistical data such as the Fisher test (P- Value= 0.0003), referenced in Table 2, Mann- Whitney Test (P- Value = 0.0047), the two tailed T- Test (P- Value= .011), and finally the Chi- square test (P- Value= .000203). This was compared to the 1mM and 100mM and control experiments all of which were not significant (Table 1, Appendix).

Table 1: Statistical Analysis of Proline

	Proline	Glycine	1mM	10mM	100mM	10mM
	Control	Control	Proline	Proline	Proline	Glycine
Probes Made	126	96	72	72	72	192
Probes Counted	63	58	33	39	39	101

Table 1: Probe Count. Approximately 46% of probes made were counting and used for the statistical data.

Table 2: Fishers Test of Proline

Control to 1mM Proline			
	Probes	Cells	Totals
Control	63	304	367
1mM Proline	33	168	201
Total	96	472	568
P- Value	0.9069		
Control to the 10mM Proline			
	Probes	Cells	Totals
Control	63	304	367
10mM Proline	39	416	455
Total	102	720	822
P- Value	0.0003		
Control to 10mM Proline Excluding 103 Outlier			
	Probes	Cells	Totals
Control	63	304	367
10mM Proline w/o 103	38	313	351
Total	101	617	718
P- Value	0.0179		
Control to 100mM Proline			
	Probes	Cells	Totals
Control	63	304	367
100mM Proline	39	188	227
Total	102	492	594
P- Value	1.0		

Table 2: Fishers Test statistical analysis of varying concentrations of Proline as compared to the control. A concentration of 10mM was found to be significant as compared to the control in aggregating a cnidocyte response.

Data Collection of Glycine

Using this data, a second series of tests were performed using the optimal concentration of 10mM with the amino acid Glycine to further support the hypothesis that a) solubilized amino acids in gelatin delivered a significant effect in eliciting feeding responses as compared to a control group consisting of gelatin- coated probes and b) 10mM acted as the optimal concentration of various amino acids to enact discharge when compared to 1mM and 100mM which may deliver adverse effects. As compared to the control group, 10mM of Glycine elicited a significant positive response to feeding mechanism as shown by the amount of cnidocytes discharged and attached to the gelatin- coated probes which can be referenced in Figure 1. Statistical analysis confirmed that the probes coated with solubilized Glycine in gelatin as determined by the Fisher test (P- Value= 0.0011), Mann- Whitney Test (P- Value = 0.0), the two tailed T- Test (P- Value= .00028), and finally the Chi- square test (P- Value= .000724). A full data analysis of Glycine at a concentration of 10mM versus the control can be referenced in Table 3.




Table 3: Statistical Analysis of Glycine

Fishers Test			
Control to 10mM Glycine			
	Probes	Cells	Totals
Control	58	386	444
10mM Proline	103	1206	1307
Total	159	1592	1751
P- Value	0.0011		
Mann- Whitney Test			
Z- Score	4.061		
P- Value	0.0		
Not/ Significant	Significant		
Student T- Test			
T- Value	3.717		
P- Value	0.0003		
Not/Significant	Significant		
Chi- Squared Test			
Chi- Squared	11.427		
P- Value	0.0007		
Not/Significant	Significant		

Table 3: Statistical analysis of Glycine. A concentration of 10mM was found to be significant as compared to the control in aggregating a cnidocyte response.

Statistical Comparisons

Further statistical analysis's using the Fishers Test, Mann- Whitney Test, T-Test and Chi Square Test were conducted to compare a) different concentrations of amino acids to one another b) day to day variances between the control and the Proline or Glycine experimental c) day to day variances between amino acids of the same concentrations of amino acids.

When comparing the samples of 1mM Proline to 10mM Proline and 10mM Proline to 100mM Proline, the results proved to show significance with a P- Value of 0.0043 and 0.0013 respectively while the 1mM Proline to the 100mM Proline was not found to show a significant difference in cnidocyte discharge with a P- Value of 0.90. The Mann- Whitney Test and Chi-Squared Test also showed significance with comparison of 1mM Proline to 10mM Proline with P- Values of 0.038 and 0.0030 respectively while the T- test and Chi Squared Test showed significance with 10mM Proline to 100mM Proline outputting P- Values of 0.035 and 0.0009 respectively. Results can be referenced in Table 4 for the Fisher test and Table 2 of the Appendix for supplemental data.

Table 4: Fishers Test Proline Concentrations

1mM Proline to 10mM Proline			
	Probes	Cells	Totals
1mM Proline	33	168	201
10mM Proline	39	416	455
Total	72	584	656
P- Value	0.0043		
1mM Proline to 10mM Proline Excluding 103 Outlier			
	Probes	Cells	Totals
1mM Proline	63	304	367
10mM Proline w/o 103	38	33	351
Total	71	48	552
P- Value	0.0649		
1mM Proline to 100mM Proline			
	Probes	Cells	Totals
1mM Proline	33	168	201
100mM Proline	39	188	277
Total	72	356	428
P- Value	0.0897		
10mM Proline to 100mM Proline			
	Probes	Cells	Totals
10mM Proline	39	416	455
100mM Proline	39	188	227
Total	78	604	682
P- Value	0.0013		
10mM Proline Excluding 103 Outlier to 100mM Proline			
	Probes	Cells	Totals
10mM Proline w/o 103	38	313	351
100mM Proline	39	188	227
Total	77	501	578
P- Value	0.033		

Table 4: Comparative statistical analysis of differing Proline concentrations using the Fisher Test. The concentration of 10mM was found to be significantly different when compared to the 1mM and 100mM concentrations while 1mM and 100mM were not significantly different to one another showing that 10mM of Proline resulted in greater cnidocyte response.

The day to day comparison was between the control and the same amino acid concentration sample as well as with the comparison between differing concentrated samples with the Fishers Test per day to show that the standardization throughout each test and date were maintained. With the exception of the first day, November 12, 2014 which had an outlier, all other days were found to be not significant for Proline indicating that the procedure and day to day testing of the *Hydra* did not adversely impact the results. Results are depicted in Table 3 of the Appendix. Glycine was conducted in a similar manner with the focus on the day to day comparisons of the control to amino acid samples and the comparison of the concentrated samples to each other. These results are seen in Table 4 of the Appendix

Discussion

Cnidocyte discharge is dictated by a complex sensory control facilitated by the opsin pathway which is hypothesized to be similar to the systems organized in vertebrates. The behavioral experiments conducted in this study demonstrated that the amino acid cues, specifically Glycine and Proline, were capable of inducing cnidocyte response. This discharge elicits a feeding response that mimics the taste one receptor responses of vertebrates to umami indicating that the feeding responses in the vertebrate systems are similar to those of invertebrates.

The optimal concentration for cnidocyte discharge was found to be at 10mM as denoted in the experiments with differing concentrations of 1mM and 100mM of Proline resulting in a P-Value of 0.003 as compared to 0.907 and 1.0 respectively. The 1mM and 100mM concentrations of Proline varied little from the discharge rates of the control possibly suggesting that those concentrations could be adversely related to the feeding mechanisms. Glycine experiments were only conducted with 10mM as a concentration and showed relative to the control, a P-Value of 0.0011 proving enhanced cnidocyte discharge as compared to the control group.

The discharge rates could have been altered on occasion due to a variety of factors including cnidocyte counting, attachment, and daily operations. The hydra were gathered each experimental day around a similar time, however, some hydra may have been exposed to different light conditions which would have induced a more phototaxis release of cnidocytes. Cnidocyte discharge is regulated by many factors including phototaxis, feeding mechanisms and locomotion. These studies did not differentiate between the different cnidocytes that were released indicating that some of the cnidocytes discharge could be attributed to causes other than

feeding responses and thus a not in relation to T1Rs. Gelatin coated probes were left to dry for approximately two days, however, depending on the thickness of the coat, drying time may have varied which would have affected the attachment of the cnidocytes to the probes.

Overall, the data shows that amino acids elicit the same feeding response in systems of vertebrates and supplemental to other data, now predates the current understanding of the evolution of T1R pathways by about 400 million years. This dating indicates that T1R could have been the most ancient sensory receptor, present in the last common ancestor of animals and could have been present in the diversification of most animals unless lost.

References

1. Anderson, Peter A.v., and Christelle Bouchard. "The Regulation of Cnidocyte Discharge." *Toxicon* 54 (2008): 1046-053.
2. Bachmanov, Alexander A., and Gary K. Beauchamp. "Taste Receptor Genes." *Annual Review of Nutrition* 27 (2007): 389-414.
3. Cannon, Quinn, and Eric Wagner. "Comparison of Discharge Mechanisms of Cnidarian Cnidae and Myxozoan Polar Capsules." *Reviews in Fisheries Science* 11.3 (2003): 185-219.
4. Cliffe, E. E., and S. G. Waley. "Effect of Analogues Of Glutathione On The Feeding Reaction Of Hydra." *Nature* (1958): 804-05.
5. Giati, F. "Investigations on the G-protein Involved in Cnidarian Phototransduction." *Marine Biology* (2011): 1-58.
6. Kulkarni, R., Galande, S. Measuring Glutathione-induced Feeding Response in Hydra. *J. Vis. Exp.* (93), e52178, doi:10.3791/52178 (2014).
7. Lenhoff, H. M. "Activation of the Feeding Reflex in Hydra Littoralis: I. Role Played by Reduced Glutathione, and Quantitative Assay of the Feeding Reflex." *The Journal of General Physiology* 45.2 (1961): 331-44.
8. Lindemann, Bernd. "Receptors and Transduction in Taste." *Nature* 413 (2001): 219-25.
9. Lindstedt, K.june. "Chemical Control of Feeding Behavior." *Comparative Biochemistry and Physiology Part A: Physiology* 39 (1970): 553-81.
10. Muroi, Yoshikage, and Toshiaki Ishii. "Umami Taste Receptor Functions as an Amino Acid Sensor via Gαs Subunit in N1E-115 Neuroblastoma Cells." *Journal of Cellular Biochemistry* 113.5 (2012): 1654-662.

11. Nelson, Greg, Jayaram Chandrashekar, Mark A. Hoon, Luxin Feng, Grace Zhao, Nicholas J. P. Ryba, and Charles S. Zuker. "An Amino-acid Taste Receptor." *Nature* 416 (2002): 199-202.
12. Plachetzki, David C, Caitlin R Fong, and Todd H Oakley. "Cnidocyte Discharge Is Regulated by Light and Opsin-mediated Phototransduction." *BMC Biology* 10.17 (2012): 17.
13. Plachetzki DC, Fong CR, Oakley TH. "The Evolution of Phototransduction from an Ancestral Cyclic Nucleotide Gated Pathway." *Proceedings: Biological Sciences*: 277 (2010): 1963-1969.
14. Plachetzki, David C., Bernard M. Degnan, Todd H. Oakley, and Berend Snel. "The Origins of Novel Protein Interactions during Animal Opsin Evolution." *PLoS ONE* 2.10 (2007): 1054.
15. Petrocellis, L. De, D. Melck, T. Bisogno, A. Milone, and V. Di Marzo. "Finding of the Endocannabinoid Signalling System in Hydra, a Very Primitive Organism: Possible Role in the Feeding Response." *Neuroscience* 92.1 (1999): 377-87.
16. Purcell, J.E, and P.A.V Anderson. "Electrical Response to Water- Soluble Components of Fish Mucus Recorded From the Cnidocytes of a Fish Predator, *Physalia physalis*." *Marine and Freshwater Behaviour and Physiology* 354.1 (1995): 101-10.
17. Purves, Dale. "Taste Receptors and the Transduction of Taste Signals." *Neuroscience*. 2nd ed. Sunderland, Massachusetts.: Sinauer Associates, 2012.
18. Shirazi- Beechy, SP, K. Daly, M. Al-Rammahi, and AW Moran. "Role of Nutrient-sensing Taste 1 Receptor (T1R) Family Members in Gastrointestinal Chemosensing." *British Journal of Nutrition* 111 (2013): 8-15.

19. Thorington GU, Hessinger DA: Control of discharge: factors affecting discharge of cnidae. In *The Biology of Nematocysts*. Edited by Hessinger DA, Lenhoff HM. San Diego, CA: Academic Press; 1988:233-254.
20. Watson, Glen M., and David A. Hessinger. "Antagonistic Frequency Tuning of Hair Bundles by Different Chemoreceptors Regulates Nematocyst Discharge." *The Journal of Experimental Biology* 187 (1993): 57-73.
21. Zhang, Yifeng, Mark A. Hoon, Jayaram Chandrashekar, Ken L. Mueller, Boaz Cook, Dianqing Wu, Charles S. Zuker, and Nicholas J.p. Ryba. "Coding of Sweet, Bitter, and Umami Tastes." *Cell* 112.3 (2003): 293-301.

APPENDIX

Table 1: Statistical Analysis of Proline

Mann- Whitney Test				
	1mM Proline	10mM Proline	10mM Excluding 103 Proline	100mM Proline
Z- Score	-0.726	-2.827	-2.657	-1.215
P- Value	0.465	0.005	0.008	0.222
Not/Significant	Not Significant	Significant	Significant	Not Significant
Student T- Test				
	1mM Proline	10mM Proline	10mM Excluding 103 Proline	100mM Proline
T- Value	0.239	2.586	2.778	0.005
P- Value	0.811	0.011	0.007	0.005
Not/ Significant	Not Significant	Significant	Significant	Not Significant
Chi- Squared Test				
	1mM Proline	10mM Proline	10mM Excluding 103 Proline	100mM Proline
Chi- Squared	0.052	13.807	5.966	0.0
P- Value	0.820	0.0002	0.014	0.996
Not/Significant	Not Significant	Significant	Significant	Not Significant

Table 1: Statistical analysis of varying concentrations of Proline as compared to the control. A concentration of 10mM was found to be significant as compared to the control and other concentrations in aggregating a cnidocyte response.

Table 2: Statistical Analysis of Proline Concentrations

Mann- Whitney Test					
	1mM v .10mM	1mM v. 10mM w/o	1mM v. 100mM	10mM v. 100mM	10mM w.o 103 v
	Proline	103 Proline	Proline	Proline	100mM Proline
Z- Score	2.0795	1.931	0.5933	1.839	-1.676
P- Value	0.0375	0.00536	0.5552	0.0658	0.0930
Not/Significant	Significant	Significant	Not Significant	Not Significant	Not Significant
Student T- Test					
	1mM v .10mM	1mM v. 10mM w/o	1mM v. 100mM	10mM v. 100mM	10mM w.o 103 v
	Proline	103 Proline	Proline	Proline	100mM Proline
T- Value	1.8433	2.114	0.2747	2.146	2.739
P- Value	0.0696	0.0382	0.7844	0.035	0.0077
Not/ Significant	Not Significant	Significant	Not Significant	Significant	Significant
Chi- Squared Test					
	1mM v .10mM	1mM v. 10mM w/o	1mM v. 100mM	10mM v. 100mM	10mM w.o 103 v
	Proline	103 Proline	Proline	Proline	100mM Proline
Chi- Squared	8.785	5.966	5.966	11.082	4.8204
P- Value	0.003	0.05899	0.833	0.0009	0.0281
Not/Significant	Significant	Not Significant	Not Significant	Significant	Significant

Table 2: Statistical analysis of varying concentrations of Proline as compared other Proline concentrations. Tests were found to be significant for for comparison between 1mM and 10mM and 10mM and 100mM while comparisons of 1mM to 100mM were not significant indicating that the 10mM concentration of Proline elicited the greatest cnidocyte discharges as compared to other concentrations.

Statistical Analysis of Proline Daily Fishers Test of Day to Day Comparison															
11/12/2014				11/14/2014				11/19/2014							
CONTROL TO PROLINE 10mM				CONTROL TO PROLINE 10mM				CONTROL TO PROLINE 100X							
	PROBES	CELLS	TOTALS		PROBE	CELLS	TOTALS		PROBES	CELLS	TOTALS				
CONTROL	11	53	64	CONTROL	12	74	86	CONTROL	8	32	40				
PROLINE 10mM	9	186	195	PROLINE 10mM	11	86	97	PROLINE 100mM	8	28	36				
TOTAL	20	239	259	TOTAL	23	160	183	TOTAL	16	60	76				
P VALUE	0.0024			P VALUE	0.6584			P VALUE	1						
CONTROL TO PROLINE 10mM W/O 103															
	PROBES	CELLS	TOTALS												
CONTROL	11	53	64												
PROLINE 10mM	8	83	91												
TOTAL	19		155												
P VALUE	0.139														
11/21/2014				11/23/2014				11/25/2014				12/3/2014			
CONTROL TO PROLINE 10mM				CONTROL TO PROLINE 10mM				CONTROL TO PROLINE 1mM				CONTROL TO PROLINE 1mM			
	PROBES	CELLS	TOTALS		PROBE	CELLS	TOTALS		PROBES	CELLS	TOTALS				
CONTROL	7	35	42	CONTROL	10	56	66	CONTROL	8	28	36	CONTROL	7	26	33
PROLINE 10mM	11	82	93	PROLINE 10mM	8	62	70	PROLINE 1mM	16	56	72	PROLINE 1mM	10	33	43
TOTAL	18	117	135	TOTAL	18	118	136	TOTAL	16	56	72	TOTAL	17	59	76
P VALUE	0.4281			P VALUE	0.6159			P VALUE	1			P VALUE	1		
CONTROL TO PROLINE 100mM				CONTROL TO PROLINE 100mM				CONTROL TO PROLINE 1mM				CONTROL TO PROLINE 100mM			
	PROBES	CELLS	TOTALS		PROBE	CELLS	TOTALS		PROBES	CELLS	TOTALS				
CONTROL	7	35	42	CONTROL	10	56	66	CONTROL	8	28	36	CONTROL	7	26	33
PROLINE 100mM	12	74	86	PROLINE 100mM	9	66	71	PROLINE 1mM	8	67	75	PROLINE 100mM	10	24	34
TOTAL	19	109	128	TOTAL	19	118	137	TOTAL	16	95	111	TOTAL	17	50	67
P VALUE	0.792			P VALUE	0.8057			P VALUE	0.1477			P VALUE	0.5763		
PROLINE 10 mM to PROLINE 100mM				PROLINE 10 mM to PROLINE 100mM				CONTROL TO PROLINE 1mM				PROLINE 1 mM to PROLINE 100mM			
	PROBES	CELLS	TOTALS		PROBE	CELLS	TOTALS		PROBES	CELLS	TOTALS				
PROLINE 10mM	11	82	93	PROLINE 10mM	9	62	71	CONTROL	8	28	36	PROLINE 1mM	10	33	43
PROLINE 100mM	12	74	86	PROLINE 100mM	8	62	70	PROLINE 1mM	7	40	47	PROLINE 100mM	10	24	34
TOTAL	23	156	179	TOTAL	17	124	141	TOTAL	15	68	83	TOTAL	20	57	77
P VALUE	0.8236			P VALUE	1			P VALUE	0.4056			P VALUE	0.606		
Amino Acid to Amino Acid															
11/25/2014															
PROLINE 1 mM A to Proline 1mMB				PROLINE 1 mM A to Proline 1mMC				PROLINE 1 mM B to Proline 1mMC							
	PROBES	CELLS	TOTALS		PROBE	CELLS	TOTALS		PROBES	CELLS	TOTALS				
PROLINE 1mM	8	28	36	PROLINE 1mM	8	28	36	PROLINE 1mM	8	67	75				
PROLINE 1mM	8	67	75	PROLINE 1mM	7	40	47	PROLINE 1mM	7	40	47				
TOTAL	16	95	111	TOTAL	15	68	83	TOTAL	15	107	122				
P VALUE	0.1477			P VALUE	0.41			P VALUE	0.5744						

Table 3: The day to day comparison between the Control and the Proline at 1mM, 10mM, and 100mM were tested with the Fisher tests as well as analysis between the day to day analysis of the same concentration being tested on the same day as seen on 11/25/2014. With the exception, November 12, 2014, all other days were found to be not significant for Proline indicating that the procedure and day to day testing of the *Hydra* did not adversely impact the results.

Statistical Analysis of Glycine Daily Fishers Test of Day to Day Comparison															
1/20/2015				1/23/2015				1/30/2015				2/4/2015			
CONTROL TO 10mM GLYCINE A				CONTROL TO 10mM GLYCINE				CONTROL TO 10mM GLYCINE				CONTROL TO 10mM GLYCINE			
	PROBES	CELLS	TOTALS		PROBES	CELLS	TOTALS		PROBES	CELLS	TOTALS		PROBES	CELLS	TOTALS
CONTROL	16	179	195	CONTROL	16	91	107	CONTROL	12	48	60	CONTROL	14	68	82
GLYCINE 10mM	14	198	212	GLYCINE 10mM	9	121	130	GLYCINE 10mM	15	98	113	GLYCINE 10mM	16	120	136
TOTAL	30	377	407	TOTAL	25	212	237	TOTAL	27	146	173	TOTAL	30	188	218
P VALUE	0.573105			P VALUE	0.056003			P VALUE	0.274722			P VALUE	0.312042		
CONTROL TO 10mM GLYCINE B				CONTROL TO 10mM GLYCINE				CONTROL TO 10mM GLYCINE				CONTROL TO 10mM GLYCINE			
CONTROL	16	179	195	CONTROL	16	91	107	CONTROL	12	48	60	CONTROL	14	68	82
GLYCINE 10mM	7	179	186	GLYCINE 10mM	15	183	198	GLYCINE 10mM	10	74	84	GLYCINE 10mM	15	233	248
TOTAL	23	358	381	TOTAL	31	274	305	TOTAL	22	122	144	TOTAL	29	301	330
P VALUE	0.085391			P VALUE	0.048322			P VALUE	0.240547			P VALUE	0.00552		
TOTALS				TOTALS				TOTALS				TOTALS			
CONTROL TO 10mM GLYCINE				CONTROL TO 10mM GLYCINE				CONTROL TO 10mM GLYCINE				CONTROL TO 10mM GLYCINE			
CONTROL	16	179	195	CONTROL	16	91	107	CONTROL	12	48	60	CONTROL	14	68	82
GLYCINE 10mM	21	377	398	GLYCINE 10mM	24	304	328	GLYCINE 10mM	25	172	197	GLYCINE 10mM	31	353	384
TOTAL	37	556	593	TOTAL	40	395	435	TOTAL	37	220	257	TOTAL	45	421	466
P VALUE	0.205072			P VALUE	0.032131			P VALUE	0.205932			P VALUE	0.021354		
Amino Acid to Amino Acid 10mM A GLYCINE to 10mM B				Amino Acid to Amino Acid 10mM A GLYCINE to 10mM B				Amino Acid to Amino Acid 10mM A GLYCINE to 10mM B				Amino Acid to Amino Acid 10mM A GLYCINE to 10mM B			
GLYCINE 10mM A	14	198	212	GLYCINE 10mM A	9	121	130	GLYCINE 10mM A	15	98	113	GLYCINE 10mM A	16	120	136
GLYCINE 10mM B	7	179	186	GLYCINE 10mM B	15	183	198	GLYCINE 10mM B	10	74	84	GLYCINE 10mM B	15	233	248
TOTAL	21	377	398	TOTAL	24	304	328	TOTAL	25	172	197	TOTAL	31	353	384
P VALUE	0.2629			P VALUE	1			P VALUE	0.8315			P VALUE	0.0761		

Table43: The day to day comparison was between the control and the Glycine 10mM concentration. Most days except 2/4/2015 show non-significant data analysis indicating that the day to day variances were minimal. Glycine to the control was found to be significant on day of 1/23/2015 and 2/4/2015.

The flavor of life: evolutionarily ancient sensory receptors for photosensitivity and taste in the cnidarian, *Hydra magnipapillata*

Connor Bell¹, Molly Hartley¹, Janine R. Appleton¹, Hannah Gianville, Julie Peterson and David Plachetzki²



Abstract
Animals sense their environment using specialized proteins called sensory receptors. We use the cnidarian freshwater polyp *Hydra magnipapillata* as a model to understand the early evolution of animal sensory systems. Ongoing work has identified a set of sensory genes that function in the hydra including the photosensitivity gene *Opsin* and Taste 1 Receptors (*T1Rs*). Here we report results from molecular cloning and *in situ* hybridization that demonstrates the dual expression of the photoreceptor gene *Opsin* and the putatively chemoreceptive gene *T1R3* in a polymodal sensory cell type in the hydra. In addition, we report results from behavioral trials that demonstrate that chemical amino acid cues from the environment are capable of exciting the discharge behavior of stinging cells called cnidocytes, mirroring the vertebrate taste sense of umami.

Background
We study the cnidarian animal *Hydra magnipapillata*, a relative of corals, jellyfish and anemones, as a model for understanding the early evolutionary histories of the animal senses

Previous work has demonstrated the importance of photosensitive opsin in mediating cnidarian feeding behavior (Plachetzki et al 2012).

Taste 1 Receptors (*T1Rs*) are responsible for umami and sweet taste sensation, but are presently only known from mammals (Shi and Zhang 2006)

Amino acids such as proline and glycine are capable of inducing *T1R* signaling, which humans perceive as savory taste

Whole genome data for representative animal genomes allow the origins of *T1Rs* to be explored

Results
Phylogenetic analyses of *T1R* genes suggest that they were present in the last common ancestor of animals and comprise one of the most ancient animal senses.

T1R genes are expressed in a polymodal sensory neuron in the hydra that also expresses photosensitive *Opsins*.

T1Rs may function in feeding behavior by modulating the cnidocyte (stinging cell) discharge

References
Plachetzki DC, Degnan BM, Oakley TH. The origins of novel protein interactions during animal opsin evolution. *PLoS One*. 2007 Oct 17;2(10):e1054. | Borowiec M, Lee E, Chiu J, and Plachetzki D. 2015. Dissecting phylogenetic signal and species bias for bias-free genome-wide data. *PLoS One*. 2012. | C. Opsin staining (green) in whole mounts. **B.** *T1R* and *Opsin* co-localize in the same cell type. **C.** Opsin staining (green) in whole mounts. **D.** *T1R* and *Opsin* co-localize in the same cell type. **E.** Confocal microscopy reveals that cnidocyte/sensory cells express both *Opsin* and *T1R*.




Figure 1. A. An emerging view of animal phylogeny drawn from exclusively whole genome sequences. This tree represents our current bioinformatics pipeline from which the phylogenetic history of animal *T1Rs* (B) was inferred. The animal phylogeny shown in A is described elsewhere (Borowiec et al 2015). For B, sequences of known mammalian *T1R* receptors were used as queries for extensive searches genomes depicted in A. Phylogenetic analyses were conducted using the best fitting model in RAxML (Stamatakis 2014).




Figure 2. Sequences of known mammalian taste receptors were used to searched genomic data for *H. magnipapillata* and isolated putative *T1R* gene sequences. We designed four gene-specific oligonucleotide primers for use in RACE and amplification of cDNA ends (RACE). Two kit-specific primers were also used, which bind specifically to synthetic adaptors that were ligated to either the 5' or the 3' end of the transcript. From there, Sanger sequencing was performed on the resulting clones and the data were assembled using GENIOUS, resulting in the full length *T1R3* clone.




Figure 3. A. Light micrograph of living specimen of *H. magnipapillata*. **B-E.** *In situ* hybridization of sensory genes in the hydra. **B.** *T1R* staining (red) in whole mount. **C.** *Opsin* staining (green) in whole mount. **D.** *T1R* and *Opsin* co-localize in the same cell type. **E.** Confocal microscopy reveals that cnidocyte/sensory cells express both *Opsin* and *T1R*.




Figure 4. The effects of amino acid cues on cnidocyte activity were measured using a cnidocyte discharge assay (Plachetzki et al 2012). Discharged cnidocytes were captured using control and amino acid containing gelatin probes and analyzed using light microscopy under DIC. Cnidocyte discharge is significantly enhanced on probes containing 10mM glycine and/or proline. This experiment shows that amino acids elicit a feeding response similar in the hydra that we hypothesize is mediated by hydra *T1Rs*.

Conclusions
T1Rs were present in the last common ancestor of animals and function in the feeding behavior of early branching animal lineages

This finding predates the current understanding of taste receptor evolution by ~ 400 million years, indicating a possible role of *T1Rs* during the diversification of animals

The co-localization of chemosensitivity (taste) and photosensitivity (vision) genes in neural cells of the hydra suggests new hypotheses for the origins of animals sensory neurons

The cnidarian polymodal sensory neuron identified here is a useful model for exploring the early evolutionary history of the animal senses.

Figure 3. A. Light micrograph of living specimen of *H. magnipapillata*. **B-E.** *In situ* hybridization of sensory genes in the hydra. **B.** *T1R* staining (red) in whole mount. **C.** *Opsin* staining (green) in whole mount. **D.** *T1R* and *Opsin* co-localize in the same cell type. **E.** Confocal microscopy reveals that cnidocyte/sensory cells express both *Opsin* and *T1R*.