

University of New Hampshire

University of New Hampshire Scholars' Repository

Center for Coastal and Ocean Mapping

Center for Coastal and Ocean Mapping

5-2015

Developing methodology for efficient eelgrass habitat mapping across lidar systems

Victoria Price

University of New Hampshire, Durham

Jennifer A. Dijkstra

University of New Hampshire, Durham, Jennifer.Dijkstra@unh.edu

Jarlath O'Neil-Dunne

University of Vermont

Christopher Parrish

Oregon State University

Erin Nagal

University of New Hampshire, Durham

See next page for additional authors

Follow this and additional works at: <https://scholars.unh.edu/ccom>

Recommended Citation

V. E. Price, Dijkstra, J. A., O'Neil-Dunne, J. P. M., Parrish, C., Nagle, E., and Pe'eri, S., "Developing methodology for efficient eelgrass habitat mapping across lidar systems", *Marine Geological and Biological Habitat Mapping (GEOHAB)*. Salvador, Brazil, 2015.

This Conference Proceeding is brought to you for free and open access by the Center for Coastal and Ocean Mapping at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Center for Coastal and Ocean Mapping by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

Authors

Victoria Price, Jennifer A. Dijkstra, Jarlath O'Neil-Dunne, Christopher Parrish, Erin Nagal, and Shachak Pe'eri

Developing Methodology for Efficient Eelgrass Habitat Mapping Across Lidar Systems.

**Victoria E. Price¹ , Jennifer A. Dijkstra¹ , Jarlath P.M. O'Neil-Dunne² ,
Christopher E. Parrish³ , Erin Nagel¹ , Shachak Pe'eri¹**

**1. University of New Hampshire Center for Coastal and Ocean Mapping,
Durham, NH USA 03824**

2. University of Vermont, Burlington VT USA 05405

3. Oregon State University, Corvallis OR USA 97331

Super Storm Sandy, the second costliest hurricane in U.S. history, made landfall on the east coast of the U.S. in October 2012. In an attempt to assess the impacts of the storm on coastal ecosystems, several U.S. mapping agencies such as the National Oceanic and Atmospheric Administration (NOAA), the U.S. Geological Survey (USGS), and the U.S. Army Corps of Engineers (USACE) commenced data collection efforts using a variety of remotely-sensed data types including aerial imagery and topobathymetric lidar. The objective of this study was to investigate the applicability of object-based image analysis techniques for benthic habitat mapping. Bathymetry and reflectance data collected by a Riegl VQ-820-G system and the AHAB Chiroptera system along with aerial imagery (Applanix DSS) were compared using an objectbased image analysis (OBIA) technique to classify dense eelgrass beds, mixed sand and macroalgae, and sand habitats. In order to determine the efficacy of this method for benthic habitat classification it was also compared to a manual method of classification from aerial imagery. The resulting habitat maps were compared between systems to determine the feasibility of using one OBIA classification rule set across lidar systems and aerial imagery. Our preliminary results using the Riegl system suggest our methodology correctly classified 85% of benthic habitats. Preliminary results using the Chiroptera also suggests similar accuracy of classification. This methodology will allow streamlined creation of habitat maps for coastal managers and researchers using large sets of data collected by multiple sensors. Testing of this OBIA methodology is ongoing as new data from various sensors becomes available.