

University of New Hampshire

University of New Hampshire Scholars' Repository

Life Sciences Faculty Scholarship

Life Sciences

9-1-2017

Draft Genome Sequence of the Marine Bacterium *Oceanimonas baumannii* ATCC 700832T

William D. McClelland
University of New Hampshire, Manchester

Ariel M. Trachtenberg
University of New Hampshire, Manchester

Marc A. Brennan
University of New Hampshire, Manchester

Kyle S. MacLea
University of New Hampshire, Manchester, kyle.maclea@unh.edu

Follow this and additional works at: https://scholars.unh.edu/unhmbiology_facpub

Recommended Citation

McClelland, W.D., Trachtenberg, A.M., Brennan, M.A., and MacLea, K.S. Draft Genome Sequence of the Marine Bacterium *Oceanimonas baumannii* ATCC 700832T. *Genome Announc*, 5:e01007-17; doi:10.1128/genomeA.01007-17, 2017.

This Article is brought to you for free and open access by the Life Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Life Sciences Faculty Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.



Draft Genome Sequence of the Marine Bacterium *Oceanimonas baumannii* ATCC 700832^T

William D. McClelland,^a Ariel M. Trachtenberg,^a Marc A. Brennan,^b
Kyle S. MacLea^{a,b,c}

Biology Program, University of New Hampshire, Manchester, New Hampshire, USA^a; Biotechnology Program, University of New Hampshire, Manchester, New Hampshire, USA^b; Department of Life Sciences, University of New Hampshire, Manchester, New Hampshire, USA^c

ABSTRACT The aerobic phenol-degrading Gram-negative rod *Oceanimonas baumannii* ATCC 700832^T was first isolated from estuary mud from the River Wear, United Kingdom, in 1983. Information on the draft genome sequence for *O. baumannii* ATCC 700832^T is included in this announcement. The predicted genome size is 3,809,332 bp, with 55.88% G+C content.

Oceanimonas baumannii strain GB6^T (=ATCC 700832^T) is an aerobic Gram-negative rod that displays up to four polar flagella and can readily grow on minimal salts medium supplemented with phenol (1). It was isolated in 1998 from estuary mud at the mouth of the River Wear in Sunderland, United Kingdom (1). At the time of its discovery, Brown et al. found that *O. baumannii* should be classified in a new genus called *Oceanomonas* (1), later corrected in spelling to *Oceanimonas* (2), along with the type strain for this new genus, (*Pseudomonas*) *doudoroffii* ATCC 27123, now called *Oceanimonas doudoroffii* (1).

O. baumannii is a chemoorganotrophic gammaproteobacterium and has a requirement for sodium. It can grow at NaCl levels of up to 7% (wt/vol) (1) or 12% (3). Its absolute requirement for NaCl is 0.15% (wt/vol) (3), and it grows at temperatures between 10°C and 30°C (1). *O. baumannii* can be grown on glycerol, D-galactose, caprate, malate, citrate, ethanol, betaine, sarcosine, phenol, succinate, L-alanine, L-proline, and L-glutamate (1).

O. baumannii ATCC 700832^T was purchased from the ATCC (Manassas, VA, USA) in lyophilized form. It was rehydrated and then cultured in marine broth or agar (ATCC medium 2216) under conditions of 26°C for 48 h at atmospheric pressure. A single colony was picked and cultured in log-phase growth to obtain genomic DNA (gDNA), using the Genomic-tip 500/G kit (Qiagen, Valencia, CA, USA). gDNA was fragmented, tagged with adapters using the Nextera DNA library prep kit (Illumina, San Diego, CA, USA), and sequenced on an Illumina HiSeq 2500. Two hundred fifty-base-pair paired-end reads were generated at the Hubbard Center for Genome Studies (Durham, NH, USA), and Trimmomatic (4) was used for bioinformatic trimming and removal of adapter sequences prior to gene analysis.

The *O. baumannii* genome was assembled using SPAdes version 3.8.0 (5) from 2,790,420 reads, with an average length of 899 bp, into 70 contigs (after removal of small low-quality contigs). The total genome length was 3,809,332 bp, and we calculated an average coverage of 80×. The largest contig found was 408,079 bp, with a G+C content of 55.88% and an N_{50} value of 232,731 bp. The G+C content is in close agreement with that published by Brown et al., at 54% (1). The genome was annotated with the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) process (6). PGAP found

Received 12 August 2017 Accepted 15 August 2017 Published 7 September 2017

Citation McClelland WD, Trachtenberg AM, Brennan MA, MacLea KS. 2017. Draft genome sequence of the marine bacterium *Oceanimonas baumannii* ATCC 700832^T. Genome Announc 5:e01007-17. <https://doi.org/10.1128/genomeA.01007-17>.

Copyright © 2017 McClelland et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kyle S. MacLea, kyle.maclea@unh.edu.

a total of 3,651 genes, 3,498 coding sequences (CDSs), 98 RNA genes, 55 pseudogenes, and 1 clustered regularly interspaced short palindromic repeat (CRISPR) array.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession no. [NQJF0000000](https://doi.org/10.1093/bioinformatics/btu170). The version described in this paper is version NQJF01000000.

ACKNOWLEDGMENTS

Sequencing and bioinformatics analyses were undertaken at the Hubbard Center for Genome Studies at UNH with the kind assistance of Kelley Thomas, Jordan Ramsdell, and Stephen Simpson. K.S.M. acknowledges Annabelle MacLea and Duncan MacLea for helpful discussions. This work was a project of the Microbiology Education through Genome Annotation–New Hampshire (MEGA-NH) program.

The Department of Life Sciences at UNH Manchester provided funds for sequencing. The funders had no role in the study design, data collection, or interpretation or the decision to submit the work for publication.

REFERENCES

1. Brown GR, Sutcliffe IC, Cummings SP. 2001. Reclassification of [*Pseudomonas*] *doudoroffii* (Baumann et al. 1983) into the genus *Oceanomonas* gen. nov. as *Oceanomonas doudoroffii* comb. nov., and description of a phenol-degrading bacterium from estuarine water as *Oceanomonas baumannii* sp. nov. *Int J Syst Evol Microbiol* 51:67–72. <https://doi.org/10.1099/00207713-51-1-67>.
2. International Journal of Systematic and Evolutionary Microbiology. 2001. Notification that new names and new combinations have appeared in volume 51, part 1, of the IJSEM. *Int J Syst Evol Microbiol* 51:269. <https://doi.org/10.1099/00207713-51-2-269>.
3. Brown GR, Sutcliffe IC, Bendell D, Cummings SP. 2000. The modification of the membrane of *Oceanimonas baumannii* when subjected to both osmotic and organic solvent stress. *FEMS Microbiol Lett* 189:149–154. <https://doi.org/10.1111/j.1574-6968.2000.tb09221.x>.
4. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
5. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
6. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciufu S, Li W. 2013. Prokaryotic Genome Annotation Pipeline. National Center for Biotechnology Information, Bethesda, MD.