

University of New Hampshire

## University of New Hampshire Scholars' Repository

---

Biological Sciences Scholarship

Biological Sciences

---

Spring 2015

### Estimating plasma volume in neonatal Holstein calves fed one or two feedings of a lacteal-based colostrum replacer using Evans blue dye and hematocrit values at various time points.

Peter S. Erickson

*University of New Hampshire, peter.erickson@unh.edu*

Rosemarie G. Cabral Ph.D.

*University of New Hampshire - Main Campus*

Colleen E. Chapman MS

*University of New Hampshire - Main Campus*

E. J. Kent

*University of New Hampshire*

Follow this and additional works at: [https://scholars.unh.edu/biosci\\_facpub](https://scholars.unh.edu/biosci_facpub)



Part of the [Dairy Science Commons](#)

---

#### Recommended Citation

Erickson, Peter S.; Cabral, Rosemarie G. Ph.D.; Chapman, Colleen E. MS; and Kent, E. J., "Estimating plasma volume in neonatal Holstein calves fed one or two feedings of a lacteal-based colostrum replacer using Evans blue dye and hematocrit values at various time points." (2015). *Canadian Journal of Animal Science*. 39.

[https://scholars.unh.edu/biosci\\_facpub/39](https://scholars.unh.edu/biosci_facpub/39)

This Article is brought to you for free and open access by the Biological Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Biological Sciences Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact [Scholarly.Communication@unh.edu](mailto:Scholarly.Communication@unh.edu).

1 **Estimating plasma volume in neonatal Holstein calves fed one or two feedings of a lacteal-**  
2 **based colostrum replacer using Evans blue dye and hematocrit values at various time**  
3 **points.**

4  
5  
6  
7 **Rosemarie G. Cabral<sup>1</sup>, Colleen E. Chapman, Emily J. Kent, and Peter S. Erickson\***  
8 *Department of Biological Sciences, University of New Hampshire, Durham, 03824, USA*

9  
10  
11 \*Corresponding author. Tel.: +1 603 862 1909:

12 Email address: [peter.erickson@unh.edu](mailto:peter.erickson@unh.edu)

13 30 O’Kane Road  
14 University of New Hampshire  
15 Durham, NH, 03824, USA

16  
17 <sup>1</sup>Present Address: Famo Feeds, Freeport, MN, USA

18  
19  
20  
21  
22  
23

24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37

## ABSTRACT

Twenty-eight Holstein calves were blocked by birth date and randomly assigned to one of two treatments to investigate the effect of colostrum replacer (**CR**) feeding regimen on plasma volume (**PV**). Treatments were: 1) one feeding of CR (**C<sub>1</sub>**; 3L of reconstituted CR 675 g of powder providing 184.5 g of IgG at birth) or 2) two feedings of CR (**C<sub>2</sub>**; 2L of reconstituted CR at birth and 1 L of reconstituted CR at six h). By 6 h of age, all calves had received 3L of CR providing 184.5 g of IgG. Plasma volume was estimated at six, 12, 18, and 24 h after birth using Evans blue dye (**EBD**). No treatment effects were noted at any time points ( $P > 0.05$ ). Mean PV for all calves regardless of treatment at six, 12, 18, and 24 h were 78.6, 89.2, 83.9, and 90.7 mL kg<sup>-1</sup> of BW, respectively. Plasma volume was correlated with hematocrit (**HCT**), initial HCT, and treatment. Hematocrit was correlated with PV, initial HCT, and body weight. Hematocrit for six, 12, 18 and 24 h after birth can be predicted with an initial precolostral HCT determination.

## RÉSUMÉ

Vingt-huit veaux de race Holstein ont été classés par leurs dates de naissance et soumis aléatoirement à l'un de deux traitements afin d'examiner l'effet d'un régime alimentaire de remplacement de colostrum (**RC**) sur le volume plasmatique (**VP**). Les traitements étaient : 1) Un dosage de RC (**C<sub>1</sub>**; 184,5 g de IgG à la naissance) ou 2) deux dosages de RC (**C<sub>2</sub>**; 2/3 de la dose totale à la naissance, 1/3 de la dose totale 6 heures plus tard). Le volume plasmatique a été estimé six, douze, dix-huit et vingt-quatre heures après la naissance. Le colorant bleu d'Evans (**CBE**) a été utilisé pour estimer le VP; l'effet du traitement n'a été observé à aucun moment sur le VP ( $P > 0,05$ ). Indépendamment du traitement, les moyennes de VP pour tous les veaux à six, douze, dix-huit et vingt-quatre heures étaient 78,6; 89,2; 83,9 et 90,7 mL kg<sup>-1</sup> de poids corporel

47 (PC), respectivement. Le VP a été corrélé à l'hématocrite (HtC), l'HtC initial et au traitement.  
48 L'hématocrite a été corrélé au VP, à l'HtC initial et au poids. L'hématocrite à six, douze, dix-  
49 huit et vingt-quatre heures après la naissance peut-être prédit par une détermination initiale de  
50 l'HtC avant l'ingestion de colostrum.

51

52 **Key words: calf, plasma volume, colostrum replacer, hematocrit**

53

54 **Abbreviations:** AEA, apparent efficiency of absorption; CR, colostrum replacer; EBD Evans  
55 Blue Dye; HCT hematocrit; I  $\mu\text{g}$  of dye injected; IgG immunoglobulin G; PV plasma volume; S  
56 absorbance of standard; U absorbance of unknown

57

58

59

60

61

62

63

64

65

66

67

68

69

## INTRODUCTION

70  
71 Apparent efficiency of absorption (**AEA**) is commonly calculated in neonatal calf research as  
72 another means of viewing Immunoglobulin G (**IgG**) absorptive data. The AEA can be estimated  
73 using the equation:  $[\text{plasma IgG (g/L)} \times \text{BW (kg)} \times 0.09/\text{IgG mass consumed (g)}] \times 100\%$   
74 (Quigley et al., 1998). Within the equation, the value 0.09 refers to PV, which has been measured  
75 in the calf at 24 and 48 h of age. Prior to the work of Quigley et al. (1998), Mollenberg et al.  
76 (1975) reported PV increasing from 53 mL (plasma)kg<sup>-1</sup> of BW at birth to 65 ml kg<sup>-1</sup> of BW at  
77 one day old after being fed 150 g of colostrum kg<sup>-1</sup> of BW in four feedings and McEwan et al.  
78 (1968) indicated that PV increased from 66 ml kg<sup>-1</sup> of BW at birth to 93 ml kg<sup>-1</sup> of BW at the end  
79 of d one after calves were fed colostrum. Quigley et al. (1998) attributed the discrepancies in PV  
80 values in the aforementioned studies to the small number of calves used. These data are also  
81 limited by PV being measured only at birth and 24 h after birth. Researchers feeding CR utilize  
82 the equation of Quigley et al. (1998). However, that equation was derived from calves fed  
83 colostrum. Most researchers studying colostrum feed calves 4L while adequate passive transfer  
84 can occur in calves fed 3 L of CR. Therefore, we question the appropriateness of using the  
85 equation of Quigley et al. (1998) for calves fed CR. An objective of this study was to provide  
86 PV of calves fed CR for the determination of AEA at different time points such as six, 12, and 18  
87 h after birth.. The second objective of this experiment was to determine if there were differences  
88 in PV due to feeding regimen of CR using EBD at six, 12, 18, and 24 h after birth in Holstein  
89 dairy calves. The third objective was to determine PV at different time points and to determine if  
90 there was any correlation between HCT values and PV determined by the EBD procedure. If  
91 HCT values are correlated with PV, then it is possible that PV could be calculated on a per calf

92 basis using HCT values at each time point. The last objective of this study was to determine if  
93 HCT could be predicted using precolostral HCT values.

## 94 **MATERIALS AND METHODS**

### 95 **Experimental Design and Treatment Diets**

96 This experiment was reviewed and approved by the University of New Hampshire  
97 Institutional Animal Care and Use Committee (# 111205 and 130105).

98 Twenty-eight Holstein calves were born from multiparous and primiparous cows and  
99 blocked by birth date. Calves were removed from dams within 30 min of birth, weighed and had  
100 their navels dipped with a 7 % tincture of iodine. Calves were individually housed for the  
101 duration of the experiment. All calves received 3 sachets of Calf's Choice Total™ Gold CR  
102 containing a total IgG mass of 184.5 g reconstituted, according to manufacturer's instructions to  
103 a final volume of 3L (The Saskatoon Colostrum Co., Saskatoon, SK). One sachet of product  
104 contains 225 g of powder. Therefore, all calves received 675 g of powder. Assuming 950 g kg<sup>-1</sup>  
105 (DM) each calf received 641 g of solids with a final volume of 3 L. Total solids equaled 210  
106 g/L. Calves were assigned to one of two treatments: 1) one feeding of CR (C<sub>1</sub>) or 2) two feedings  
107 of CR (C<sub>2</sub>). Calves fed a single feeding of CR received the total dose at 0 h (3L CR, 184.5 g  
108 IgG). Calves assigned to two feedings received 2/3 of the total CR dose at 0 h (2L CR, 123 g  
109 IgG) and the remaining 1/3 of CR at six h (1L CR, 61.5 g IgG). All calves had consumed 3L of  
110 CR by six h. Calves were fed via nipple bottle with the first feeding of CR given at 0 h (within  
111 30 min of birth). If the calf did not consume the entire volume of CR, it was administered via  
112 esophageal tube to ensure the total dose was received.

113

114

115

## 116 **Evans Blue Dye Procedure**

117           Blood samples for PV analysis were collected at six, 12, 18, and 24 h of age. The stock  
118 solution of EBD was prepared by weighing 2 g of 750 g kg<sup>-1</sup> EBD into a volumetric flask mixed  
119 with 100 mL of 9 g L<sup>-1</sup> sterile NaCl and stored at 3°C. At all sampling times, 10 mL of blood was  
120 collected into Vacutainer tubes containing freeze-dried Na-heparin (Kendall, Mansfield, MA,  
121 USA) via jugular venipuncture using a 22-gauge needle. After the pre-injection blood sample  
122 was collected, approximately 1.5 mL of stock solution was injected into the jugular vein. The  
123 syringe was weighed prior to and after injection so exact amount of dye administered was  
124 known. After injection, the dye was allowed to equilibrate for 10 min before a post-injection  
125 blood sample was obtained into 10 mL Na-heparinized Vacutainer tubes. Blood samples were  
126 centrifuged (CentraMP4R; International Equipment Company; Needham HTS, MA, USA) at  
127 1,310 × g at -4°C for 20 min (within 15 min of collection). The plasma was harvested and stored  
128 at -20°C until PV analysis via spectrophotometry. Absorbance of the spectrophotometer  
129 (Beckman DU 520; Brea, CA, USA) was set to 620 nm. Upon analysis, plasma samples were  
130 thawed and 1 mL of pre-injection plasma was transferred into a cuvette and the  
131 spectrophotometer was blanked against the sample. Fifty µl of standard dye solution (prepared  
132 by adding 1 mL of stock solution to a 100-mL volumetric flask which was diluted with 9 g L<sup>-1</sup>  
133 NaCl solution) was added to the plasma blank, vortexed and absorbance read. Finally,  
134 absorbance of the post-injection sample was obtained. The PV was calculated using the equation:  
135  $[1 / ((U \div S) \times 7.142857)] \times I$ , where U = absorbance of unknown (post-injection sample); S =  
136 absorbance of standard (pre-injection sample + 50 µl of standard dye solution); and I = µg of dye  
137 injected (mL of stock injected × 15 mg dye mL<sup>-1</sup> stock × 1000), 7.142857 is equal to the amount

138 of dye in the standard plasma sample (50  $\mu$ L of dye containing 0.15 mg dye  $\text{mL}^{-1}$  of standard  
139 dye solution added to 1 mL of plasma =  $7.5 \mu\text{g} \div 1.05\text{mL}$ ).

140

#### 141 **Blood Collection for Hematocrit**

142 A subsample of blood collected for PV analysis was obtained for evaluation of packed  
143 cell volume. An initial sample was obtained at birth before CR feeding for HCT analysis as well.  
144 Capillary tubes were centrifuged (Haematokrit 210; Andreas Hettich GmbH & Co; Germany) at  
145  $16,060 \times g$  at  $25^\circ\text{C}$  for 5 min.

#### 146 **Statistical Analysis**

147 Plasma volume and HCT values at six, 12, 18, and 24 h were analyzed using the mixed  
148 procedure of SAS® (Version 9.3; SAS Institute, Inc. Cary, NC, USA) as a randomized complete  
149 block design according to the following model:

$$150 Y_{ijk} = \mu + B_i + C_j + H_k + E_{ijk}$$

151 Where:

152 Y = the dependent variable;

153  $\mu$  = the overall mean;

154  $B_i$  = the random effect of block  $i$  ( $i = 1, \dots, 14$ );

155  $C_j$  = the fixed effect of the  $j^{\text{th}}$  CR feeding regimen ( $k = 1, 2$ );

156  $H_k$  = the fixed effect of the  $k^{\text{th}}$  hour ( $k = 6, 12, 18, \text{ and } 24$ );

157  $E_{ij}$  = the residual error  $\sim N(0, \sigma^2_e)$ .

158

159 Degrees of freedom were calculated using the Kenward-Rogers option of the MIXED procedure

160 of SAS® (Version 9.3; SAS Institute, Inc. Cary, NC, USA). Least square means were

161 determined for each treatment. Significant treatment effects were noted at  $P \leq 0.05$  and trends

162 were noted at  $0.05 < P \leq 0.10$ . Regression analysis was conducted using the regression procedure

163 of SAS using PV ( $\text{g kg}^{-1}$  BW) and HCT. Variables used in the regression analysis for PV were:

164 sex, HCT, initial HCT, BW and treatment. Variables used in the regression analysis for HCT



165 were: sex, initial HCT, BW, treatment, hour (for overall value), and PV (mL). Correlation  
166 coefficients were determined using the correlation procedure of SAS (Version 9.3; SAS Institute,  
167 Inc., Cary NC, USA),

168

169

## Results

170 Mean BW was  $46 \pm 4.61$  kg, mean initial HCT was  $39.4 \pm 6.82$  %, mean HCT was  $35.2 \pm$   
171  $5.83$  %, and mean PV was  $3,866 \pm 2022$  mL (Tables 1 and 2). Three calves on the C<sub>1</sub> and two  
172 calves on C<sub>2</sub> treatment groups were fed using an esophageal tube feeder. The calves on treatment  
173 C<sub>1</sub> required all of the CR to be fed in this manner. Whereas, one calf on C<sub>2</sub> required all of the CR  
174 fed in this manner while the other calf consumed its first feeding via nipple bottle, but was fed  
175 the total six h dose with the esophageal feeder. Initially, 28 samples per time point were  
176 obtained; however, due to sampling error or hemolysis, 4 calves were removed from analysis (2  
177 per treatment). Analyses of PV (mL, and mL kg<sup>-1</sup> of BW) along with HCT data are presented in  
178 Table 1. Mean PV for all calves regardless of treatment at six, 12, 18, and 24 h were 78.2, 89.2,  
179 83.9, and 90.7 mL kg<sup>-1</sup> of BW, respectively. Treatment did not affect PV expressed as either mL  
180 or mL kg<sup>-1</sup> of BW at any time point (Table 1). Hematocrit did not vary at any time point (Table  
181 1). Mean, minimum and maximum values for PV, BW, initial HCT and HCT are in Table 2.  
182 Correlation coefficients (Table 3) for PV showed strong relationships between PV, HCT, initial  
183 HCT and treatment. Prediction equations for PV as a percent of BW had coefficients of multiple  
184 determinations ( $r^2$ ) values that ranged from 0.19 for the 12 h prediction equation to 0.32 for the  
185 6 h equation suggesting a weak relationship (Table 4).

186 Prediction equations for HCT had  $r^2$  values that ranged from 0.69 for the 6 h equation to  
187 0.84 for the 24 h equation (Table 5). Body weight and initial HCT can be used to predict HCT at

188 12 and 24 h. Whereas, these variables plus sex of the calf can predict HCT at six and 18 h  
189 (Figure 1).

## 190 **Discussion**

191 Evans blue dye has been used as a marker of PV in cattle (Dalton and Fisher, 1961;;  
192 Payne et al., 1967; Möllenberg et al., 1975; Thornton and English, 1978; Besser and Osborn,  
193 1993 ), sheep (English, 1966; Mackie, 1976; van Waversveld and van Bruchem, 1985), horses  
194 (McKeever et al., 1988), and pigs (McCance and Widdowson, 1959). Estimates of PV using  
195 EBD and other methods typically indicate close correlation; however, van Waversveld and van  
196 Brushem (1985) reported that EBD estimated total blood volume but not PV. However a strong  
197 correlation was found between PV and hematocrit ( $P = 0.006$ ). Quigley et al. (1998), with 146  
198 calves sampled, also reported large variability in PV among calves even with the same BW (39  
199 kg) with PV as  $\text{mL kg}^{-1}$  of BW ranging from 60 – 131. This is similar to calves used on the  
200 current study with a BW of 40 kg having PV ranging from 67 – 131.

201 Research in human physiology has indicated that PV can be estimated by using a  
202 patient's height, weight, and HCT; however, equations are only reliable at normal HCT.  
203 Sprenger et al. (1987) were able to modify equations and allow for accurate calculation of PV  
204 even with the use of pathological HCT but PV was measured using 51CR-method and not EBD.  
205 Quigley et al. (1998) reported that birth BW was the best predictor of PV, though the relationship  
206 was not linear; however, multiple regression analysis using HCT and BW yielded no correlation  
207 with PV. Our results may have been affected by the use of EBD and perhaps a more accurate  
208 method of PV estimation should be used in the future.

209 There was no difference in PV between treatments across the 4 sampling times.

210 Regardless of treatment, mean PV for all calves at six, 12, 18, and 24 h were 78.2, 89.2, 83.9

211 90.7 mL kg<sup>-1</sup> of BW, respectively. Quigley et al. (1998) reported PV at 24 h at 98.6 mL kg<sup>-1</sup> of  
212 BW. The authors also adjusted for loss of EBD from plasma caused by single-point sampling by  
213 multiplying PV as mL kg<sup>-1</sup> of BW by 9.06 (correction factor determined as the mean  
214 overestimations reported by Mackie (1976) and Möllenberg et al. (1975)), which resulted in a  
215 new mean PV of 89 mL kg<sup>-1</sup> of BW. Adjusting PV at 24 h by this same method reduces our  
216 estimate of PV to 82.5 mL kg<sup>-1</sup> of BW. Adjusting for EBD loss at six, 12 and 18 h results in  
217 new PV values of : 71.1, 81.1, 76.3 ml kg<sup>-1</sup> . Quigley et al. (1998) also indicate that age at  
218 sampling averaged 24.3 h but ranged from 22.75 – 28.75 h. All calves used in the current study  
219 had samples obtained within ±15 min from designated time point (six 12, 18, or 24 h). The  
220 differences in sampling time may have impacted PV values. Colostrum intake type and volume  
221 (3L of CR vs, 1.5-4L colostrum) were also inconsistent between studies. The current study  
222 utilized CR with esophageal tube feeding used to ensure the total volume of CR was consumed.  
223 Quigley et al. (1998) utilized colostrum with intake averaging 3.76 L but ranging from 1.50 –  
224 4.00 L. Differences in volume of colostrum consumed may have affected hydration status and,  
225 therefore, PV of their calves.

226 Calves on the current study were also sampled at four total time points, as compared to  
227 one time point in the previously mentioned study. This may have affected the integrity of the  
228 blood vessel leading to an increase in loss of EBD during the equilibration period. Samples  
229 obtained at six, 12, 18 h and 24 h may have been slightly inaccurate because of the rate of loss of  
230 EBD from circulation. Younger animals lose EBD from circulation more rapidly than older  
231 animals (Thornton and English, 1978). By 24 h of age, calves were more apt to struggle during  
232 administration of dye and blood sampling. This may have increased EBD loss from circulation  
233 therefore adding to sample inaccuracy.

234 Results from this experiment provide researchers with values to allow determination of  
235 AEA when CR is fed. Also, determination of AEA at different time points over the first 24 h of  
236 life which will allow for a more accurate determination of when IgG absorption is maximized.

237 Hematocrit was not different between treatments. Hematocrit is a measure of packed cell  
238 volume, which is an indicator of hydration status. Though the total volume was identical,  
239 spreading the feeding between 2 time periods may have affected hydration status. Overall, HCT  
240 decreased from birth to 24 h indicating an increase in hydration status. Hematocrit, sex,  
241 treatment and initial HCT were correlated with PV in this study. Initial BW, sex and precolostral  
242 HCT can be used to predict HCT at six, 12, 18 and 24 h after birth in Holstein calves receiving  
243 CR.

### 244 **Conclusion**

245 Feeding CR in either one or split into two feedings did not affect PV, or HCT  
246 measurements. Plasma volume was highly correlated to sex, HCT, initial HCT and feeding  
247 regimen. Prediction equations to determine PV utilizing BW, sex, feeding regimen, initial HCT  
248 and HCT had correlation coefficients that ranged from 0.25 to 0.40 indicating a weak  
249 relationship. However, a better relationship was found for predicting HCT with  $r^2$  ranging from  
250 0.69 to 0.84. An initial precolostral blood sample along with BW can accurately predict HCT at  
251 12, and 24 h. Those values along with an adjustment for sex of the calf can accurately predict  
252 hematocrit at six and 18 h after birth. Researchers evaluating CR can utilize these data to  
253 accurately determine AEA and pinpoint when IgG absorption is maximized..

254

255

256

257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283

**ACKNOWLEDGEMENTS**

This project was supported by the Multi-State Hatch NH00512-R (NC-1042). The authors would like to thank the staff of the Fairchild Teaching and Research Center (University of New Hampshire) for the use of calves and their assistance throughout the study and Dr. Nadine Berenguier, Professor of French, Department of Languages, Literatures and Cultures (University of New Hampshire) for translating the abstract. Partial funding was provided by the New Hampshire Agricultural Experiment Station. This is Scientific Contribution number 2544.

284 Besser, T. E. and Osborn, D.. 1993. Effect of bovine serum albumin on passive transfer of  
285 immunoglobulin G1 to newborn calves. *Vet. Immunol. Immunopathol.* 37:321-327.

286 Dalton, R. G., and Fisher, E.W.. 1961. Plasma and blood volumes in Ayrshire cattle. *Br. Vet. J.*  
287 117:115-119.

288 English, P. B. 1966. A study of water and electrolyte metabolism in sheep. II. The volumes of  
289 distribution of antipyrine, thiosulfate, and T1824 (Evans blue) and values for certain  
290 extracellular fluid constituents. *Res. Vet. Sci.* 7:258-275.

291 Mackie, W. S. 1976. Plasma volume measurements in sheep using Evans' blue and continuous  
292 blood sampling. *Res. Vet. Sci.* 21:108-109.

293 McCance, R. A. and Widdowson, E.M.. 1959. The effect of colostrum on the composition and  
294 volume of the plasma of new-born piglets. *J. Physiol.* 145:547-550.

295 McEwen, S. A., Martin, W., Clarke, R.C.,and Tamblyn, S.E.. 1988. A prevalence survey of  
296 *Salmonella* in raw milk in Ontario, 1986– 87. *J. Food Prot.* 51:963–965.

297 McKeever, K. H. Schurg, W.A.,and Convertino, V.A.. 1988. A modified Evans blue dye method  
298 for determining plasma volume in the horse. *Equine Vet. Sci.* 8:208-212.

299 Möllerberg, L., Ekman, L., and Jacobsson, S.. 1975. Plasma and blood volume in the calf from  
300 birth till 90 days of age. *Acta. Vet. Scand.* 16:178-185.

301 Payne, E., Ryley, J.W., and Gartner, R.J.W. 1967. Plasma, blood, and extracellular fluid in  
302 grazing Hereford cattle. *Res. Vet. Sci.* 8:20-26.

303 Quigley, J. D. III, Drewry, J.J.,and Martin, K.R.. 1998. Estimation of plasma volume in Holstein  
304 and Jersey calves. *J. Dairy Sci.* 81:1308-1312.

305 Sprenger, K. B., Huber, K., Kratz, W., and Henze, E.. 1987. Nomograms for the prediction of  
306 patient's plasma volume in plasma exchange therapy from height, weight, and  
307 hematocrit. *J. Clin. Apher.* 3:185-190.

308 Thornton, J. R. and English, P.B.,. 1978. Body water of calves: Change in distribution with  
309 diarrhoea. *Br. Vet. J.* 134: 445-453.

310 Van Waversveld, J. and van Bruchem, J. 1985. Estimation of blood (plasma) volume in sheep  
311 with Evans blue and bromsulphalein. *Neth. J. Agric. Sci.* 33:325-327.

312

313

314 Table 1. Calves, body weight, plasma volume and hematocrit for calves fed colostrum replacer  
 315 either all at birth or split between two feedings.

Item	Treatment <sup>1</sup>		
	C <sub>1</sub>	C <sub>2</sub>	SE <sup>2</sup>
BW, kg (+/- SD)	44.90 +/- 4.0	46.70 +/- 5.1	
Initial HCT, %	38.25 +/- 3.36	41.09 +/- 3.50	
PV <sup>3</sup> , 6h, mL	3,522	3,590	289
PV, 12h, mL	3,612	3,967	289
PV, 18h, mL	3,221	4,376	289
PV, 24h, mL	3542	3,615	289
PV, 6h, mL kg <sup>-1</sup> BW	78.5	78.7	8.2
PV, 12h, mL kg <sup>-1</sup> BW	90.2	88.2	8.2
PV, 18h, mL kg <sup>-1</sup> , BW	80.0	87.9	8.2
PV, 24h, mL kg <sup>-1</sup> BW	81.4	100.0	8.2
HCT <sup>5</sup> , 6h, %	36.8	38.0	1.3
HCT, 12h, %	34.3	35.1	1.3
HCT, 18h, %	33.8	35.4	1.3
HCT, 24h, %	34.2	35.0	1.3

316  
 317 <sup>1</sup>Treatment C<sub>1</sub> = all colostrum replacer is fed at time 0 h, C<sub>2</sub> = colostrum replacer fed 2/3 at time

318 0 h and 1/3 at 6h.

319 <sup>2</sup>Standard error.

320 <sup>3</sup>Repeated measures by hour for plasma volume (mL) was not different (P< 0.13).

321 <sup>4</sup>Repeated measures by hour for plasma volume (% of body weight) was not different (P = 0.54).

322 <sup>5</sup>Repeated measures by hour for hematocrit was not different (P= 0.99).

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344



345 Table 2. Mean, minimum and maximum values and standard deviations for data used in  
346 development of prediction equations.

Variable	n	Mean	SD	Minimum	Maximum
PV, mL	95	3,430	1,333	1376	7176
BW, kg	97	45.87	4.68	40.0	56.0
Initial HCT, %	97	39.63	6.90	24.0	54.0
HCT, %	94	35.37	5.78	20.0	45.0

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362 Table 3. Correlation coefficients for plasma volume, sex, body weight, hematocrit, initial  
 363 hematocrit, and treatment.

Item	PV <sup>1</sup> (mL)	Sex	BW <sup>2</sup>	HCT <sup>3</sup>	Hi <sup>4</sup>	Trt <sup>5</sup>
PV (mL)	1.000	0.03	-0.12	-0.21	-0.26	0.13
		P = 0.74	P = 0.28	P = 0.06	P = 0.02	P = 0.24
Sex		1.000	-0.51	-0.01	0.07	-0.08
			P < 0.0001	P = 0.92	P = 0.49	P = 0.39
BW			1.000	0.33	0.09	0.20
				P = 0.0005	P = 0.38	P = 0.04
HCT				1.000	0.81	0.08
					P < 0.0001	P = 0.41
Hi					1.000	0.06
						P = 0.51
Trt						1.000

364

365 <sup>1</sup>Plasma volume in mL.

366 <sup>2</sup>Body weight.

367 <sup>3</sup>Hematocrit.

368 <sup>4</sup>Initial Hematocrit.

369 <sup>5</sup>Treatment- feeding regimen (one feeding of CR or 2 feedings of CR).

370

371

372 Table 4. Prediction equations for plasma volume (% of BW) at various time points<sup>1</sup>.

Hour <sup>2</sup>	Intercept	Sex <sup>3</sup>	BW <sup>4</sup> kg	Initial HCT <sup>5</sup> %	HCT <sup>6</sup>	Trt <sup>7</sup>	r <sup>2</sup>
6	21.80	-	-0.28	-0.11	-	1.66	0.32
12	19.93	-	-	-0.29	-		0.19
18	14.38	2.51	-	-	-0.28	-	0.22
24	11.58	2.80	-	-	-0.31	2.56	0.31

373 <sup>1</sup>Variables with blank spaces indicate that P >= 0.25.

374 <sup>2</sup>Hour corresponds to time after birth.

375 <sup>3</sup>Sex ,1 = bull, 2 = heifer.

376 <sup>4</sup>Body weight.

377 <sup>5</sup>Initial Hematocrit.

378 <sup>6</sup>Hematocrit.

379 <sup>7</sup>Treatment 1 = fed at 0h, 2 = fed in 2 feedings.

380

381

382

383

384

385

386

387

388

389 Table 5. Prediction equations for hematocrit<sup>1</sup>

Hour	Intercept	Sex <sup>2</sup>	BW <sup>3</sup> , kg	Initial HCT, %	r <sup>2</sup>
6	-3.74	2.05	0.33	0.58	0.69
12	-2.59	-	0.26	0.71	0.74
18	-13.7	2.08	0.43	0.64	0.77
24	-12.14	-	0.39	0.73	0.84

390

391 <sup>1</sup>Variables with blank spaces indicate that P >= 0.25.

392

393 <sup>2</sup> Sex ,1 = bull, 2 = heifer.

394

395 <sup>3</sup>Body weight.

396

397

398

399

400

401

402

403

404

405


406

407

408

409

410



411  
412  
413  
414  
415  
416

Figure 1. Predicted hematocrit versus measured hematocrit.  
Cabral et al