

University of New Hampshire

## University of New Hampshire Scholars' Repository

---

Life Sciences Faculty Scholarship

Life Sciences

---

7-19-2018

### Genome Sequence of the Halophilic Bacterium *Kangiella spongicola* ATCC BAA-2076T

Aria Underriner  
*Seacoast School of Technology*

Tyler Silverwood  
*Seacoast School of Technology*

Carolyn Kelley  
*Seacoast School of Technology*

Kyle S. MacLea  
*University of New Hampshire, Manchester, kyle.maclea@unh.edu*

Follow this and additional works at: [https://scholars.unh.edu/unhmbiology\\_facpub](https://scholars.unh.edu/unhmbiology_facpub)

#### Comments

This is an Author's Original Manuscript/Accepted Manuscript of an article published by American Society for Microbiology in Microbiology Resource Announcements in 2018, available online: <https://dx.doi.org/10.1128/MRA.00847-18>

---

#### Recommended Citation

Underriner A, Silverwood T, Kelley C, MacLea KS. 2018. Genome sequence of the halophilic bacterium *Kangiella spongicola* ATCC BAA-2076T. *Microbiol Resour Announc* 7:e00847-18. <https://doi.org/10.1128/MRA.00847-18>.

This Article is brought to you for free and open access by the Life Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Life Sciences Faculty Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact [Scholarly.Communication@unh.edu](mailto:Scholarly.Communication@unh.edu).



# Genome Sequence of the Halophilic Bacterium *Kangiella spongicola* ATCC BAA-2076<sup>T</sup>

Aria Underriner,<sup>a</sup> Tyler Silverwood,<sup>a</sup> Carolyn Kelley,<sup>a</sup>  Kyle S. MacLea<sup>b,c,d</sup>

<sup>a</sup>Biomedical Science and Technology Program, Seacoast School of Technology, Exeter, New Hampshire, USA

<sup>b</sup>Biology Program, University of New Hampshire, Manchester, New Hampshire, USA

<sup>c</sup>Biotechnology Program, University of New Hampshire, Manchester, New Hampshire, USA

<sup>d</sup>Department of Life Sciences, University of New Hampshire, Manchester, New Hampshire, USA

**ABSTRACT** The Gram-negative genus *Kangiella* contains a number of halophilic species that display high levels of iso-branched fatty acids. *Kangiella spongicola* was isolated from a marine sponge, *Chondrilla nucula*, from the Florida Keys in the United States. A genome assembly of 2,825,399 bp with a 44.31% G+C content was generated from strain A79<sup>T</sup> (=ATCC BAA-2076<sup>T</sup>).

Bacteria within the family *Alcanivoraceae* and class *Gammaproteobacteria* include the genus *Kangiella*. *Kangiella* is notable for being a newly described genus of halophilic bacteria with an unusually high production of iso-branched fatty acids (1, 2). *Kangiella spongicola* ATCC BAA-2076<sup>T</sup> is a member of this genus recently discovered from the marine sponge *Chondrilla nucula*, collected from the Florida Keys in the United States (2). This organism is a Gram-negative nonmotile rod-shaped bacterium that is able to survive in environments with up to 15% NaCl and a wide range of growth temperatures and pH values. The genome sequence generated for *K. spongicola* may be a useful data point in the understanding of sponge microbiota in marine environments, especially in combination with the five other members of the genus (1, 3, 4) that have been sequenced thus far (5, 6).

*Kangiella spongicola* ATCC BAA-2076<sup>T</sup> was bought in lyophilized form from ATCC (Manassas, VA, USA). It was rehydrated in marine broth 2216 (BD, Franklin Lakes, NJ, USA) and incubated in a shaking incubator at 30°C for 24 h. Inoculated marine broth was spread on marine agar 2216 (BD), incubated at the same temperature, and, from the streak plate, a single colony was selected from which to grow a larger liquid culture for production of genomic DNA (gDNA) using a DNA minikit (Qiagen, Valencia, CA, USA). Pure gDNA was tagged with sequence adapters concurrent with fragmentation using the Nextera library prep kit (Illumina, San Diego, CA, USA). Tagged fragments were then sequenced on an Illumina HiSeq 2500 (Hubbard Center for Genome Studies, Durham, NH, USA) instrument. The generated 250-bp paired-end read sequences were bioinformatically trimmed before assembly using Trimmomatic (7). Paired and trimmed reads were assembled into a draft genome using the default settings of SPAdes version 3.11.1 (8). Small contigs and contaminants were removed, and the 140 remaining contigs were analyzed with the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) process for gene prediction and annotation (9) and found to have a total sequence length of 2,825,399 bp, representing an average coverage of 326×. Its closest relatives, *K. korensis* and *K. aquimarina*, are of similar size at 2.85 Mbp and 2.68 Mbp, respectively. Of the 140 contigs, the bulk of the sequence data were found in four contigs with lengths of 1,220,304 bp, 675,233 bp, 545,949 bp, and 83,637 bp. Of the remaining 136 contigs, all were less than 6,001 bp in length. The  $N_{50}$  value was 675,233 bp with an  $L_{50}$  value of 2, as determined by QUAST (10). The G+C content of 44.31% was consistent

Received 12 June 2018 Accepted 26 June 2018 Published 19 July 2018

**Citation** Underriner A, Silverwood T, Kelley C, MacLea KS. 2018. Genome sequence of the halophilic bacterium *Kangiella spongicola* ATCC BAA-2076<sup>T</sup>. *Microbiol Resour Announc* 7:e00847-18. <https://doi.org/10.1128/MRA.00847-18>.

**Editor** Jason Stajich, University of California, Riverside

**Copyright** © 2018 Underriner et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kyle S. MacLea, [Kyle.MacLea@UNH.edu](mailto:Kyle.MacLea@UNH.edu).

A.U. and T.S. contributed equally to this work.

with the initial report of 44.9% (2). A total of 2,544 genes, 2,460 coding DNA sequences (CDS), 36 pseudogenes, 5 rRNAs, and 38 tRNAs were annotated using NCBI PGAP.

**Data availability.** This *Kangiella spongicola* ATCC BAA-2076<sup>T</sup> whole-genome shotgun sequence (WGS) project has been deposited in DDBJ/ENA/GenBank under accession number [Q1CH00000000](https://www.ncbi.nlm.nih.gov/nuccore/Q1CH00000000). The version described in this paper is the first version, Q1CH01000000.

## ACKNOWLEDGMENTS

Sequencing and bioinformatics analysis were undertaken at the Hubbard Center for Genome Studies at UNH, supported by NH-INBRE, with the kind assistance of Kelley Thomas, Toni Westbrook, and Stephen Simpson. K.S.M. thanks lab associate Oliver Harmon for the inspiration to sequence a bacterium from a marine sponge. This work was a project of the Microbiology Education through the Genome Annotation-New Hampshire (MEGA-NH) program.

The Department of Life Sciences at UNH Manchester provided funds for purchasing the bacterial strain. Sequencing costs were supported by New Hampshire-INBRE through an Institutional Development Award (IDeA), P20GM103506, from the National Institute of General Medical Sciences of the NIH. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

## REFERENCES

1. Yoon J-H, Oh T-K, Park Y-H. 2004. *Kangiella koreensis* gen. nov., sp. nov. and *Kangiella aquimarina* sp. nov., isolated from a tidal flat of the Yellow Sea in Korea. *Int J Syst Evol Microbiol* 54:1829–1835. <https://doi.org/10.1099/ijs.0.63156-0>.
2. Ahn J, Park J-W, McConnell JA, Ahn Y-B, Häggblom MM. 2011. *Kangiella spongicola* sp. nov., a halophilic marine bacterium isolated from the sponge *Chondrilla nucula*. *Int J Syst Evol Microbiol* 61:961–964. <https://doi.org/10.1099/ijs.0.021733-0>.
3. Yoon J-H, Kang S-J, Lee S-Y, Lee J-S, Oh T-K. 2012. *Kangiella geojedonensis* sp. nov., isolated from seawater. *Int J Syst Evol Microbiol* 62:511–514. <https://doi.org/10.1099/ijs.0.029314-0>.
4. Lee S-Y, Park S, Oh T-K, Yoon J-H. 2013. *Kangiella sediminilitoris* sp. nov., isolated from a tidal flat sediment. *Int J Syst Evol Microbiol* 63: 1001–1006. <https://doi.org/10.1099/ijs.0.040691-0>.
5. Han C, Sikorski J, Lapidus A, Nolan M, Glavina Del Rio T, Tice H, Cheng J-F, Lucas S, Chen F, Copeland A, Ivanova N, Mavromatis K, Ovchinnikova G, Pati A, Bruce D, Goodwin L, Pitluck S, Chen A, Palaniappan K, Land M, Hauser L, Chang Y-J, Jeffries CD, Chain P, Saunders E, Brettin T, Göker M, Tindall BJ, Bristow J, Eisen JA, Markowitz V, Hugenholtz P, Kyrpides NC, Klenk H-P, Detter JC. 2009. Complete genome sequence of *Kangiella koreensis* type strain (SW-125T). *Stand Genomic Sci* 1:226–233. <https://doi.org/10.4056/sigs.36635>.
6. Choe H, Kim S, Oh J, Nasir A, Kim BK, Kim KM. 2015. Complete genome of *Kangiella geojedonensis* KCTC 23420<sup>T</sup>, putative evidence for recent genome reduction in marine environments. *Mar Genomics* 24:215–217. <https://doi.org/10.1016/j.margen.2015.05.015>.
7. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
8. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
9. Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Ciufu S, Li W. 2013. Prokaryotic Genome Annotation Pipeline. National Center for Biotechnology Information, Bethesda, MD.
10. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUASt: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.