

University of New Hampshire

University of New Hampshire Scholars' Repository

Earth Sciences Scholarship

Earth Sciences

3-20-1995

Global perspective of nitrate flux in ice cores

Qinzhao Yang

University of New Hampshire - Main Campus

Paul A. Mayewski

University of New Hampshire - Main Campus

Sallie I. Whitlow

University of New Hampshire - Main Campus

Mark S. Twickler

University of New Hampshire - Main Campus

Michael Morrison

University of New Hampshire - Main Campus

See next page for additional authors

Follow this and additional works at: https://scholars.unh.edu/earthsci_facpub



Part of the [Atmospheric Sciences Commons](#)

Recommended Citation

Yang, Q., P. A. Mayewski, S. Whitlow, M. Twickler, M. Morrison, R. Talbot, J. Dibb, and E. Linder (1995), Global perspective of nitrate flux in ice cores, *J. Geophys. Res.*, 100(D3), 5113–5121, doi:10.1029/94JD03115.

This Article is brought to you for free and open access by the Earth Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Earth Sciences Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

Authors

Qinzhao Yang, Paul A. Mayewski, Sallie I. Whitlow, Mark S. Twickler, Michael Morrison, R. Talbot, Jack E. Dibb, and Ernst Linder

Global perspective of nitrate flux in ice cores

Qinzhao Yang,¹ Paul A. Mayewski,¹ Sallie Whitlow,¹ Mark Twickler,¹
Michael Morrison,¹ Robert Talbot,² Jack Dibb,¹ and Ernst Linder³

Abstract. The relationships between the concentration and the flux of chemical species (Cl^- , NO_3^- , SO_4^{2-} , Na^+ , K^+ , NH_4^+ , Mg^{2+} , Ca^{2+}) versus snow accumulation rate were examined at GISP2 and 20D in Greenland, Mount Logan from the St. Elias Range, Yukon Territory, Canada, and Sentik Glacier from the northwest end of the Zanskar Range in the Indian Himalayas. At all sites, only nitrate flux is significantly ($\alpha = 0.05$) related to snow accumulation rate. Of all the chemical series, only nitrate concentration data are normally distributed. Therefore we suggest that nitrate concentration in snow is affected by postdepositional exchange with the atmosphere over a broad range of environmental conditions. The persistent summer maxima in nitrate observed in Greenland snow over the entire range of record studied (the last 800 years) may be mainly due to NO_x released from peroxyacetyl nitrate by thermal decomposition in the presence of higher OH concentrations in summer. The late winter/early spring nitrate peak observed in modern Greenland snow may be related to the buildup of anthropogenically derived NO_y in the Arctic troposphere during the long polar winter.

Introduction

Ice core records from polar regions provide a unique record of environmental and climatic change [e.g., *Dansgaard and Oeschger*, 1989; *Lorius et al.*, 1989; *Mayewski et al.*, 1993a, 1994] and illustrate the influence of anthropogenic emissions on the chemistry of the remote atmosphere [e.g., *Neftel et al.*, 1985; *Mayewski et al.*, 1986, 1990a; *Mayewski and Legrand*, 1990]. They have also contributed to our understanding of atmospheric chemistry, atmospheric circulation and biogeochemical cycling [e.g., *Legrand et al.*, 1988, 1991; *Delmas and Legrand*, 1989; *McElroy*, 1989; *Mayewski et al.*, 1993a, 1994].

Nitrate is one of the major chemical species in polar snow. As such, measurements of nitrate in ice cores may provide information for understanding the complex atmospheric nitrogen cycle [e.g., *Neubauer and Heumann*, 1988; *Legrand and Kirchner*, 1990; *Mayewski and Legrand*, 1990; *Mayewski et al.*, 1990a; *Mulvaney and Wolff*, 1993]. However, the sources, transport pathways, and scavenging processes of nitrate deposited on glaciers are not well understood [*Wolff*, 1995]. Understanding these relationships will improve our interpreta-

tion of nitrate data retrieved from worldwide ice cores.

The relationships between snow accumulation rate and nitrate concentration/flux are important because they can be used to assess the dominant processes of nitrate incorporation into snow. The seasonal variation of nitrate concentration in snow is presumably related to the source of nitrate precursor in the atmosphere.

Previous studies conducted by *Herron* [1982], *Legrand and Delmas* [1986], and *Legrand and Kirchner* [1990] investigated relationships between nitrate concentration and flux versus snow accumulation rate in Greenland and Antarctica. However, there is a discrepancy in their conclusions. *Herron* [1982] concluded that nitrate concentration decreased with increasing snow accumulation rate, whereas *Legrand and Delmas* [1986] and *Legrand and Kirchner* [1990] found that nitrate concentration was independent of snow accumulation rate. In this paper we reevaluate this relationship using high-resolution snow chemical series from central Greenland as well as other series from southern Greenland, northwest Canada, and central Asia.

Annual nitrate summer peaks in snow have been reported for both Greenland and Antarctica [*Finkel et al.*, 1986; *Mayewski et al.*, 1987, 1990b; *Davidson et al.*, 1988; *Neubauer and Heumann*, 1988; *Legrand and Kirchner*, 1990; *Whitlow et al.*, 1992; *Mulvaney and Wolff*, 1993]. However, the cause for the timing of the peak is not yet well understood. We investigated cause(s) for the peak in Greenland snow to improve our understanding of the relationship between seasonal variations of nitrate measured in snow and possible sources of NO_x in the atmosphere.

Methodology

In this paper we first examine the relationship between the concentration of major cations and anions

¹ Glacier Research Group, Institute for the Study of Earth, Oceans and Space, University of New Hampshire, Durham.

² Atmospheric Chemistry Research Group, Institute for the study of Earth, Oceans and Space, University of New Hampshire, Durham.

³ Mathematics Department, University of New Hampshire, Durham.

Copyright 1995 by the American Geophysical Union.

Paper number 94JD03115.
0148-0227/95/94JD-03115\$05.00

(Cl^- , NO_3^- , SO_4^{2-} , Na^+ , K^+ , NH_4^+ , Mg^{2+} , Ca^{2+}) and their flux versus snow accumulation rate using 36 detailed sections (averaging 10 samples per year over sections ranging from 2 to 10 years) sampled from the GISP2 core located at Summit, Greenland, to see if nitrate behaves differently from other major ions. The same analysis is performed on several other glacio-chemical series, namely, 20D from southern Greenland; Mount Logan from the St. Elias Range; Yukon Territory, northwest Canada; and Sentik Glacier from the northwestern end of the Zaskar Range in Ladakh, Indian Himalayas. Ice core sites and some basic information about sites and sampling details are presented in Figure 1 and Table 1. Although the Sentik Glacier nitrate time series is shorter than the others, it is the only continuous high-resolution chemical record available from a low-latitude/high-elevation site.

Samples retrieved from GISP2, 20D, and Mount Logan were processed in a cold room at temperatures that did not exceed -12°C by individuals wearing precleaned polyethylene gloves, nonparticulating clean suits, and particle masks. All tools and containers used for sampling were precleaned in ultrapure water. Container blanks prepared on a frequent basis showed that these containers were free of contamination. All analyses were performed by ion chromatography using a DionexTM model 4040 with techniques described previously [e.g., Mayewski *et al.*, 1990b; Buck *et al.*, 1992]. Samples retrieved from Sentik Glacier were processed in the field, and nitrate was measured using a Technicon Auto AnalyzerTM system [Lyons and Mayewski, 1983; Mayewski *et al.*, 1984].

Each core was dated by using a variety of different seasonal indicators (e.g., major anions and cations, stable isotopes), historically documented volcanic events plus total beta activity signatures related to known

nuclear fallout events. Maximum dating error is ± 1 year throughout most of the cores [Lyons and Mayewski, 1983; Holdsworth and Peake, 1985; Mayewski *et al.*, 1986, 1993b]. Flux (F) of each chemical species is calculated based on that individual annual mean concentration (C) times snow accumulation rate (A) of the same year ($F = C \times A$).

Results and Discussion

Chemical Concentration/Flux versus Snow Accumulation Rate

The relationship between chemical (Cl^- , NO_3^- , SO_4^{2-} , Na^+ , K^+ , NH_4^+ , Mg^{2+} , Ca^{2+}) concentration ($\mu\text{g}/\text{kg}$) and snow accumulation rate ($\text{kg m}^{-2} \text{yr}^{-1}$) was examined using the GISP2 chemical data sets (exclusive of the anthropogenic period) (Figure 2). On the basis of the data shown in Figure 2, we examined each individual species using statistical methods of polynomial and linear regression to determine if there is a relationship. From Figure 2 it is clear that none of the concentration series of chemical species are statistically related to snow accumulation rate. This suggests that the annual mean concentration of chemical species in snow is independent of the annual snow accumulation rate at Summit, Greenland.

In Figure 3 we compare chemical flux ($\text{kg km}^{-2} \text{yr}^{-1}$) versus snow accumulation rate ($\text{kg m}^{-2} \text{yr}^{-1}$). A table of t values (Table 2) provides the significance of the slope coefficients for all chemical flux versus snow accumulation rate relationships in Figure 3. Nitrate has the highest t value and correlation coefficient (r value). For the other species, t values exceed the critical value of 1.645 (significance level $\alpha = 0.05$), but their r values are lower than 0.4, suggesting that less than 20% of to-




Figure 1. Location map showing the sampling sites at Summit, Greenland; 20D, south Greenland; Mount Logan; Sentik Glacier; and south pole.

Table 1. Ice Core Site Details

	Core				
	GISP2	20D ^a	Mount Logan ^b	Sentik Glacier ^c	South Pole ^d
Location	72.58°N, 38.47°W	65.01°N, 44.87°W	60.63°N, 140.50°W	35.00°N, 76.01°E	90.00°S
Elevation, meters above sea level	3207	2616	5340	4908	2880
Accumulation rate, m yr ⁻¹ , water equivalent	0.23	0.41	0.33	0.60	0.085
Mean annual temperature, °C	-31.5	-20.2	-28.9	-3	-50
Total samples analyzed	1524	1357	1853	103	759
Sampling resolution, samples/year	~10	~6.3	~6.4	~6.2	~4.3
Period covered by core (AD)	1989-1200 ^e	1984-1767	1980-1688	1980-1963	1980-1805

^aMayewski et al. [1986, 1990a].

^bHoldsworth and Peake [1985].

^cMayewski et al. [1984], Lyons and Mayewski [1983].

^dLegrand and Kirchner [1990].

^eRepresented by 36 uniformly distributed sections throughout the period. Each section ranges from 3 to 10 years.

tal flux variance can be explained by a linear fit to the relationship. Only nitrate flux has a statistical linear relationship with snow accumulation rate over the range measured (120 to 350 kg m⁻² yr⁻¹ water equivalent and 9.3 to 25.6 kg km⁻² yr⁻¹ of NO₃⁻).

The best fit linear regression of nitrate flux to snow accumulation rate for Summit, Greenland is

$$F = 0.75 + 0.071A \quad (r = 0.81) \quad (1)$$

where F is nitrate flux in kg km⁻² yr⁻¹, and A is the snow accumulation rate in kg m⁻² yr⁻¹.




Figure 2. Chemical concentration (Na⁺, NH₄⁺, K⁺, Mg²⁺, Ca²⁺, Cl⁻, NO₃⁻, SO₄²⁻) versus snow accumulation rate for the GISP2 detailed sections.




Figure 3. Chemical flux (Na^+ , NH_4^+ , K^+ , Mg^{2+} , Ca^{2+} , Cl^- , NO_3^- , SO_4^{2-}) versus snow accumulation rate for the GISP2 detailed sections. The solid line is the least squares fitted regression line to each species.

Herron [1982] investigated snow pits and ice cores recovered from different sites in Greenland and Antarctica and suggested that the relationship between nitrate concentration and snow accumulation rate may be expressed as

$$\text{NO}_3^- = \text{CA}^{-\frac{1}{2}} \quad (2)$$

where C is a constant, and A is the snow accumulation rate in $\text{kg m}^{-2} \text{yr}^{-1}$.

Legrand and Delmas [1986] and Legrand and Kirchner [1990] found a discrepancy in Herron's [1982] relationship for Antarctica. Their data suggested that

nitrate concentration was not related to snow accumulation rate, but nitrate flux was related to snow accumulation rate by the following expression:

$$F = 0.99 + 0.082A \quad (r = 0.92) \quad (3)$$

Our data sets (Figure 2) show that nitrate concentration is independent of snow accumulation, in harmony with the findings of Legrand and Delmas [1986] and Legrand and Kirchner [1990]. Furthermore, we found that the relationship between nitrate flux and snow accumulation at Summit, Greenland, agrees closely with the relationship at south pole [Legrand and Kirchner, 1990].

Table 2. Calculated t-Distribution Values and Correlation Coefficients (r) for the Regression Lines Displayed in Figure 2

	Na^+	NH_4^+	K^+	Mg^{2+}	Ca^{2+}	Cl^-	NO_3^-	SO_4^{2-}
GISP2	n=117							
t value	1.71	2.78	1.75	4.22	4.67	2.92	12.54	3.13
r	0.15	0.3	0.16	0.36	0.39	0.25	0.82	0.28

n, Data points used in regression.

To investigate the apparent uniqueness of the GISP2 nitrate data compared to other chemical species, we examined their concentration probability plots [Looney and Gullede, 1985]. The “straightness” of the probability plot is tested by using their correlation coefficients [Looney and Gullede, 1985; Minitab, 1993]. Nitrate is the only species whose concentration data are normally distributed at a significance level of 95% in Greenland snow (Table 3).

To test if a similar relationship between nitrate flux and snow accumulation rate exists at our other study sites, we used chemical data series from three additional sites: 20D (exclusive of the anthropogenic portion of the record), Mount Logan, and Sentik Glacier. At all three sites the relationship between nitrate flux and snow accumulation rate has the highest correlation coefficient (significant at the 95% level) of any of the species (Table 4).

The best fit linear regression equations of nitrate flux to snow accumulation rate at the three sites are

$$20D : F = 1.3 + 0.053A \quad (r = 0.71) \quad (4)$$

$$\text{Mount Logan} : F = -1.9 + 0.054A \quad (r = 0.79) \quad (5)$$

$$\text{Sentik Glacier} : F = 1.8 + 0.080A \quad (r = 0.74) \quad (6)$$

Since there is a relatively strong linear relationship between nitrate flux and snow accumulation rate at all three sites, we investigated the distributions of nitrate concentration versus the other chemical species concentration at these sites. The normality of chemical concentration probability plots from 20D, Mount Logan, and Sentik Glacier were tested as described for the GISP2 data set (Table 3). Nitrate is the only chemical species for which the correlation coefficient is higher than the critical value (at significance level of 95%) for each of the data sets.

Nitrate concentrations in snow at these sites are distinct from the other chemical species concentrations in that they are normally distributed. A recent surface snow chemistry study at Summit, Greenland [J. Dibb, unpublished data, 1994] indicates that the summer surface snowpack loses nitrate at some time after deposition, as has been reported for lower accumulation sites

in Antarctica [Mayewski and Legrand, 1990]. The normally distributed nitrate concentration may be a consequence of postdepositional alteration, potentially via exchange between the surface snow and the atmosphere [Neubauer and Heumann, 1988]. Hence we suggest that the exchange of nitrate between the snow and the atmosphere results in sufficient smoothing of the series to make them normally distributed (Table 3).

Two depositional styles dominate the incorporation of chemical species into snow, namely, wet and dry [Barrie, 1985; Davidson, 1989]. Equations (1) and (4)-(6) have small (<10%) *y*-intercept values relative to the individual mean nitrate flux of 17, 24, 17, and 74 $\text{km}^{-2} \text{yr}^{-1}$ in the snow at GISP2, 20D, Mount Logan, and Sentik Glacier, respectively. If the value of the *y* intercept of these equations reflects the flux contributed by dry depositional processes [Legrand and Kirchner, 1990], then nitrate is incorporated into snow predominantly by wet deposition at these sites. However, as discussed previously, the nitrate mean annual concentration is affected by the postdepositional processes. For nitrate the *y*-intercept value may not be an indication of the contribution of dry deposition in snow.

Since the annual mean concentration of nitrate is independent of snow accumulation rate at our study sites and at south pole [Legrand and Kirchner, 1990], the presence of a statistically significant linear relationship between nitrate flux and snow accumulation rate suggests that the supply of atmospheric nitrate is large enough so that it is never depleted over the accumulation range studied here. If this is the case, we would suggest that annual mean nitrate concentration in snow is proportional to its annual mean concentration in the atmosphere despite postdepositional process, as long as the depositional environment does not vary significantly.

Sources for Nitrate Summer Peak

As mentioned previously, one of the important characteristics of nitrate concentrations in Greenland and Antarctic snow is that there is a persistent nitrate summer peak [Finkel et al., 1986; Mayewski et al., 1987, 1990b; Neubauer and Heumann, 1988; Steffensen, 1988;

Table 3. Normality Tests for Chemical Concentration Data Distribution

	Na ⁺	NH ₄ ⁺	K ⁺	Mg ²⁺	Ca ²⁺	Cl ⁻	NO ₃ ⁻	SO ₄ ²⁻
GISP2	n=117	r critical=0.988 ^a						
	r 0.903	0.717	0.831	0.966	0.909	0.931	0.996	0.907
20D	n=174	r critical=0.990						
	r 0.926	0.822	0.649	0.935	0.916	0.917	0.991	0.954
Mount Logan	n=220	r critical=0.992						
	r 0.683	0.825	0.847	0.884	0.834	0.954	0.995	0.955
Sentik Glacier	n=16	r critical=0.942						
	r 0.893	*	*	*	*	0.915	0.965	*

n, Data points used in test. Asterisk, species not measured.
^a(After Table 13.1 in *Minitab Reference Manual* [1993].

Table 4. Calculated t-Distribution Values and Correlation Coefficients (r) Between Chemical Flux and Snow Accumulation Rate at 20D, Mount Logan, and Sentik Glacier

	Na ⁺	NH ₄ ⁺	K ⁺	Mg ²⁺	Ca ²⁺	Cl ⁻	NO ₃ ⁻	SO ₄ ²⁻
20D n=174								
t value	6.98	2.22	2.38	6.59	5.66	7.81	12.88	7.39
r	0.43	0.15	0.16	0.41	0.36	0.47	0.66	0.45
Mount Logan n=220								
t value	4.95	6.21	6.4	4.95	5.48	6.79	18.89	11.29
r	0.32	0.39	0.4	0.32	0.35	0.42	0.79	0.61
Sentik Glacier n=16								
t value	1.59	*	*	*	*	2.01	4.26	*
r	0.38	*	*	*	*	0.46	0.74	*

n: Data points used in regression. Asterisk, species not measured.

Davidson, 1989; Whitlow *et al.*, 1992; Mulvaney and Wolff, 1993]. However, the mechanism(s) that causes this peak is not well understood. To investigate the source of the nitrate in snow, we start our investigation of the cause(s) for the annual nitrate summer peak in Greenland snow by examining controls on the formation of seasonal variations in nitrate in the atmosphere.

Nitric acid (HNO₃) is formed primarily by the reaction of NO₂ + OH in the presence of sunlight and is well known as a main sink for atmospheric NO_x [Logan, 1983; Warneck, 1988; Ehhalt *et al.*, 1992]. HNO₃ is also formed by the nighttime reaction between N₂O₅ and H₂O [Russell *et al.*, 1986; Mozurkewich and Calvert, 1988]. Various processes and sources can contribute to the nitrate budget at high latitude, including combustion of fossil fuel (especially in the northern hemisphere [Logan, 1983; Ehhalt *et al.*, 1992]), biomass burning, lightning, ammonia oxidation and soil exhalation, in addition to processes acting within the stratosphere (galactic cosmic rays, N₂O oxidation [Levy *et al.*, 1980]). A few scientists have investigated the possible origin of nitrate in polar precipitation, suggesting an important role for lightning and N₂O stratospheric oxidation [Legrand and Kirchner, 1990; Wolff, 1995] and sometimes processes acting in the high atmosphere [Parker and Zeller, 1979; Zeller *et al.*, 1986]. Using a modeling approach, Legrand *et al.* [1989] suggested that lightning, soil exhalation, and N₂O stratospheric oxidation are likely major contributors to the natural Antarctic nitrate budget. The major sources of NO_x in the northern hemisphere troposphere are combustion of fossil fuels, lightning formed at high altitude/low latitude, and NO exhalation from soils [Logan, 1983; Legrand and Kirchner, 1990; Mayewski *et al.*, 1990a].

With the rather short residence time of NO_x in the troposphere (less than a day in summer and a few days in winter [Singh, 1987]), it is unlikely that it is directly transported from primary sources to interior central Greenland. Most of the nitric acid and particulate nitrate is believed to be removed during transport [Barrie, 1985; Lyons *et al.*, 1990; Bottenheim *et al.*, 1993]. However, the formation of peroxyacetyl nitrate (PAN) has recently received considerable attention as a carrier

and reservoir of NO_x for the northern high latitudes due to its longer residence time (up to months in winter) and strong thermal decomposition characteristics [Cox and Roffey, 1977; Singh, 1987; Honrath and Jaffe, 1992; Singh *et al.*, 1992a]. Singh and Hanst [1981], and Singh and Salas [1983] suggested that PAN not only is associated with polluted air but also is an important constituent of the natural atmosphere. The production rate of OH radicals varies seasonally, with a maximum in summer in the troposphere [Warneck, 1988]. Honrath and Jaffe [1992] reported a well-defined seasonal cycle of NO_y, with a maximum (560-620 parts per trillion, by volume) in early spring and minimum (median 70 parts per trillion, by volume) in summer at Barrow, Alaska. The transition occurs sharply in April, when a sevenfold to eightfold decrease in NO_y concentration was observed. Davidson *et al.* [1993] also conclude that the minima in summer of atmospheric CO and CH₄ at Dye-3, Greenland, is caused in part by high concentrations of OH which are related to relatively high airborne concentrations of NO_x and O₃.

To explain the summer peak seen in nitrate concentration in polar snow we propose the following: During the winter, cold temperature and low insolation at middle to high latitudes favor long distance transport of PAN into polar regions. Further, since there is no sunlight during the polar winter, the secondary sources of NO_x and OH are limited. Under such conditions, snow falling during the polar winter has low nitrate concentrations due to a lack of NO_x and OH. With the appearance of sunlight and increasing atmospheric temperatures during the polar spring, the NO_x released from PAN reacts with OH to form HNO₃, yielding a summer nitrate peak in Greenland snow.

The less pronounced summer nitrate peaks at Mount Logan and Sentik Glacier may be a consequence of the types of air masses reaching these sites. Air masses reaching Mount Logan and Sentik Glacier are pristine free troposphere due to high elevations [Monaghan and Holdsworth, 1990; Mayewski *et al.*, 1993b; Wake *et al.*, 1993]. Additionally, these sites maintain some sunlight during winter so that the nitrate formed/released from PAN is deposited throughout the year.

As reported by *Finkel et al.* [1986] and *Whitlow et al.* [1992], our snow pit chemistry study around the GISP2 site reveals that there is sometimes a late winter/early spring nitrate peak in modern Greenland snow. Since this earlier peak has not been observed from our GISP2 detailed sections and *Finkel et al.*'s [1986] data before the year 1900 A.D., it may result from the input of anthropogenic material. In general, any NO_y species (NO , NO_2 , HO_2NO_2 , HNO_3 , the PANs, HNO_2 , N_2O_5 , NO_3 , and gaseous nitrates [*Singh*, 1987]) may end up as nitrate to be removed from the atmosphere. PAN concentrations are elevated in the polluted air [*Singh*, 1987; *Warneck*, 1988] which is transported and accumulated in "Arctic haze" during winter. PAN has also been reported to be a significant component of NO_y in the Arctic boundary layer during spring [*Barrie and Bottenheim*, 1991]. But NO_x released from PAN is restricted in the polar regions due to lower temperatures in late winter/early spring [*Singh et al.*, 1992b].

In addition to the role of NO_x and PANs, N_2O_5 chemistry may also play a role in the formation of the modern spring nitrate peak in Greenland snow since the low light to dark conditions favor the formation of N_2O_5 [*Calvert et al.*, 1985; *Leaitch et al.*, 1988]. There are two potential ways for N_2O_5 to yield nitrate spring peaks in snow. First, NO_2 is released from N_2O_5 due to an increased temperature in spring [*Atkinson et al.*, 1986], and it then reacts with OH to form HNO_3 . Second, N_2O_5 scavenges to HNO_3 on aqueous aerosols in late winter/spring [*Calvert et al.*, 1985; *Mozurkewich and Calvert*, 1988]. Although there are no N_2O_5 data sets available from the ambient Greenland atmosphere and we are unable to provide quantitative relationship between nitrate in snow and N_2O_5 in the atmosphere, these two processes may also be important in explaining the difference between the preanthropogenic nitrate summer peak and the anthropogenic era nitrate spring peak.

Conclusions

By examining the GISP2, 20D, Mount Logan, and Sentik Glacier ice core chemical species, we find that the mean annual concentrations of all chemical species are independent of the annual snow accumulation rate. Only the nitrate concentration data series reveal a normally distributed pattern. Based on previous work [*Neubauer and Heumann*, 1988; *Mayewski and Legrand*, 1990], we suggest that the cause of this pattern is post-depositional alternation. As a consequence, nitrate flux has a relatively strong linear relationship to the rate of snow accumulation at all sites studied.

The presence of a nitrate summer peak for the past several thousand years in Greenland snow is believed to be associated with the higher OH radical concentrations and an available NO_x source released from PAN in summertime.

The observed nitrate peak in early spring in modern (anthropogenic era) Greenland snow may be due to the buildup of NO_y as Arctic haze in the troposphere dur-

ing the long polar winter. NO_x released/formed from a component of NO_y (N_2O_5 , PAN, HO_2NO_2) reacting with OH to form HNO_3 in the atmosphere is believed to cause this peak.

Acknowledgments. We would like to acknowledge the assistance of the Polar Ice Coring Office (University of Alaska, Fairbanks) and the Greenland Ice Sheet Project 2 Science Management Office (University of New Hampshire). The comments of reviewers are greatly appreciated. Support for this work was provided by the National Science Foundation.

References

- Atkinson, R., A. M. Winer, and J. N. Pitts Jr., Estimation of night-time N_2O_5 concentrations from ambient NO_2 and NO_3 radical concentrations and the role of N_2O_5 in nighttime chemistry, *Atmos. Environ.*, **20**, 331-339, 1986.
- Barrie, L. A., Scavenging ratios, wet deposition, and in-cloud oxidation: An application to the oxides of sulphur and nitrogen, *J. Geophys. Res.*, **90**, 5789-5799, 1985.
- Barrie, L. A., and J. W. Bottenheim, Sulfur and nitrogen in the Arctic atmosphere, in *Pollution of the Arctic Atmosphere*, edited by W. Sturges, Elsevier, New York, 1991.
- Bottenheim, J. W., L. A. Barrie, and E. Atlas, The partitioning of nitrogen oxides in the lower Arctic troposphere during spring 1988, *J. Atmos. Chem.*, **17**, 15-27, 1993.
- Buck, C. F., P. A. Mayewski, M. J. Spencer, S. Whitlow, M. S. Twickler, and D. Barrett, Determination of major ions in snow and ice cores by ion chromatography, *J. Chromatogr.*, **594**, 225-228, 1992.
- Calvert, J. G., A. Lazrus, G. L. Kok, B. G. Heikes, J. G. Walega, J. L. Cantrell, and C. A. Cantrell, Chemical mechanisms of acid generation in the troposphere, *Nature*, **317**, 27-35, 1985.
- Cox, R. A., and M. J. Roffey, Thermal decomposition of peroxyacetylnitrate in the presence of nitric oxide, *Environ. Sci. Technol.*, **11**, 900-906, 1977.
- Dansgaard, W., and H. Oeschger, Past environmental long-term records from the Arctic, in *The Environmental Record in Glaciers and Ice Sheets*, edited by H. Oeschger and C. C. Langway Jr., pp. 287-318, John Wiley, New York, 1989.
- Davidson, C. I., Mechanisms of wet and dry deposition of atmospheric contaminants to snow surfaces, in *The Environmental Record in Glaciers and Ice Sheets*, edited by H. Oeschger and C. C. Langway Jr., pp. 29-51, John Wiley, New York, 1989.
- Davidson, C. I., J. R. Harrington, M. J. Stephenson, F. P. Small, F. P. Boscoe, and R. E. Grandly, Seasonal variations in sulfate, nitrate and chloride in the Greenland ice sheet: Relation to atmospheric concentrations, *Atmos. Environ.*, **23**, 2483-2493, 1988.
- Davidson, C. I., et al., Chemical constituents in the air and snow at Dye-3, Greenland, I, Seasonal variations, *Atmos. Environ.*, **27**(A), 2709-2722, 1993.
- Delmas, R. J., and M. Legrand, Long-term changes in the concentrations of major chemical compounds (soluble and insoluble), in *The Environmental Record in Glaciers and Ice Sheets*, edited by H. Oeschger and C. C. Langway Jr., pp. 319-342, John Wiley, New York, 1989.
- Ehhalt, D. H., F. Rohere, and A. Wahne, Sources and distribution of NO_x in the upper troposphere at northern midlatitudes, *J. Geophys. Res.*, **97**, 3725-3738, 1992.
- Finkel, R. C., C. C. Langway Jr., and H. B. Clausen, Changes in precipitation chemistry at Dye 3, Greenland, *J. Geophys. Res.*, **91**, 9849-9855, 1986.

- Herron, M., Impurity sources of F, Cl, NO₃ and SO₄²⁻ in Greenland and Antarctic precipitation, *J. Geophys. Res.*, **87**, 3052-3060, 1982.
- Holdsworth, G., and E. Peake, Acid content of snow from a mid-troposphere sampling site on Mount Logan, Yukon Territory, Canada, *Ann. Glaciol.*, **7**, 153-160, 1985.
- Honrath, R. E., and D. A. Jaffe, The seasonal cycle of nitrogen oxides in the Arctic troposphere at Barrow, Alaska, *J. Geophys. Res.*, **97**, 20,615-20,630, 1992.
- Leitch, W. R., W. Bottenheim, and J. W. Strapp, Possible contribution of N₂O₅ scavenging to HNO₃ observed in winter stratiform cloud, *J. Geophys. Res.*, **93**, 12,569-12,584, 1988.
- Legrand, M. R., and R. J. Delmas, Relative contributions of tropospheric and stratospheric sources to nitrate in Antarctic snow, *Tellus*, **38**(B), 236-249, 1986.
- Legrand, M. R., and S. Kirchner, Origins and variations of nitrate in south polar precipitation, *J. Geophys. Res.*, **95**, 3493-3507, 1990.
- Legrand, M., C. Lorius, N. I. Barkov, and V. N. Petrov, Vostok (Antarctica) ice core: Atmospheric chemistry changes over the last climatic cycle (160,000 years), *Atmos. Environ.*, **22**, 317-331, 1988.
- Legrand, M. R., F. Stordal, I. S. A. Isaksen, and B. Rognerud, A model study of the stratospheric budget of odd nitrogen including effects of solar cycle variations, *Tellus*, **41**, 413-426, 1989.
- Legrand, M., C. Feniet-Saigne, E. S. Saltzman, C. Germain, N. I. Barkov, and V. N. Petrov, Ice-core record of oceanic emissions of dimethylsulphide during the last climatic cycle, *Nature*, **350**, 144-146, 1991.
- Levy, H., J. D. Muhlman, and W. J. Moxim, A stratospheric source of reactive nitrogen in the unpolluted troposphere, *Geophys. Res. Lett.*, **7**, 441-444, 1980.
- Logan, J. A., Nitrogen oxides in the troposphere: Global and regional budgets, *J. Geophys. Res.*, **88**, 10,785-10,807, 1983.
- Looney, S. W., and T. R. Gullledge Jr., Use of the correlation coefficient with normal probability plots, *Am. Stat.*, **39**, 75-79, 1985.
- Lorius, C., G. Raisbeck, J. Jouzel, and D. Raynaud, Long-term environmental records from Antarctic ice cores, in *The Environmental Record in Glaciers and Ice Sheets*, edited by H. Oeschger and C. C. Langway Jr., pp. 343-362, John Wiley, New York, 1989.
- Lyons, W. B., and P. A. Mayewski, Nitrate plus nitrite concentrations in a Himalayan ice core, *Geophys. Res. Lett.*, **10**, 1160-1163, 1983.
- Lyons, W. B., P. A. Mayewski, M. J. Spencer, and M. S. Twickler, Nitrate concentrations in snow from remote areas: Implications for the global NO_x flux, *Biogeochem.*, **9**, 211-222, 1990.
- Mayewski, P. A., and M. R. Legrand, Recent increase in nitrate concentration of Antarctic snow, *Nature*, **346**, 258-260, 1990.
- Mayewski, P. A., W. B. Lyons, N. Ahmad, G. Smith, and M. Pourchet, Interpretation of the chemical and physical time-series retrieved from Sentik Glacier, Ladakh Himalayas, India, *J. Glaciol.*, **30**, 66-76, 1984.
- Mayewski, P. A., W. B. Lyons, M. J. Spencer, M. S. Twickler, W. Dansgaard, B. Koci, C. I. Davidson, and R. E. Honrath, Sulfate and nitrate concentration from a south Greenland ice core, *Science*, **232**, 975-977, 1986.
- Mayewski, P. A., M. J. Spencer, W. B. Lyons, and M. S. Twickler, Seasonal and spatial trends in south Greenland snow chemistry, *Atmos. Environ.*, **21**, 863-869, 1987.
- Mayewski, P. A., W. B. Lyons, M. J. Spencer, M. S. Twickler, C. F. Buck, and S. Whitlow, An ice-core record of atmospheric response to anthropogenic sulphate and nitrate, *Nature*, **346**, 554-556, 1990a.
- Mayewski, P. A., M. J. Spencer, M. S. Twickler, and S. Whitlow, A glaciochemical survey of the Summit region, Greenland, *Ann. Glaciol.*, **14**, 186-190, 1990b.
- Mayewski, P. A., L. D. Meeker, S. Whitlow, M. S. Twickler, M. C. Morrison, R. B. Alley, P. Bloomfield, and K. Taylor, The atmosphere during the Younger Dryas, *Science*, **261**, 195-197, 1993a.
- Mayewski, P. A., G. Holdsworth, M. J. Spencer, S. Whitlow, M. S. Twickler, M. C. Morrison, K. K. Ferland, and L. D. Meeker, Ice core sulfate from three Northern Hemisphere sites: Source and temperature forcing implications, *Atmos. Environ.*, **27A**, (17/18), 2915-2919, 1993b.
- Mayewski, P. A., et al., Changes in atmospheric circulation and ocean ice cover over the North Atlantic during the last 41,000 years, *Science*, **263**, 1747-1751, 1994.
- McElroy, M. B., Studies of polar ice: Insights for atmospheric chemistry, in *The Environmental Record in Glaciers and Ice Sheets*, edited by H. Oeschger and C. C. Langway Jr., pp. 363-377, John Wiley, New York, 1989.
- Minitab Reference Manual Release 9 (Windows), Minitab, Incorporated, State College, Penn., 1993.
- Monaghan, M. G., and G. Holdsworth, The origin of non-sea-salt sulphate in the Mount Logan ice core, *Nature*, **343**, 245-248, 1990.
- Mozurkewich, M., and J. G. Calvert, Reaction probability of N₂O₅ on aqueous aerosols, *J. Geophys. Res.*, **93**, 15,889-15,896, 1988.
- Mulvaney, R., and E. W. Wolff, Evidence for winter/spring denitrification of stratosphere in the nitrate record of Antarctic firn cores, *J. Geophys. Res.*, **98**, 5213-5220, 1993.
- Neftel, A., J. Beer, H. Oeschger, and F. Zurcher, Sulphate and nitrate concentrations in snow from south Greenland 1895-1978, *Nature*, **314**, 611-613, 1985.
- Neubauer, J., and K. G. Heumann, Nitrate trace determinations in snow and firn core samples of ice shelves at the Weddell sea, Antarctica, *Atmos. Environ.*, **22**, 537-545, 1988.
- Parker, B. C., and E. J. Zeller, Nitrogenous chemical composition of Antarctic ice and snow, *Antarctic J.*, **XIV**, 80-82, 1979.
- Russell, A. G., G. R. Cass, and J. H. Seinfeld, On some aspects of nighttime atmospheric chemistry, *Environ. Sci. Technol.*, **20**, 1167-1172, 1986.
- Singh, H. B., Reactive nitrogen in the troposphere, *Environ. Sci. Technol.*, **21**, 320-327, 1987.
- Singh, H. B., and P. L. Hanst, Peroxyacetyl nitrate (PAN) in the unpolluted atmosphere: An important reservoir for nitrogen oxides, *Geophys. Res. Lett.*, **8**, 941-944, 1981.
- Singh, H. B., and L. Salas, Peroxyacetyl nitrate in the free atmosphere, *Nature*, **302**, 326-328, 1983.
- Singh, H. B., D. Herlth, D. O'Hara, K. Zahnle, J. D. Bradshaw, S. T. Sandholm, R. Talbot, P. J. Crutzen, and M. Kanakidou, Relationship of peroxyacetyl nitrate to active and total odd nitrogen at northern high latitudes: Influence of reservoir species on NO_x and O₃, *J. Geophys. Res.*, **97**, 16,523-16,530, 1992a.
- Singh, H. B., D. O'Hara, D. Herlth, J. D. Bradshaw, S. T. Sandholm, G. L. Gregory, G. W. Sachse, D. R. Blake, P. J. Crutzen, and M. A. Kanakidou, Atmospheric measurements of peroxyacetyl nitrate and other organic nitrates at high latitudes: Possible sources and sinks, *J. Geophys. Res.*, **97**, 16511-16522, 1992b.
- Steffensen, J. P., Analysis of the seasonal variation in dust, Cl⁻, NO₃⁻, and SO₄²⁻ in two central Greenland firn cores, *Ann. Glaciol.*, **10**, 171-177, 1988.
- Wake, C. P., P. A. Mayewski, Z. C. Xie, P. Wang, and Z. Q. Li, Regional distribution of monsoon and desert dust signal record in Asian glacier, *Geophys. Res. Lett.*, **20**, 1411-1414, 1993.

- Warneck, P., *Chemistry of the Natural Atmosphere*, Academic, San Diego, Calif., 1988.
- Whitlow, S., P. A. Mayewski, and J. Dibb, A comparison of major chemical species seasonal concentration and accumulation at the South Pole and Summit, Greenland, *Atmos. Environ.*, 26(A), 2045-2054, 1992.
- Wolff, E. A., Nitrate in polar ice, paper presented at the *NATO ARW on Ice Core Studies of Global Biogeochemical Cycles*, edited by R. Delmas, Annecy, France, in press, 1995.
- Zeller, E. J., B. C. Dreschhoff, and C. M. Laird, Nitrate flux on the Ross Ice Shelf, Antarctica, and its relation to solar cosmic rays, *Gephys. Res. Lett.*, 13, 1264-1267, 1986.
- responding author), Glacier Research Group, Institute for the Study of Earth, Oceans and Space, University of New Hampshire, Durham, NH 03824. (e-mail: jed@unh.edu; p_mayewski@unh.edu; mcm@unh.edu; mst@unh.edu; siw@unh.edu; qinzhao.yang@grg.unh.edu)
- Robert Talbot, Atmospheric Chemistry Research Group, Institute for the Study of Earth, Oceans and Space, Morse Hall, University of New Hampshire, Durham, NH 03824. (e-mail: rwt@christa.unh.edu)
- Ernst Linder, Mathematics Department, Kingsbury Hall, University of New Hampshire, Durham, NH 03824. (e-mail: elinder@christa.unh.edu)

Jack Dibb, Paul A. Mayewski, Michael Morrison, Mark Twickler, Sallie Whitlow, and Qinzhao Yang (cor-

(Received April 25, 1994; revised October 19, 1994; accepted October 21, 1994.)