



3-31-2007

2006 Great Bay Organic Nitrogen (PON & DON) and Light Extinction (PAR) Monitoring Program

Jonathan Pennock
University of New Hampshire - Main Campus

Follow this and additional works at: <https://scholars.unh.edu/prep>



Part of the [Marine Biology Commons](#)

Recommended Citation

Pennock, Jonathan, "2006 Great Bay Organic Nitrogen (PON & DON) and Light Extinction (PAR) Monitoring Program" (2007). *PREP Reports & Publications*. 50.
<https://scholars.unh.edu/prep/50>

This Report is brought to you for free and open access by the Institute for the Study of Earth, Oceans, and Space (EOS) at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in PREP Reports & Publications by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

**2006 Great Bay
Organic Nitrogen (PON & DON) and
Light Extinction (PAR) Monitoring Program**

A Final Report to

The New Hampshire Estuaries Project

Submitted by

Dr. Jonathan Pennock
University of New Hampshire
Jackson Estuarine Laboratory
85 Adams Point Road
Durham, NH, 03824

March 31, 2007

This report was funded by a grant from the New Hampshire Estuaries Project, as authorized by the U.S. Environmental Protection Agency pursuant to Section 320 of the Clean Water Act.



Table of Contents

Introduction	2
Project Goals and Objectives	2
Methods	3
Results and Discussion	3
Conclusions and Recommendations	4
References	4
Appendix 1 (Meta-Data)	5
Appendix 2 (Data CD)	Inside Back Cover

Introduction

Nitrogen is most often considered to be the limiting nutrient for plant growth in marine waters. As a result, knowledge of nitrogen loading and ambient water-column concentrations are considered to be critical to understanding the response of aquatic ecosystems to nutrient over-enrichment—a process known as eutrophication when it results in the excess production of organic matter.

Plant production in many estuarine systems may also be limited by light availability as a result of high levels of turbidity in the water resulting from sediments, dissolved organic matter, and phytoplankton in the water column. Light limitation resulting from human-induced increases in turbidity is known to be particularly deleterious to seagrass production/distribution in some ecosystems and also play an important role in determining how phytoplankton respond to nutrient enrichment.

EPA is developing water quality criteria for estuaries that require knowledge of both total nitrogen and light availability (measured as photosynthetically active radiation, PAR). Through the National Estuarine Research Reserve (NERR) System-Wide Monitoring Program (SWMP), inorganic nutrient concentrations, chlorophyll-a concentration, and a number of hydrographic and water quality parameters are sampled on a monthly basis at 7 sites in the Great Bay system.

This project takes advantage of these existing monitoring activities to collect and analyze for particulate organic nitrogen (PON), dissolved organic nitrogen (DON) and photosynthetically active radiation (PAR) at existing sample sites in the New Hampshire seacoast region. When combined with existing dissolved inorganic nitrogen measurements, PON and DON allow the entire Total Nitrogen (TN) pool to be quantified. PAR measurements provide an estimate of the light availability in the system.

Project Goals and Objectives

UNH completed this project under contract to the NH Estuaries Project (Project ID #05-M-9). The project goals and objectives per the contract were to:

- (1) Conduct PON, DON, and PAR monitoring for the April – December sampling season at the Lamprey River (LR), Squamscott River (SQ), Chapman’s Landing (CL), Oyster River (OR), Great Bay (GB) and Coastal Marine Lab (CML).

The goal was to provide 216 measurements for PON and DON and 108 measurements of PAR during the sampling period. The final work product was agreed to be an Excel data file containing

hydrographic, PON, DON and PAR data for all of these sites.

Methods

The methods for this project followed the procedures specified in the approved QA Project Plan (Pennock and Trowbridge, 2003).

Results and Discussion

Overall, as shown in the table below, we were able to obtain 70 discrete estimates for the attenuation coefficient (PAR), 132 measurements for particulate nitrogen (PN) and 131 measurements of particulate carbon (PC), 214 measurements of total dissolved nitrogen (TDN) and 214 measurements of dissolved organic nitrogen (DON). The discrepancy between the number of PN/PC samples (132) and the goal (216) was because PN/PC was not measured in high tide samples from GRBCL, GRBCML, GRBLR, and GRBOR as planned in the work scope. Many PAR measurements were missed due to sensor malfunctions, ice, or shallow water.

We performed replicate analyses on all parameters except for PAR (triplicate PAR analyses was performed at 5 stations during the 2006 sampling period.) Laboratory analyses fell within the accepted guidelines detailed in the approved QA Project Plan (Pennock and Trowbridge, 2003).

Parameter	Location	Stations	Analyses	Mean	SD
PAR	All Data	107	70	-1.76	
PAR	QA	5	15	-2.16	0.25
POC	All Data	71	131	1.1119	
PON	All Data	71	132	0.1236	
TDN	All Data	107	214	0.4438	
DON	All Data	107	214	0.3189	

The CD included with this report contains data files in an Excel format that is consistent with NERR SWMP CDMO requirements. Included are the following required parameters for this project: Record #; Funding Source for Sampling Effort; Sample Date; Site/Station Name; Tidal Stage; Bottom Depth; Temperature; Salinity; Oxygen Concentration; Oxygen Percent Saturation; pH; Total Dissolved Nitrogen (TDN); Dissolved Organic Nitrogen (DON); Particulate Organic Carbon (POC); Particulate Organic Nitrogen (PON); and Attenuation Coefficient (K_d).

Please note that for analyses that were above zero and below the MDL, a value of one-half the MDL was inserted into the raw data file. Nutrient values that fall below prescribed MDL's have been changed to '-9999' in the processed data files, per NERR CDMO regulations. The attached CD contains an excel file with 4 worksheets (1. Raw data for all 2006 samples, 2. Processed data for all 2006 samples, 3. Processed data for relevant NHEP samples, 4. Processed data for relevant NERR SWMP samples.)

Differences between POC, PON and TDN replicate samples were compared to data quality objectives of 30% RPD and 0.5, 0.1 and 0.5 mg/L, respectively. Replicate samples that failed both tests should be rejected.

Data contained in this file are scheduled to be submitted to NERR CDMO by May 15th, 2007. It is subject to review and potential revision by the NERR CDMO.

Conclusions and Recommendations

The PON, DON and PAR monthly monitoring program provides important data on nitrogen concentration and light availability in the Great Bay estuary. When combined with the NERRS SWMP program, these data provide comprehensive coverage of the Great Bay estuary and allow total nitrogen concentrations to be calculated for use in nutrient criteria measurements.

References

Pennock and Trowbridge (2003) UNH Nutrient and Light Extinction Monitoring Program Quality Assurance Project Plan, Version 4, Final. University of New Hampshire and NH Estuaries Project, August 8, 2003.

2006 Organic Nitrogen (PON & DON) and Light Extinction (PAR) Monitoring Program Meta-Data (Appendix 1)

Research Methods

Monthly monitoring is conducted during ice-free seasons (generally April through December) as part of the GB NERR System-Wide Monitoring Program and National Coastal Assessment at the following locations:

Funding Source	Station Name	Station ID	Tide Stage	Lat Deg	Lat Minute	Long Deg	Long Minute
NERR	Adams Point	AP	L&H	43	5.495	70	51.821
NERR	Great Bay	GB	L	43	4.367	70	52.311
NERR	Lamprey River	LR	L&H	43	4.697	70	56.092
NERR	Oyster River	OR	L&H	43	8.400	70	54.100
NERR	Squamscott Railroad Bridge	SQ	L	43	3.182	70	54.754
NERR	Squamscott Chapman's Landing	CL	L&H	43	2.500	70	55.569
NERR	Coastal Marine Lab	CML	L&H	43	4.350	70	42.640

Samples are generally collected by small boat except for the Coastal Marine Lab site, which is accessed by vehicle. During certain periods of particularly stormy weather or when the channel markers are not present (generally early and late season), a subset of the normal stations may be sampled by vehicle; these cases are noted in the meta-data.

At each station, a hand-held YSI multi-probe is used to measure temperature, salinity, dissolved oxygen, and dissolved oxygen percent saturation in the surface water (~0.5m). Nutrient samples are collected in acid-washed and DI-water rinsed 1-liter HDPE bottles at ~0.5m depth, placed on ice in a cooler and processed upon return to the laboratory. Photosynthetically Active Radiation (PAR) profiles are made at as many stations as possible using a LiCor Quantum Irradiance Meter.

All nutrient processing and analysis methods and PAR profiling methods are detailed in the 2003 UNH Nutrient and Light Extinction Monitoring Program Quality Assurance Project Plan (Pennock & Trowbridge, 2003).

Deviations in Sampling Procedures for 2006

January

- Icing Conditions; only AP samples collected (no PAR).

February

- Icing Conditions; only AP samples collected (no PAR).

March

- Icing Conditions; only AP samples collected (no PAR).

April

- NERR samples collected by vehicle due to the lack of channel markers.
- Limited PAR measurements taken as a result of vehicle-based sampling.

May

- Limited vehicle-based sampling due to extreme flooding conditions.

June

- No PAR taken at ORL; too shallow.

July

- Many missing PAR values due to Licor instrument malfunction.

August

- Many missing PAR values due to Licor instrument malfunction.

September

- No PAR taken at OR (low tide); too shallow.

October

- No PAR taken at ORL, CLL, CLH; too shallow (CL accessed by vehicle).

November

- Vehicle-based sampling; limited PAR measurements as a result of shallow water.

December

- No PAR taken, rough sampling conditions.