University of New Hampshire

University of New Hampshire Scholars' Repository

Space Science Center

Institute for the Study of Earth, Oceans, and
Space (EOS)

2000

COMPTEL time-averaged all-sky point source analysis

W Collmar

Max-Planck-Institut für extraterrestrische Physik

V Schonfelder

Max-Planck-Institut für extraterrestrische Physik

A W. Strong

Max-Planck-Institut für extraterrestrische Physik

H Bloemen

Space Research Organization of the Netherlands

W Hermsen

Space Research Organization of the Netherlands (SRON)

See next page for additional authors

Follow this and additional works at: <https://scholars.unh.edu/ssc>

Part of the [Astrophysics and Astronomy Commons](#)

Recommended Citation

COMPTEL time-averaged all-sky point source analysis Collmar, W. and Schönfelder, V. and Strong, A. W. and Bloemen, H. and Hermsen, W. and McConnell, M. and Ryan, J. and Bennett, K., AIP Conference Proceedings, 510, 591-595 (2000), DOI:<http://dx.doi.org/10.1063/1.1303271>

This Conference Proceeding is brought to you for free and open access by the Institute for the Study of Earth, Oceans, and Space (EOS) at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Space Science Center by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

Authors

W Collmar, V Schonfelder, A W. Strong, H Bloemen, W Hermsen, Mark L. McConnell, James M. Ryan, and K Bennett

COMPTEL time-averaged all-sky point source analysis

W. Collmar, V. Schönfelder, A. W. Strong, H. Bloemen, W. Hermsen, M. McConnell, J. Ryan, and K. Bennett

Citation: [AIP Conference Proceedings](#) **510**, 591 (2000); doi: 10.1063/1.1303271

View online: <http://dx.doi.org/10.1063/1.1303271>

View Table of Contents: <http://scitation.aip.org/content/aip/proceeding/aipcp/510?ver=pdfcov>

Published by the [AIP Publishing](#)

Articles you may be interested in

[The spectral properties of the GRB prompt gammaray emission observed by the Suzaku Wideband Allsky Monitor](#)

AIP Conf. Proc. **1000**, 101 (2008); 10.1063/1.2943418

[AGILE sky exposure and sensitivity maps](#)

AIP Conf. Proc. **587**, 759 (2001); 10.1063/1.1419495

[All sky monitor for energetic \$\gamma\$ -rays](#)

AIP Conf. Proc. **558**, 594 (2001); 10.1063/1.1370831

[Preliminary results from a new analysis method for EGRET data](#)

AIP Conf. Proc. **510**, 606 (2000); 10.1063/1.1303274

[Improved COMPTEL maps of the milky way](#)

AIP Conf. Proc. **510**, 586 (2000); 10.1063/1.1303270

COMPTEL Time-Averaged All-Sky Point Source Analysis

W. Collmar*, V. Schönfelder*, A.W. Strong*, H. Bloemen⁺,
W. Hermsen⁺, M. McConnell[†], J. Ryan[†], K. Bennett[@]

*Max-Planck-Institut für extraterrestrische Physik, Postfach 1603, 85740 Garching, Germany

⁺SRON-Utrecht, Sorbonnelaan 2, NL-3584 CA Utrecht, The Netherlands

[†]University of New Hampshire, Durham NH 03824-3525, USA

[@]Astrophysics Division, ESTEC, NL-2200 AG Noordwijk, The Netherlands

Abstract. We use all COMPTEL data from the beginning of the CGRO mission (April '91) up to the end of CGRO Cycle 6 (November '97) to carry out all-sky point source analyses in the four standard COMPTEL energy bands for different time periods. We apply our standard maximum-likelihood method to generate all-sky significance and flux maps for point sources by subtracting off the diffuse emission components via model fitting. In addition, fluxes of known sources have been determined for individual CGRO Phases/Cycles to generate lightcurves with a time resolution of the order of one year. The goal of the analysis is to derive quantitative results – significances, fluxes, light curves – of our brightest and most significant sources such as 3C 273, and to search for additional new COMPTEL sources, showing up in time-averaged maps only.

INTRODUCTION

The imaging COMPTEL experiment aboard CGRO is the pioneering satellite experiment of the MeV-sky ($\sim 1 - 30$ MeV). For a detailed description of COMPTEL see [1]. One of COMPTEL's prime goals is the generation of all-sky maps, which provide a summary on the MeV-sky in total. This goal has been achieved by e.g. [2], [3] who generated maximum-entropy all-sky images and by [4], who generated the first COMPTEL all-sky maximum-likelihood maps, which – compared to maximum-entropy ones – have the advantage of providing quantitative results like significances and fluxes of source features. Here we present all-sky maximum-likelihood maps from which models of the diffuse emission have been removed. Our emphasis is on AGN. For a discussion on the method see [5] in these proceedings.

The main analysis goals are 1) to derive a summary of known COMPTEL point sources, 2) to search for further point sources, 3) to derive time-averaged quantitative parameters ('first order') of our brightest point sources, i.e., significances, fluxes, MeV-spectra, and possible time variability, and 4) to further investigate our data and analysis methods.

CP510, *The Fifth Compton Symposium*, edited by M. L. McConnell and J. M. Ryan
© 2000 American Institute of Physics 1-56396-932-7/00/\$17.00

DATA AND ANALYSIS METHOD

Using all data from the beginning of the CGRO mission (April '91) to the end of CGRO Cycle VI (Nov. '97), we generated a consistent database of relevant COMPTEL data sets (events, exposure, geometry) for individual CGRO viewing periods (VPs) in the 4 standard energy bands (0.75-1, 1-3, 3-10, 10-30 MeV) in galactic coordinates by applying consistent data selections. This database was supplemented by relevant data sets containing models describing the galactic diffuse γ -ray emission (HI, CO, and inverse-Compton components) and the isotropic extragalactic γ -ray background emission. To check for time variability of γ -ray sources these data sets were combined for different time periods: the six individual CGRO Phases/Cycles, the sum of all data (CGRO Phases I-VI; April '91 - Nov. '97) as well as the first (CGRO Phases I-III; April '91 - Oct. '94) and the second half (CGRO Phases IV-VI; Oct. '94 - Nov. '97). Each set of all-sky data is analysed by our standard maximum-likelihood method which simultaneously 'handles' individual VPs, generates, iteratively, a background model (see [6]), and finally generates significance and flux maps and/or significances and fluxes for individual sources. Because we are interested in point sources, the diffuse emission is always removed in the fitting procedure (e.g. Figure 1). For the derivation of the source fluxes (see Figure 2 as an example), the point sources of interest (e.g. 3C 273, Cyg X-1) have additionally been included in the fitting procedure. We like to mention however, that the results derived by such all-sky fits should be considered correct to first order only. To derive final/optimal results for a particular source, a dedicated analysis has to be carried out, which e.g. makes several cross checks by applying different background models and would take into account the presence of other source features in the region of interest. Also, along the galactic plane the results depend on the 'goodness' of the applied diffuse emission models for the MeV-band.

RESULTS

The significance maps in Figure 1, which contain all data of the first 6.5 years of the COMPTEL mission, are the first COMPTEL all-sky point source maps in the continuum bands. They provide a summary of the on-average brightest and most significant MeV-sources. Similar maps focussing on the Galactic plane only are given elsewhere in these proceedings ([5]). The Crab – for display reasons removed in all maps of Figure 1 – is by far the most significant COMPTEL point source. In the 1-3 MeV band for example it reaches a significance of $\sim 110\sigma$ (i.e. a likelihood ratio of ~ 12000) for the CGRO Phase I-VI period. With significances of $\sim 11\sigma$, $\sim 10\sigma$, and $\sim 6\sigma$ in the 1-3, 3-10, and 10-30 MeV bands is the quasar 3C 273 found to be on average the second most significant point source. Its fluxes in these bands are between 10% and 15% of the Crab flux. Several other extragalactic (e.g. 3C 279, PKS 0528+134, Cen A) and galactic (e.g. Cyg X-1, PSR 1509-58, a known but unidentified source at l;b: $18^\circ; 0^\circ$) sources are visible as well. In addition there

FIGURE 1. COMPTEL time-averaged maximum-likelihood point source all-sky maps in the 1-3 (upper), 3-10 (middle), and 10-30 MeV (lower panel) energy bands for the time period April '91 to November '97, i.e. CGRO phases I-VI. The galactic and extragalactic diffuse emission as well as the emission from the Crab have been subtracted off via model fitting. For the 1-3 MeV and the 3-10 MeV significance maps the contour lines start at a likelihood ratio value of 12 ($\sim 3.5\sigma$ for a known source; χ^2_1 -statistics) and for the 10-30 MeV map at a likelihood ratio value of 9 (3.0σ for a known source) with steps of 6 for all maps. The most significant source features are labeled.

FIGURE 2. Left: COMPTEL light curves of the quasar 3C 273 in the 4 standard bands as derived from all-sky flux fitting. Each flux point is averaged over an individual CGRO phase. The error bars are 1σ and the upper limits are 2σ . Right: COMPTEL MeV spectra of 3C 273 for the sum of all CGRO phase I-III (April '91 - Oct. '94), IV-VI (Oct. '94 - Nov. '97), and I-VI (April '91 - Nov. '97) data as derived from all-sky flux fitting. The error bars are 1σ and the upper limits are 2σ .

are indications for previously unknown source features like at l;b: $75^\circ; +65^\circ$ in the 1-3 MeV map and at l;b: $85^\circ; -65^\circ$ in the 3-10 MeV map for example. Such spots are promising candidates for further dedicated analyses. This time-averaged approach suppresses sources which flare up only in short time periods. Therefore the maps show fewer sources than are listed in the COMPTEL source catalog (see [7]).

For all bright and significant COMPTEL sources we have derived fluxes in our 4 standard energy bands for the different time periods mentioned above, and have combined them to MeV light curves and spectra. Some results for 3C 273 are shown as an example in Figure 2. In the 1-10 MeV energy band 3C 273 is detected in each CGRO Phase/Cycle i.e. in time periods of typically 1 year. The flux turns out to be rather stable and varies only within a factor of ~ 2 in the 1-3 MeV and within a factor of ~ 4 in the 3-10 MeV energy band. The spectra show the same trend. Whereas the flux below 3 MeV turns out to be same for both halves, there is an indication that at the upper COMPTEL energies (>3 MeV) the source was brighter during the second half. All three spectra clearly show the spectral turnover

TABLE 1. Detection significances (σ) in 3 different energy bands of some known COMPTEL AGN sources (plus Crab) as derived by this all-sky analysis for the sum of data of the first 6.5 years of the COMPTEL mission.

γ -ray Source	1-3 MeV	3-10 MeV	10-30 MeV	Source Type
Crab	~ 110	~ 76	~ 44	pulsar + nebula
3C 273	11.5	10.8	6.2	FSRQ ^a
3C 279	5.3	5.7	4.9	FSRQ
PKS 0528+134	— ^b	3.0	6.0	FSRQ
PKS 1622-297	<3	<3	4.1	FSRQ
3C 454.3	<3	5.3	<3	FSRQ
Cen A	5.7	5.3	3.0	Radio Galaxy

^a Flat-Spectrum Radio Quasar

^b Unknown due to uncertainties in removing the strong Crab signal

occurring at MeV-energies. However, we emphasize that for final conclusions a dedicated source analysis has to be carried out.

SUMMARY

We have applied the maximum-likelihood method to COMPTEL all-sky data of different time periods. By simultaneously fitting models for the different diffuse emission components, this analysis method provided quantitative all-sky results – significances and fluxes – on point sources. An all-sky summary on their time-averaged fluxes and significances is thereby provided. After the Crab – pulsar plus nebula – the quasar 3C 273 was found to be the most significant COMPTEL MeV-source, having time-averaged fluxes of the order of 10% to 15% of the Crab. Additional evidence for previously unknown source features has been found as well.

ACKNOWLEDGMENTS: The COMPTEL project is supported by the German government through DARA grant 50 QV 9096 8, by NASA under contract NAS5-26645, and by the Netherlands Organisation for Scientific Research (NWO).

REFERENCES

1. Schönfelder, V., et al., *ApJS* **86**, 657 (1993).
2. Strong, A.W., et al., *Proc. 2nd INTEGRAL Workshop, ESA SP-382*, 533 (1997).
3. Bloemen, H., et al., *Proc. 3rd INTEGRAL Workshop*, in press (1999).
4. Blom, J.J., et al., *COMPTEL High-Latitude Gamma-Ray Sources (Ph.D. Thesis)*, ISBN 90-9010945-5, (1997).
5. Bloemen, H., et al., these proceedings.
6. Bloemen, H., et al., *ApJS* **92**, 419 (1994).
7. Schönfelder, V., et al., *ApJS* submitted (1999).