

Pleistocene History and Glacio-tectonic Features  
in the Lac Mégantic Region, Québec<sup>1</sup>

by

William W. Shilts  
Geological Survey of Canada

The road log which follows provides detailed descriptions of field localities which define the glacial stratigraphy of the Lac-Mégantic area. Evidence for three glaciations separated by two nonglacial intervals includes three stratigraphically distinct tills, lacustrine deposits related to Glacial Lake Gayhurst, a complex sequence of lake sediments and fine gravels (Massawippi Formation), late-glacial recessional moraines, and striae indicating at least two different ice-flow directions.

In addition, large-scale slump blocks involving thick sections of lacustrine sediments and till, and complex deformation structures in ice-contact sediments will be examined.

<sup>1</sup>Publication authorized by the Director, Geological Survey of Canada.






Figure 1



Quaternary Stratigraphic Column, Southeastern Quebec

Age                      Unit                      Chronologic Control

Late Wisconsin	post-Lennoxville sediments	12,640 ± 190 (GSC-312; peat) 12,570 ± 220 (GSC-419; peat) 11,500 ± 160 (GSC-475-2; marine shells) 9,180 ± 180 (GSC-856; wood in colluvium) 12,000 ± 230 (GSC-936; marine shells) 11,200 ± 200 (GSC-1248; peat) 11,000 ± 240 (GSC-1289; peat) 11,200 ± 160 (GSC-1294; peat) 14,900 ± 220 (GSC-1339; peat)
	Lennoxville Till	
Middle Wisconsin	Gayhurst Formation	>20,000 B.P. (GSC-1137) ca. 4000 varves
	Chaudière Till	
Early Wisconsin	Massawippi Formation	>54,000 B.P. (Y-1683) >41,500 B.P. (GSC-507) >40,000 B.P. (GSC-1084)
	Johnville Till	
Sangamon and/or Illinoian	pre-Johnville sediments	



Stop DescriptionsStop #1: Grande Coulée Section

Lennoxville Till; 5 m - 7 m; At east end of the section till is sandy and silty and oxidized to its base; sandy facies pinches out on the west side of the section and is represented by a compact, grey, unoxidized till facies typical of most exposures of Lennoxville Till in the region; sandy facies may represent an end moraine deposited in the proglacial lake which abutted the Lennoxville glacier during its retreat down the Chaudière valley; fabrics indicate ice flow from N to N 40°W.

Gayhurst Formation (?); 7 m thick; Interlaminated sand, silt, and clay, tentatively correlated with Gayhurst Formation sediments occurring farther south in the Chaudière valley, on the basis of stratigraphic position beneath Lennoxville Till; Chaudière Till, which underlies Gayhurst Formation sediments elsewhere, has not been identified in this section; note alternating oxidized and unoxidized beds throughout the section.

Massawippi Formation; 0.4 m thick; Fissile, structureless sand contains finely divided plant fragments dated >40,000 B.P (GSC-1084); Massawippi Formation contains pollen (predominantly Picea and Pinus) which suggest presence of a northern boreal forest at its time of deposition; i.e. a climate cooler than present; Massawippi Formation is exposed only in a lens at the west end of the section where it rests on a diamicton about 0.3 m thick. This diamicton can be traced to the east end of the section at about the same altitude and is used as a marker bed to clarify the relationships of older and younger stratigraphic units to the Massawippi Formation.

Johnville Till; 1.6 m thick; Compact, non-calcareous, grey till with strong fabric from N 40° W; only occurs in the central part of the section where it is directly overlain by the diamicton which underlies the Massawippi Formation; Johnville Till is thought to be of early Wisconsin age and correlative with the Bécancour Till of the St. Lawrence Lowlands.

Pre-Johnville Sediments;

A) Fluvial Gravel; 3 m thick; Coarse, massive gravel with strong stone imbrication indicating eastward current; gravel clasts are thickly coated with iron oxide at the west end of the section; unit can be traced from the west to east end of the section; overlain by the diamicton and Massawippi Formation at the west end of the section, directly by Johnville Till in the center, and by the diamicton and Gayhurst Formation (?) at the east.

B) Laminated silt and clay; 1.6 m thick; Possible glacial-lake sediment; non-calcareous throughout and oxidized to within 30 cm of its base; traceable with the same thickness and altitude throughout the section.



C) Fluvial Gravel; 1.3 m thick; Similar to (A) but clasts are not coated with iron oxide and are finer; rare erratics of gneiss from the Canadian Shield (70 mi to the north) found in this gravel, indicating derivation from a glaciated terrain.

D) Coarse sand grading downward into fine sand with clay partings; >4.4 m; May be a lacustrine or flood-plain sediment; unoxidized fine sand below river level contains finely-divided plant debris.

The pre-Johnville sediments are the oldest sediments recognized in southeastern Quebec and are tentatively assigned a pre-Wisconsin age. They may be the most complete record yet described of pre-Bécancour-Johnville deposition east of Toronto.

#### Stop #2: Chaudière River Section north of St. Martin-de-Beauce

Lennoxville Till; 18 m thick; Compact, grey, fissile, calcareous, pebbly, silty till; oxidized upward 5 m from base and downward 3 m from top; selenite crystals, apparently formed during breakdown of pyrite and calcite, were found near the base of the unit; fabrics indicate that the Lennoxville glacier was flowing from N 60° W - N 80° W during deposition of entire unit.

Lennoxville Till, sandy facies; 4.5 m thick; Sandy, loose, gravelly, oxidized till; similar to sandy tills of New England; texture is probably partially a result of incorporation of underlying sandy lake sediments; bottom contact irregular due to thrust faults which have carried lake sediment into the till.

Gayhurst Formation (?); 7 m thick; Interlaminated fine sand and clay; upper portion of the sediment is disturbed and sheared into overlying sandy till.

Chaudière Till; >1m thick; Grey, hard, calcareous, sparingly stony till with shear structures; fabric suggests that depositing glacier moved from about N 20° E.

#### Stop #3: Samson River Section

The area traversed to reach the Samson River section is on the axis of the indicator ribbon of ultrabasic rocks which extends southeast from the Thetford Mines area. This indicator train is very narrow and does not have the classic "fan" shape of indicator trains described in New England. Boulder piles on the field at the start of the traverse have 10% and 14% ultrabasic cobbles, but ultrabasic frequency drops to less than 2% 6 miles north and south of the field (the width of the source area outcrop is 14 miles).



The petrology of the upper tills at the Samson River section has been studied in detail. 155 till samples from a 10 m x 4 m vertical face have had complete textural analyses (on particles <6 mm diameter); clay-mineral analyses, calcite/dolomite (Chittick) analyses, and <2 $\mu$  carbonate (X-ray) analyses. The weight percents of magnetite and total heavy minerals in the fine-sand fractions, and trace element concentrations (Ni, Cr, Zr, Ti, Cu, V) in the <63 $\mu$  fraction have been calculated for most samples. The purpose of this exercise was to evaluate the vertical and lateral variations of these parameters in relation to primary (sedimentation) and secondary (weathering) processes. Preliminary results of this study will be discussed at the section.

- Stratigraphy: -

Lennoxville Till

A. Till; 1 m to 3 m thick; Sandy, compact, pebbly till, oxidized to base; includes lenses of lake sediment sheared up from beds immediately at base; fabric and high ultrabasic pebble frequencies indicate that this unit was deposited by a glacier flowing from N 50° W - N 80° W.

B. Lake Sediment; 0 m to 1.6 m thick; Highly contorted and sheared, interlaminated fine sand and clay; oxidized where less than 0.5 m thick.

C. Till; 5 m thick; Compact, grey, pebbly, silty till with a boulder pavement at the top; fabrics and pebble lithologies indicate the unit was deposited by a glacier with flow from N 20° W - N 30° W; oxidation extends along joints almost to the base; underlying deformed, clayey lake sediments grade gradually upward into till.

Gayhurst Formation; 13 m thick; Grey, calcareous, interlaminated silt and clay; contains several 0.3 m- to 1.0 m-thick zones of massive silt-clay or stony till-like sediment; the stony bands contain convolute laminations, and thin, horizontally bedded silt and clay laminae above and below them are undisturbed; the massive zones are interpreted as "turbidity" current deposits resulting from slumping from basin sides or an ice front or subaqueous slumping of sediments already deposited on the lake floor.

Chaudière Till; 1.5 m thick; Grey, calcareous, compact, clayey till; although the fabric is similar to those in the higher Lennoxville Tills (NW) the unit contains no ultrabasic pebbles; clayey texture caused by incorporation of clayey lake sediments which crop out at the base of the section; till includes clasts of these sediments and has a maroon tinge which is typical of the lake sediments.



## Pre-Chaudière Sediments

A. Fine Gravel; 3 m thick; Fine gravel with large-scale crossbedding dipping south (opposite present river flow); gravel contains 5 cm to 10 cm rounded clasts of underlying lacustrine laminated silt and clay.

B. Lake Sediment; >1 m thick; Interlaminated, calcareous, silt and clay; clay laminae are chocolate brown to maroon at this locality; this unit crops out at several places just above and below river level along the Samson River.

Approximately 0.25 miles upstream from the section a diapir of the lowest (maroon) silt and clay can be seen intruding the overlying fine gravel. The origin of this structure is not known, but it illustrates the high plasticity of clay and silt in the Lac-Mégantic area.

### Stop #4: Gayhurst Dam Borrow Pit

The section described here is a composite of the borrow pit, the deep gully on the far (east) side of the river, and a borehole drilled to bedrock just north of the gully. The stratigraphy in the borehole was determined by taking split-tube and Shelby tube samples at 1 m intervals and at changes in lithology. The three sites serve as the type section of the Gayhurst Formation.

#### - Stratigraphy: -

Lennoxville Till; ~10 m thick; Clay-till member of Lennoxville Till; compact, grey, calcareous clayey till with less than 10% sand and coarser fragments; note that there are no boulders in the till and that the till pebbles include no granodiorite component; the ground surface behind the section is mantled with large, predominantly granodiorite boulders - a one-boulder-thick ablation deposit; Fabric is N 20° E.

## Gayhurst Formation

A. Upper Member; 6.2 m thick; Calcareous, laminated silt and clay; bottom 0.3 m is contorted with ball and pillow structures and convolute lamination suggesting subaqueous slumping; about 600 graded couplets are present in this unit.

B. Middle Member; 22 m thick; Fine sand grading upward to coarse sand and fine gravel with large-scale crossbedding; the top of this unit (altitude = 369 m a.s.l.) is thought to approximate the altitude of the water surface of the low level phase of Glacial Lake Gayhurst at this site; the upper contact can be traced at approximately 370 m several miles downstream from here.



C. Lower Member; 59 m thick; Fine-grained lacustrine sediment; contains approximately 3400 graded, calcareous silt and clay couplets; upper 7 m and lower 10 m are very fine sand and silt; graded couplets contain sparse, finely disseminated plant fragments; several zones show evidence of subaqueous slumping.

Chaudière Till; 11.8 m thick; Two grey, compact, calcareous, silty, sandy till units separated by 2.4 m of highly deformed, brecciated silt and clay laminae; upper part of till derived from northwest; lower part derived from north or northeast.

#### Pre-Chaudière Sediments

- A) Lake sediment; 2.4 m thick; intensely deformed laminated silt and clay.
- B) Fine Gravel; 1.2 m thick; fine gravel with abundant clasts derived from sources to northwest.
- C) Johnville (?) Till; 2.7 m thick; stony, compact sediment; no sample recovered.
- D), Lake sediment (?); 1.8 m thick; silt and clay; no sample recovered.
- E) Weathered bedrock (?); 1.8 m thick; no sample recovered.
- F) Pyritiferous black slate (Compton Fm., Devonian (?)); 2.8 m of core recovered.


#### Stop #5: Drolet Slump Blocks

The asymmetric ridges on which we will eat lunch are thought to be large, coherent slump blocks composed of Gayhurst Formation sediments capped by the clayey lentil of Lennoxville Till. The blocks have dropped as much as 50 m vertically along normal fault planes, apparently in response to erosion by the Chaudière River. Although rotational slumping is quite common everywhere in the Lac-Mégantic region, these blocks are unusual for their size (more than 1 mile long) and location more than two miles from the site of river erosion.

The Drolet River occupies a broad, deep bedrock depression (see accompanying figures).



# Rivière Drolet, P.Q.: Cross Sections



Diagrammatic Cross Section  
of Rotational Slump:  
Rivière Chaudière, P. Q.




Figure 2



Stop #6: Mégantic Moraine Complex

The smooth ridges at this locality are thought to be till moraines deposited during a halt of the Lennoxville glacier. The till exposed in the cut through the moraine ridge is quite sandy and loose and the complex grades into ice contact stratified deposits at lower altitudes, several tens of meters to the northwest.

Smooth till or gravel ridges (usually nested) are common moraine forms in the Lac-Mégantic region, but they are generally less than a mile long. It is not known whether the present ridge form is entirely a product of ice deposition or whether the form was accentuated by meltwater flowing between ridges after their formation.

Ice-front positions during retreat of the Lennoxville glacier have been drawn by connecting linear trends of assemblages of ice-front features such as till moraine ridges, ice-contact subaerial outwash, ice-contact deltaic sediments, and meltwater channels.

Stop #7: Striae West of Lac-Mégantic

The striae on this outcrop of interbedded Compton Formation slate and sandstone reflect two directions of movement for the Lennoxville glacier. The "tails" of rock in the lee of pyrite cubes indicate the sense of movement - toward the Boundary Mountains (SE).

Outcrops in the Lac-Mégantic region commonly show two or more distinct azimuth groupings of striae with very few intervening directions.

Note the "offset" striae on this outcrop; one part of the outcrop has been uplifted a couple of centimeters above the other after the striae were cut. Such offsetting is common in this region. Failure is along bedding planes in the slate. On the basis of such evidence, Oliver, Johnson and Dorman (1970) have suggested that the cumulative postglacial displacement along such faults may have been "substantial".

Stop #8: Woburn Esker Gravel Pit

This pit has exposed in the past an excellent assemblage of structural features resulting from melting of buried ice. The pit face exposes two main areas of slumping over melting ice. Note that the faults formed in response to subsidence are high-angle reverse faults with small compensating normal faults. The significance of this observation will be discussed briefly.

This stop is about  $1\frac{1}{2}$  to 2 miles from the Coburn Gore customs houses. The field trip ends here and those travelling to Rangely-Stratton need only follow route 34 east to the border where it joins the Arnold Trail (rte. 27) which passes through Rangely and Stratton. Drive very carefully as this road is tortuous, narrow, and heavily travelled by logging trucks.



References

- Behling, R.E., 1965, A detailed study of the Wisconsin stratigraphic section of the upper Lamoille vally, north central Vermont: unpub. M.S. thesis, Miami University, Oxford, Ohio, 125 p.
- Borns, H.W., Jr., 1966, An end-moraine complex in southeastern Maine: Geol. Soc. America, Northeastern Sec. Ann. Mtg., Philadelphia, p. 13-14.
- Borns, H.W., 1963, Preliminary report on the age and distribution of the late Pleistocene ice in north central Maine: Amer. J. Sci., v. 261, pp. 738-740.
- Brady, J.G. and Dean, R.S., 1966, The composition and properties of ceramic clays and shales of Québec; Dept. of Energy, Mines and Resources, Mines Branch, Ottawa, Research Report, R 187, 107 p.
- Cannon, William F., 1964, The petrography of the tills of northern Vermont: unpub. M.S. Thesis, Miami University, Oxford, Ohio.
- Chalmers, R., 1898, Surface geology and auriferous deposits of southeastern Quebec: Geol. Surv. of Canada, Ann. Report, vol. 10, pt. V., 160 p.
- Cooke, H.C., 1937, Further note on northward moving ice: Am. Jour. Sci., 5th ser., v. 34, p. 221.
- Duquette, G., 1960, Preliminary report on Gould area: Quebec Dept. Mines, Prelim. Rept. 432, 10 p.
- Flint, R.F., 1951, Highland centers of former glacial outflow in northeastern North America: Geol. Soc. America Bull., v. 62, p. 21-38.
- Gadd, N.R., 1960, Surficial geology of the Bécancour map-area, Quebec: Geol. Surv. Canada, Paper 59-8, 34 p.
- Gadd, N.R., 1964, Surficial geology, Beauceville Map-area, Quebec: Geol. Surv. Canada, paper 64-12.
- Gadd, N.R., 1964, Moraines in the Appalachian region of Quebec: Bull. Geol. Soc. Amer., vol. 75, pp. 1249-1254.
- Gadd, N.R., 1965, Surficial geology, Chaudière River valley: Geol. Survey of Canada, Paper 65-1, pp. 115-117.
- Gadd, N.R., 1965, Buried valleys and gold placers, Beauce County: Geol. Surv. Canada, Paper 65-1, pp. 117-119.



- Gadd, N.R., 1967, The Pattern of Glacial Recession in Southeastern Québec: (Abs.), Northeast section of the Geol. Soc. of America 1967, Annual meeting, pp. 28-29.
- Gadd, N.R., (in press), Pleistocene geology of the Central St. Lawrence Lowland: Geol. Surv. Canada, Memoir 359.
- Gorman, W.A., 1955, Preliminary report on St. Georges-St. Zacharie area: Quebec Dept. Mines, Prelim. Rept. 314, 5 p.
- Lasalle, P., 1966, Late Quaternary vegetation and glacial history in the St. Lawrence Lowlands, Canada: Leidse Geol. Mededel., pt. 38, p. 91-128.
- McDonald, B.C., 1966, Pleistocene geology studies, Richmond-Sherbrooke region, southeastern Québec: Geol. Surv. of Canada, Paper 66-1, p. 167.
- McDonald, B.C., 1966, Surficial geology, Richmond-Dudswell, Quebec: Geol. Surv. Canada, Map 4-1966.
- McDonald, B.C., 1966, Auriferous till in the Eastern Townships, southeastern Quebec: Geol. Surv. Canada, Paper 66-2, p. 51-54.
- McDonald, B.C., 1967, Pleistocene events and chronology in the Appalachian region of southeastern Quebec, Canada: Ph.D. dissert., Yale Univ., New Haven, 161 p.
- McDonald, B.C. 1967, Surficial geology, Sherbrooke-Orford-Memphremagog, Quebec: Geol. Surv. Canada, Map 5-1966.
- McDonald, B.C. 1967, Wisconsin stratigraphy and ice-movement directions in southeastern Quebec, Canada, (abs.): Geol. Soc. America, Northeastern Sec. Ann. Mtg., Boston, p. 41-42.
- McDonald, B.C., 1967, Glacial geology of Eaton River watershed and adjacent areas of southeastern Quebec: Geol. Surv. of Canada Paper 67-1, pp. 166-167.
- McDonald, B.C., 1968, Deglaciation and differential postglacial rebound in the Appalachian region of southeastern Quebec: Jour. Geology, V. 76, p. 664-677.
- McDonald, B.C. 1968, Composition of some garnets used in provenance studies of tills; Jour. Sed. Pet., vol. 38, no. 3, pp. 956-957.
- McDonald, B.C. 1969, Surficial geology of La Patrie-Sherbrooke area, Quebec, including Eaton River watershed: Geol. Surv. Canada, Paper 67-52, 21 p.



- McDonald, B.C. and Shilts, W.W., (in press) Quaternary events and stratigraphy, southeastern Quebec: Geol. Soc. America Bull.
- McGerrigle, H.W., 1936; Gold placer deposits of the Eastern Townships: Que. Bur. Mines, Annual Rept. for 1935, pt. E.
- Marleau, R.A., 1968, Woburn-East Mégantic-Armstrong area, Frontenac and Beauce counties: Quebec Dept. Nat. Res., Geol. Rept. No. 131, 55 p.
- Oliver, J., Johnson, T., and Dorman, J., 1970, Postglacial faulting and seismicity in New York and Quebec: Can. Jour. Earth Sci., V. 7, no. 2, pt. 2, p. 579-590.
- Prest, V.K., Grant, D.R., and Rampton, V.N., 1967, Glacial Map of Canada, Geol. Surv. of Canada, Map 1253A.
- Shilts, W.W., 1967, X-Ray diffraction as a stratigraphic tool in the Pleistocene of Northern Vermont and southeastern Québec: (Abs.) Program, Geol. Assn. of Canada International Meetings, Kingston, Ontario, 1967.
- Shilts, W.W., 1968, Pleistocene geology, Lake Mégantic area, southeastern Québec: Geol. Surv. of Canada, Paper 68-1, p. 123.
- Shilts, W.W., 1969, Quaternary geology of the upper Chaudière River drainage basin, Quebec: Geol. Surv. Canada, Paper 69-1A, p. 218-220.
- Shilts, W.W., 1969, Pleistocene geology of the Lac Mégantic region, southeastern Quebec, Canada: Ph. D. dissert., Syracuse Univ., Syracuse, 154 p.
- Shilts, W.W., 1970, Indicator studies; Pleistocene geology; Lac Mégantic region, Quebec: Geol. Surv. Canada, Paper 70-1A, p. 185-186.
- Shilts, W.W., (in press) Geochemical anomalies in till; southeastern Quebec: Geol. Surv. Canada, Paper 70-1B,
- Stewart, David, P., 1961, The glacial geology of Vermont: Vermont Geol. Survey, Bull. No. 19, 124 p.
- Stewart, D.P. and MacClintock, P., 1964, The Wisconsin stratigraphy of northern Vermont: Am. Jour. Sci., v. 262, p. 1089-1097.
- Stewart, D.P., and MacClintock, P., 1969, The surficial geology and Pleistocene history of Vermont: Vermont Geol. Surv., Bull. 31, 251 p.



- Terasmae, J., 1958, Non-glacial deposits in the St. Lawrence Lowlands, Quebec; Geol. Surv. of Canada, Bull. 46, contributions to Canadian Palynology, Part II, pp. 13-28.
- Terasmae, J., 1965, Geological Survey palynological studies: Geol. Surv. Canada, Paper 65-1, p. 158-159.
- Thornes, J.B., 1965, Some observations on the late-glacial stages in the Coaticook valley, southern Quebec: Cahiers de Géog. de Québec, 9th year, no. 18, p. 223-238.
- Van Ingen, R., 1967, (Abs.) Granite Studies related to ore formation in the Eastern Townships, Québec; in Proceedings, Symposium on Geochemical Prospecting, Ottawa, April, 1966; Geol. Surv. of Canada, paper 66-54, pp. 277-278.