

University of New Hampshire

University of New Hampshire Scholars' Repository

Molecular, Cellular and Biomedical Sciences
Scholarship

Molecular, Cellular and Biomedical Sciences

3-17-2016

Draft Genome Sequence of *Photorhabdus luminescens* subsp. *laumondii* HP88, an Entomopathogenic Bacterium Isolated from Nematodes

Shimaa Ghazal

University of New Hampshire, Durham, Shimaa.Ghazal@unh.edu

Rediet Oshone

University of New Hampshire, Durham

Stephen Simpson

University of New Hampshire, Durham

Krystalynne Morris

University of New Hampshire, Durham

Feseha Abebe-Akele

University of New Hampshire, Durham

Follow this and additional works at: https://scholars.unh.edu/mcbs_facpub

Comments

See next page for additional authors

This is an article published by American Society for Microbiology in Genome Announcements in 2016, available online: <https://dx.doi.org/10.1128/genomeA.00154-16>

Recommended Citation

Ghazal, S. R. Oshone, S. Simpson, K. Morris, F. Abebe-Akele, W. K. Thomas, K. M. Khalil, and L. S. Tisa. 2016. Draft Genome Sequence of *Photorhabdus luminescens* subsp. *laumondii* HP88, an Entomopathogenic Bacterium Isolated from Nematodes. *Genome Announc.* 4(2):e00154-16. doi:10.1128/genomeA.00154-16.

This Article is brought to you for free and open access by the Molecular, Cellular and Biomedical Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Molecular, Cellular and Biomedical Sciences Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

Authors

Shimaa Ghazal, Rediet Oshone, Stephen Simpson, Krystalynne Morris, Feseha Abebe-Akele, W. Kelley Thomas, Kamal M. Khalil, and Louis S. Tisa

Draft Genome Sequence of *Photorhabdus luminescens* subsp. *laumondii* HP88, an Entomopathogenic Bacterium Isolated from Nematodes

Shimaa Ghazal,^{a,b} Rediet Oshone,^a Stephen Simpson,^a Krystalynne Morris,^a Feseha Abebe-Akele,^a W. Kelley Thomas,^a Kamal M. Khalil,^b Louis S. Tisa^a

University of New Hampshire, Durham, New Hampshire, USA^a; Genetic Engineering & Biotechnology Division, Genetics and Cytology Department, Applied Microbial Genetics Laboratory, National Research Centre, Dokki, Cairo, Egypt^b

***Photorhabdus luminescens* subsp. *laumondii* HP88 is an entomopathogenic bacterium that forms a symbiotic association with *Heterorhabditis* nematodes. We report here a 5.27-Mbp draft genome sequence for *P. luminescens* subsp. *laumondii* HP88, with a G+C content of 42.4% and containing 4,243 candidate protein-coding genes.**

Received 3 February 2016 Accepted 5 February 2016 Published 17 March 2016

Citation Ghazal S, Oshone R, Simpson S, Morris K, Abebe-Akele F, Thomas WK, Khalil KM, Tisa LS. 2016. Draft genome sequence of *Photorhabdus luminescens* subsp. *laumondii* HP88, an entomopathogenic bacterium isolated from nematodes. *Genome Announc* 4(2):e00154-16. doi:10.1128/genomeA.00154-16.

Copyright © 2016 Ghazal et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Louis S. Tisa, louis.tisa@unh.edu.

Photorhabdus species are Gram-negative motile bioluminescent bacteria that maintain two distinct lifestyles as insect pathogens and in a symbiotic relationship with the entomopathogenic *Heterorhabditis* nematodes (see references 1–7 for a review). The life cycles of *Photorhabdus* and its nematode host *Heterorhabditis* are best described as a cyclic association that begins and ends with infective juvenile (IJ) nematodes. A monoculture of *Photorhabdus* is maintained within the anterior region of the IJ nematode's intestine (8, 9). The nematodes actively seek and infect insect hosts by entering through natural openings or by burrowing directly through the insect cuticle. Once inside the insect, the nematodes regurgitate the bacteria into the hemolymph (8). The bacteria release highly virulent toxins (10, 11), which results in insect death in <48 h. As the bacteria enter the stationary phase of their growth cycle, they secrete extracellular enzymes that aid in breaking down insect tissue, thereby providing nutrients for both the bacteria and nematodes. The bacteria also generate essential growth factors for the nematode growth and development. The growth and development of *Heterorhabditis* nematodes have an obligate requirement for their specific bacterial symbiont (12). The bacteria also release antibiotics to prevent secondary invaders and putrefaction of the insect carcass (13, 14). After several days of feeding, the nematodes and bacteria reassociate and leave in search of a new insect host.

Based on molecular analysis, the *Photorhabdus* genus is divided into three bacterial species: *P. luminescens*, *P. temperata*, and *P. asymbiotica* (15, 16). Our understanding of these bacteria has been greatly enhanced by the genome sequencing of strains from all three established species: *P. luminescens* TT01 (17), *P. asymbiotica* ATCC 43949 (18, 19), *P. temperata* NC19 (20), *P. temperata* Meg1 (21), *P. luminescens* BA1 (22), *P. asymbiotica* Kingcliff (23), and *P. temperata* M121 (24). Here, we

present a draft genome sequence for *P. luminescens* subsp. *laumondii* HP88, which was isolated from *Heterorhabditis bacteriophora* nematodes found in Utah (25).

The draft genome of *P. luminescens* strain HP88 was generated at the Hubbard Genome Center (University of New Hampshire, Durham, NH) using Illumina technology (26) techniques. A standard Illumina shotgun library was constructed and sequenced using the Illumina HiSeq 2000 platform, which generated 7,680,248 reads (260-bp insert size) totaling 1,120.0 Mbp. The Illumina sequence data were assembled using CLC Genomics Workbench (version 8.5) and AllPaths-LG (version r41043) (27). The final draft assembly contained 287 contigs, with an N_{50} of 34.4 kb. The total size of the genome is 5.27 Mbp, and the final assembly is based on 949 Mb of Illumina draft data, providing an average 163× coverage of the genome.

The high-quality draft genome of *P. luminescens* strain HP99 was resolved to 287 contigs consisting of 5,268,230 bp, with a G+C content of 42.4%. The assembled *P. luminescens* strain HP88 genome was annotated via the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) and resulted in 4,243 candidate protein-coding genes.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [LJPB00000000](https://www.ncbi.nlm.nih.gov/nuclink/LJPB00000000). The version described in this paper is version LJPB01000000.

ACKNOWLEDGMENTS

This work was supported in part by USDA NIFA grant 2009-35302-05257 to L.S.T. and by the College of Life Science and Agriculture at the University of New Hampshire–Durham. S.G. was supported by the Egyptian Cultural and Educational Bureau, Washington, DC. Sequencing was performed on an Illumina HiSeq 2500 purchased with NSF MRI grant DBI-1229361 to W.K.T.

FUNDING INFORMATION

This work, including the efforts of Shima Ghazal, was funded by Egyptian Cultural and Education Bureau. This work, including the efforts of W. Kelley Thomas, was funded by National Science Foundation (NSF) (DBI-1229361). This work, including the efforts of Louis S. Tisa, was funded by U.S. Department of Agriculture (USDA) (NIFA 2009-35302-05257).

REFERENCES

- Boemare N, Givaudan A, Brehelin M, Laumond C. 1997. Symbiosis and pathogenicity of nematode-bacterium complexes. *Symbiosis* 22:21–45.
- Forst S, Dowds B, Boemare N, Stackebrandt E. 1997. *Xenorhabdus* and *Photorhabdus* spp.: bugs that kill bugs. *Annu Rev Microbiol* 51:47–72. <http://dx.doi.org/10.1146/annurev.micro.51.1.47>.
- Forst S, Neelson K. 1996. Molecular biology of the symbiotic pathogenic bacteria *Xenorhabdus* spp. and *Photorhabdus* spp. *Microbiol Res* 60: 21–43.
- Goodrich-Blair H, Clarke DJ. 2007. Mutualism and pathogenesis in *Xenorhabdus* and *Photorhabdus*: two roads to the same destination. *Mol Microbiol* 64:260–268. <http://dx.doi.org/10.1111/j.1365-2958.2007.05671.x>.
- Clarke DJ. 2008. *Photorhabdus*: a model for the analysis of pathogenicity and mutualism. *Cell Microbiol* 10:2159–2167. <http://dx.doi.org/10.1111/j.1462-5822.2008.01209.x>.
- Waterfield NR, Ciche T, Clarke D. 2009. *Photorhabdus* and a host of hosts. *Annu Rev Microbiol* 63:557–574. <http://dx.doi.org/10.1146/annurev.micro.091208.073507>.
- Burnell AM, Stock SP. 2000. *Heterorhabditis*, *Steinernema* and their bacterial symbionts—lethal pathogens of insects. *Nematology* 2:31–42. <http://dx.doi.org/10.1163/156854100508872>.
- Ciche TA, Ensign JC. 2003. For the insect pathogen *Photorhabdus luminescens*, which end of a nematode is out? *Appl Environ Microbiol* 69: 1890–1897. <http://dx.doi.org/10.1128/AEM.69.4.1890-1897.2003>.
- Endo BY, Nickle WR. 1991. Ultrastructure of the intestinal epithelium, lumen, and associated bacteria in *Heterorhabditis-bacteriophora*. *J Helminthol Soc Wash* 58:202–212.
- Bowen D, Rocheleau TA, Blackburn M, Andreev O, Golubeva E, Bhartiya R, Ffrench-Constant RH. 1998. Insecticidal toxins from the bacterium *Photorhabdus luminescens*. *Science* 280:2129–2132. <http://dx.doi.org/10.1126/science.280.5372.2129>.
- Ffrench-Constant RH, Bowen DJ. 2000. Novel insecticidal toxins from nematode-symbiotic bacteria. *Cell Mol Life Sci* 57:828–833. <http://dx.doi.org/10.1007/s000180050044>.
- Thomas GM, Poinar GO. 1979. *Xenorhabdus* gen. nov., a genus of entomopathogenic, nematophilic bacteria of the family *Enterobacteriaceae*. *Int J Syst Bacteriol* 29:352–360. <http://dx.doi.org/10.1099/00207713-29-4-352>.
- Akhurst RJ. 1982. Antibiotic activity of *Xenorhabdus* spp., bacteria symbiotically associated with insect pathogenic nematodes of the families *Heterorhabditidae* and *Steinernematidae*. *J Gen Microbiol* 128:3061–3065. <http://dx.doi.org/10.1099/00221287-128-12-3061>.
- Richardson WH, Schmidt TM, Neelson KH. 1988. Identification of an anthraquinone pigment and a hydroxystilbene antibiotic from *Xenorhabdus luminescens*. *Appl Environ Microbiol* 54:1602–1605.
- Fischer-Le Saux M, Viallard V, Brunel B, Normand P, Boemare NE. 1999. Polyphasic classification of the genus *Photorhabdus* and proposal of new taxa: *P. luminescens* subsp. *luminescens* subsp. nov., *P. luminescens* subsp. *akhurstii* subsp. nov., *P. luminescens* subsp. *laumondii* subsp. nov., *P. temperata* sp. nov., *P. temperata* subsp. *temperata* subsp. nov. and *P. asymbiotica* sp. nov. *Int J Syst Bacteriol* 49:1645–1656. <http://dx.doi.org/10.1099/00207713-49-4-1645>.
- Boemare N. 2002. Interactions between the partners of the entomopathogenic bacterium nematode complexes, *Steinernema*-*Xenorhabdus* and *Heterorhabditis*-*Photorhabdus*. *Nematology* 4:601–603. <http://dx.doi.org/10.1163/15685410260438863>.
- Duchaud E, Rusniok C, Frangeul L, Buchrieser C, Givaudan A, Taourit S, Bocs S, Boursaux-Eude C, Chandler M, Charles JF, Dassa E, Derose R, Derzelle S, Freyssinet G, Gaudriault S, Médigue C, Lanois A, Powell K, Siguier P, Vincent R, Wingate V, Zouine M, Glaser P, Boemare N, Danchin A, Kunst F. 2003. The genome sequence of the entomopathogenic bacterium *Photorhabdus luminescens*. *Nat Biotechnol* 21:1307–1313. <http://dx.doi.org/10.1038/nbt886>.
- Wilkinson P, Waterfield NR, Crossman L, Corton C, Sanchez-Contreras M, Vlisidou I, Barron A, Bignell A, Clark L, Ormond D, Mayho S, Bason N, Smith F, Simmonds M, Churcher C, Harris D, Thompson NR, Quail M, Parkhill J, Ffrench-Constant RH. 2009. Comparative genomics of the emerging human pathogen *Photorhabdus asymbiotica* with the insect pathogen *Photorhabdus luminescens*. *BMC Genomics* 10:302. <http://dx.doi.org/10.1186/1471-2164-10-302>.
- ffrench-Constant R, Waterfield N, Daborn P, Joyce S, Bennett H, Au C, Dowling A, Boundy S, Reynolds S, Clarke D. 2003. *Photorhabdus*: towards a functional genomic analysis of a symbiont and pathogen. *FEMS Microbiol Rev* 26:433–456. <http://dx.doi.org/10.1111/j.1574-6976.2003.tb00625.x>.
- Hurst S, Rowedder H, Michaels B, Bullock H, Jackobeck R, Abebe-Akele F, Durkavovic U, Gately J, Janicki E, Tisa LS. 2015. Elucidation of the *Photorhabdus temperata* genome and generation of a transposon mutant library to identify motility mutants altered in pathogenesis. *J Bacteriol* 197:2201–2216. <http://dx.doi.org/10.1128/JB.00197-15>.
- Hurst SG, IV, Ghazal S, Morris K, Abebe-Akele F, Thomas WK, Badr UM, Hussein MA, AbouZaied MA, Khalil KM, Tisa LS. 2014. Draft genome sequence of *Photorhabdus temperata* strain meg1, an entomopathogenic bacterium isolated from *Heterorhabditis megidis* nematodes. *Genome Announc* 2(6):e01273-14. <http://dx.doi.org/10.1128/genomeA.01273-14>.
- Ghazal S, Hurst SG, IV, Morris K, Abebe-Akele F, Thomas WK, Badr UM, Hussein MA, AbouZaied MA, Khalil KM, Tisa LS. 2014. Draft genome sequence of *Photorhabdus luminescens* strain BA1, an entomopathogenic bacterium isolated from nematodes found in Egypt. *Genome Announc* 2(2):e00396-14. <http://dx.doi.org/10.1128/genomeA.00396-14>.
- Wilkinson P, Paszkiewicz K, Moorhouse A, Szubert JM, Beatson S, Gerrard J, Waterfield NR, Ffrench-Constant RH. 2010. New plasmids and putative virulence factors from the draft genome of an Australian clinical isolate of *Photorhabdus asymbiotica*. *FEMS Microbiol Lett* 309: 136–143. <http://dx.doi.org/10.1111/j.1574-6968.2010.02030.x>.
- Park GS, Khan AR, Hong SJ, Jang EK, Ullah I, Jung BK, Choi J, Yoo NK, Park KJ, Shin JH. 2013. Draft genome sequence of entomopathogenic bacterium *Photorhabdus temperata* strain M1021, isolated from nematodes. *Genome Announc* 1(5):e00747-13. <http://dx.doi.org/10.1128/genomeA.00747-13>.
- Poinar GO, Georgis R. 1990. Characterization and field application of *Heterorhabditis bacteriophora* strain HP88 (*Heterorhabditidae*: *Rhabditida*). *Rev Nematol* 13:387–393.
- Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* 5:433–438. <http://dx.doi.org/10.1517/14622416.5.4.433>.
- Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, Berlin AM, Aird D, Costello M, Daza R, Williams L, Nicol R, Gnirke A, Nusbaum C, Lander ES, Jaffe DB. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci USA* 108:1513–1518. <http://dx.doi.org/10.1073/pnas.1017351108>.