

9-14-2017

Permanent Draft Genome Sequence of Photorhabdus temperata Strain Hm, an Entomopathogenic Bacterium Isolated from Nematodes

Shimaa M. Ghazal

University of New Hampshire, Durham, Shimaa.Ghazal@unh.edu

Erik Swanson

University of New Hampshire, Durham

Stephen Simpson

University of New Hampshire, Durham

Krystalynne Morris

University of New Hampshire, Durham

Feseha Abebe-Akele

University of New Hampshire, Durham

See next page for additional authors

Follow this and additional works at: https://scholars.unh.edu/mcbs_facpub

Recommended Citation

GHAZAL, S., E. SWANSON, S. Simpson, K. Morris, F. Abebe-Akele, W. K. Thomas, K. M. Khalil, and L. S. Tisa. 2017. Draft Genome Sequence of Photorhabdus temperta strain Hm, an Entomopathogenic Bacterium Isolated from Nematodes. GenomeA 5:e00974-17 doi: 10.1128/genomeA.00974-17

This Article is brought to you for free and open access by the Molecular, Cellular and Biomedical Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Molecular, Cellular and Biomedical Sciences Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact nicole.hentz@unh.edu.

Authors

Shimaa M. Ghazal, Erik Swanson, Stephen Simpson, Krystalynne Morris, Feseha Abebe-Akele, W. Kelley Thomas, Kamal M. Khalil, and Louis S. Tisa



Permanent Draft Genome Sequence of *Photorhabdus temperata* Strain Hm, an Entomopathogenic Bacterium Isolated from Nematodes

Shimaa Ghazal,^{a,b} Erik Swanson,^a Stephen Simpson,^a Krystalynne Morris,^a Feseha Abebe-Akele,^a W. Kelley Thomas,^a Kamal M. Khalil,^b Louis S. Tisa^a

University of New Hampshire, Durham, New Hampshire, USA^a; Applied Microbial Genetics Laboratory, Genetics and Cytology Department, Genetic Engineering & Biotechnology Division, National Research Center, Dokki, Cairo, Egypt^b

ABSTRACT *Photorhabdus temperata* strain Hm is an entomopathogenic bacterium that forms a symbiotic association with *Heterorhabditis* nematodes. Here, we report a 5.0-Mbp draft genome sequence for *P. temperata* strain Hm with a G+C content of 44.1% and containing 4,226 candidate protein-encoding genes.

Members of the genus *Photorhabdus* maintain two distinct lifestyles as insect pathogens and in a symbiotic relationship with the entomopathogenic *Heterorhabditid* nematodes (for reviews, see references 1–3). The *Heterorhabditis* nematodes carry a monoculture of *Photorhabdus* within the anterior region of the infective juvenile (IJ) nematode's intestine (4, 5) and actively seek insect prey in the soil. The nematodes infect a wide range of insect hosts by entering through natural openings or by burrowing directly through the insect cuticle. Once inside the insect, the nematodes regurgitate the bacteria into the hemolymph (4). The bacteria kill the insect within 48 h by releasing highly virulent toxins (6–9). As the bacteria enter the stationary phase of their growth cycle, they secrete extracellular enzymes that aid in breaking down insect tissue, thereby providing nutrients for both the bacteria and the nematodes. The bacteria also generate essential growth factors for nematode growth and development. The growth and development of *Heterorhabditis* nematodes have an obligate requirement for their specific bacterial symbiont (10). The bacteria also release antibiotics to prevent secondary invaders and putrefaction of the insect carcass (11, 12). After several days of feeding, the nematodes and bacteria reassociate and leave in search of a new insect host.

Members of *Photorhabdus* are classified taxonomically into one of three species, *P. luminescens*, *P. temperata*, or *P. asymbiotica* (13–15). Several subspecies are recognized. Our understanding of these bacteria has been greatly enhanced by genome sequencing of the three established species, including that of *P. luminescens* TT01 (16), *P. asymbiotica* ATCC 43949 (17), *P. temperata* subsp. *khanii* NC19 (18), *P. temperata* Meg1 (19), *P. luminescens* BA1 (20), *P. luminescens* subsp. *laumondii* HP88 (21), *P. asymbiotica* Kingcliff (22), *P. temperata* subsp. *temperata* M121 (23), *P. luminescens* subsp. PB45.5 (24), and *P. asymbiotica* subsp. *australis* PB68.1 (24). Here, we present a draft genome sequence for *P. temperata* strain Hm, which was isolated from *Heterorhabditis* nematodes found in Georgia (10).

The draft genome sequence of *P. temperata* strain Hm was generated at the Hubbard Genome Center (University of New Hampshire, Durham, NH) using Illumina technology (25) techniques. A standard Illumina shotgun library was constructed and sequenced using the Illumina HiSeq2500 platform, which generated 15,461,198 reads (150-bp insert size) totaling 2,303.7 Mbp. The Illumina sequence data were trimmed by

Received 2 August 2017 Accepted 8 August 2017 Published 14 September 2017

Citation Ghazal S, Swanson E, Simpson S, Morris K, Abebe-Akele F, Thomas WK, Khalil KM, Tisa LS. 2017. Permanent draft genome sequence of *Photorhabdus temperata* strain Hm, an entomopathogenic bacterium isolated from nematodes. *Genome Announc* 5:e00974-17. <https://doi.org/10.1128/genomeA.00974-17>.

Copyright © 2017 Ghazal et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Louis S. Tisa, louis.tisa@unh.edu.

Trimmomatic version 0.32 (26) and assembled using SPAdes version 3.5 (26) and ALLPaths-LG version r52488 (27). The final draft assembly contained 151 contigs with an N_{50} contig size of 71.2 kb and $356.8\times$ coverage of the genome. The final assembled genome contained a total sequence length of 5,003,482 bp with a G+C content of 44.1%. The assembled *P. temperata* strain Hm genome was annotated via the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) and resulted in 4,226 candidate protein-encoding genes and 68 tRNA and 4 rRNA regions.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number [MBJU00000000](https://www.ncbi.nlm.nih.gov/nuclink/MBJU00000000). The version described in this paper is MBJU01000000.

ACKNOWLEDGMENTS

This work was supported in part by USDA NIFA grant 2009-35302-05257 and by the College of Life Science and Agriculture at the University of New Hampshire-Durham. S.G. was supported by the Egyptian Cultural and Educational Bureau, Washington, DC, USA.

Sequencing was performed on an Illumina HiSeq2500 instrument purchased with NSF MRI grant DBI-1229361 to W.K.T.

REFERENCES

- Goodrich-Blair H, Clarke DJ. 2007. Mutualism and pathogenesis in *Xenorhabdus* and *Photorhabdus*: two roads to the same destination. *Mol Microbiol* 64:260–268. <https://doi.org/10.1111/j.1365-2958.2007.05671.x>.
- Clarke DJ. 2008. *Photorhabdus*: a model for the analysis of pathogenicity and mutualism. *Cell Microbiol* 10:2159–2167. <https://doi.org/10.1111/j.1462-5822.2008.01209.x>.
- Waterfield NR, Ciche T, Clarke D. 2009. *Photorhabdus* and a host of hosts. *Annu Rev Microbiol* 63:557–574. <https://doi.org/10.1146/annurev.micro.091208.073507>.
- Ciche TA, Ensign JC. 2003. For the insect pathogen *Photorhabdus luminescens*, which end of a nematode is out? *Appl Environ Microbiol* 69:1890–1897. <https://doi.org/10.1128/AEM.69.4.1890-1897.2003>.
- Endo BY, Nickle WR. 1991. Ultrastructure of the intestinal epithelium, lumen, and associated bacteria in *Heterorhabditis*-Bacteriophora. *J Helminthol Soc Wash* 58:202–212.
- Bowen D, Rocheleau TA, Blackburn M, Andreev O, Golubeva E, Bhartiya R, Ffrench-Constant RH. 1998. Insecticidal toxins from the bacterium *Photorhabdus luminescens*. *Science* 280:2129–2132. <https://doi.org/10.1126/science.280.5372.2129>.
- Bowen DJ, Ensign JC. 1998. Purification and characterization of a high-molecular-weight insecticidal protein complex produced by the entomopathogenic bacterium *Photorhabdus luminescens*. *Appl Environ Microbiol* 64:3029–3035.
- Ffrench-Constant R, Bowen D. 1999. *Photorhabdus* toxins: novel biological insecticides. *Curr Opin Microbiol* 2:284–288. [https://doi.org/10.1016/S1369-5274\(99\)80049-6](https://doi.org/10.1016/S1369-5274(99)80049-6).
- Ffrench-Constant RH, Bowen DJ. 2000. Novel insecticidal toxins from nematode-symbiotic bacteria. *Cell Mol Life Sci* 57:828–833. <https://doi.org/10.1007/s000180050044>.
- Thomas GM, Poinar GO. 1979. *Xenorhabdus* gen. nov., a genus of entomopathogenic, nematophilic bacteria of the family *Enterobacteriaceae*. *Int J Syst Bacteriol* 29:352–360. <https://doi.org/10.1099/00207713-29-4-352>.
- Akhurst RJ. 1982. Antibiotic activity of *Xenorhabdus* spp., bacteria symbiotically associated with insect pathogenic nematodes of the families *Heterorhabditidae* and *Steinernematidae*. *J Gen Microbiol* 128:3061–3065. <https://doi.org/10.1099/00221287-128-12-3061>.
- Richardson WH, Schmidt TM, Nealson KH. 1988. Identification of an anthraquinone pigment and a hydroxystilbene antibiotic from *Xenorhabdus luminescens*. *Appl Environ Microbiol* 54:1602–1605.
- Fischer-Le Saux M, Viillard V, Brunel B, Normand P, Boemare NE. 1999. Polyphasic classification of the genus *Photorhabdus* and proposal of new taxa: *P. luminescens* subsp. *luminescens* subsp. nov., *P. luminescens* subsp. *akhurstii* subsp. nov., *P. luminescens* subsp. *laumondii* subsp. nov., *P. temperata* sp. nov., *P. temperata* subsp. *temperata* subsp. nov. and *P. asymbiotica* sp. nov. *Int J Syst Bacteriol* 49:1645–1656. <https://doi.org/10.1099/00207713-49-4-1645>.
- Tailliez P, Pagès S, Ginibre N, Boemare N. 2006. New insight into diversity in the genus *Xenorhabdus*, including the description of ten novel species. *Int J Syst Evol Microbiol* 56:2805–2818. <https://doi.org/10.1099/ijs.0.64287-0>.
- Tounsi S, Blight M, Jaoua S, Pimenta AD. 2006. From insects to human hosts: identification of major genomic differences between entomopathogenic strains of *Photorhabdus* and the emerging human pathogen *Photorhabdus asymbiotica*. *Int J Med Microbiol* 296:521–530. <https://doi.org/10.1016/j.ijmm.2006.06.004>.
- Duchaud E, Rusniok C, Frangeul L, Buchrieser C, Givaudan A, Taourit S, Bocs S, Boursaux-Eude C, Chandler M, Charles JF, Dassa E, Deroose R, Derzelle S, Freyssinet G, Gaudriault S, Médigue C, Lanois A, Powell K, Siguier P, Vincent R, Wingate V, Zouine M, Glaser P, Boemare N, Danchin A, Kunst F. 2003. The genome sequence of the entomopathogenic bacterium *Photorhabdus luminescens*. *Nat Biotechnol* 21:1307–1313. <https://doi.org/10.1038/nbt886>.
- Wilkinson P, Waterfield NR, Crossman L, Corton C, Sanchez-Contreras M, Vlisidou I, Barron A, Bignell A, Clark L, Ormond D, Mayho M, Bason N, Smith F, Simmonds M, Churcher C, Harris D, Thompson NR, Quail M, Parkhill J, Ffrench-Constant RH. 2009. Comparative genomics of the emerging human pathogen *Photorhabdus asymbiotica* with the insect pathogen *Photorhabdus luminescens*. *BMC Genomics* 10:302. <https://doi.org/10.1186/1471-2164-10-302>.
- Hurst S, Rowedder H, Michaels B, Bullock H, Jackobeck R, Abebe-Akele F, Durakovic U, Gately J, Janicki E, Tisa LS. 2015. Elucidation of the *Photorhabdus temperata* genome and generation of a transposon mutant library to identify motility mutants altered in pathogenesis. *J Bacteriol* 197:2201–2216. <https://doi.org/10.1128/JB.00197-15>.
- Hurst SG, Ghazal S, Morris K, Abebe-Akele F, Thomas WK, Badr UM, Hussein MA, Abouzaied MA, Khalil KM, Tisa LS. 2014. Draft genome sequence of *Photorhabdus temperata* strain Meg1, an entomopathogenic bacterium isolated from *Heterorhabditis megidis* nematodes. *Genome Announc* 2(6):e01273-14. <https://doi.org/10.1128/genomeA.01273-14>.
- Ghazal S, Hurst SG, Morris K, Abebe-Akele F, Thomas WK, Badr UM, Hussein MA, Abouzaied MA, Khalil KM, Tisa LS. 2014. Draft genome sequence of *Photorhabdus luminescens* strain BA1, an entomopathogenic bacterium isolated from nematodes found in Egypt. *Genome Announc* 2(2):e00396-14. <https://doi.org/10.01128/genomeA.00396-14>.
- Ghazal S, Oshone R, Simpson S, Morris K, Abebe-Akele F, Thomas WK, Khalil KM, Tisa LS. 2016. Draft genome sequence of *Photorhabdus luminescens* subsp. *laumondii* HP88, an entomopathogenic bacterium isolated from nematodes. *Genome Announc* 4(2):e00154-16. <https://doi.org/10.1128/genomeA.00154-16>.
- Wilkinson P, Paszkiewicz K, Moorhouse A, Szubert JM, Beatson S, Gerrard J, Waterfield NR, Ffrench-Constant RH. 2010. New plasmids and putative virulence factors from the draft genome of an Australian clinical isolate

- of *Photorhabdus asymbiotica*. FEMS Microbiol Lett 309:136–143. <https://doi.org/10.1111/j.1574-6968.2010.02030.x>.
23. Park GS, Khan AR, Hong SJ, Jang EK, Ullah I, Jung BK, Choi J, Yoo NK, Park KJ, Shin JH. 2013. Draft genome sequence of entomopathogenic bacterium *Photorhabdus temperata* strain M1021, isolated from nematodes. Genome Announc 1(5):e00154-16. <https://doi.org/10.1128/genomeA.00747-13>.
 24. Tobias NJ, Mishra B, Gupta DK, Sharma R, Thines M, Stinear TP, Bode HB. 2016. Genome comparisons provide insights into the role of secondary metabolites in the pathogenic phase of the *Photorhabdus* life cycle. BMC Genomics 17:537. <https://doi.org/10.1186/s12864-016-2862-4>.
 25. Bennett S. 2004. Solexa Ltd. Pharmacogenomics 5:433–438. <https://doi.org/10.1517/14622416.5.4.433>.
 26. Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
 27. Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, Berlin AM, Aird D, Costello M, Daza R, Williams L, Nicol R, Gnirke A, Nusbaum C, Lander ES, Jaffe DB. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. Proc Natl Acad Sci U S A 108:1513–1518. <https://doi.org/10.1073/pnas.1017351108>.