

University of New Hampshire

University of New Hampshire Scholars' Repository

Molecular, Cellular and Biomedical Sciences
Scholarship

Molecular, Cellular and Biomedical Sciences

1-1-2017

Draft Genome Sequence of the Symbiotic Frankia Sp. Strain KB5 Isolated from Root Nodules of Casuarina equisetifolia

Celine Pesce

University of New Hampshire, Durham, Celine.Pesce@unh.edu

Erik Swanson

University of New Hampshire, Durham

Stephen Simpson

University of New Hampshire, Durham

Krystalynne Morris

University of New Hampshire, Durham

W. Kelley Thomas

University of New Hampshire, Durham, Kelley.Thomas@unh.edu

See next page for additional authors

Follow this and additional works at: https://scholars.unh.edu/mcbs_facpub

Comments

This is an article published by Ivyspring Interantional Publisher in Journal of Genomics in 2017, available online:

<https://dx.doi.org/10.7150/jgen.20887>

Recommended Citation

Pesce, C. E. Swanson, S. Simpson, K. Morris, W.K. Thomas, L.S. Tisa, and A. Sellstedt. 2017. Draft Genome Sequence for Frankia sp. Strain KB5 Isolated from the Root Nodules of Casuarina equisetifolia. J. Genomics 5: 64-67. doi: 10.7150/jgen.20887

This Article is brought to you for free and open access by the Molecular, Cellular and Biomedical Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Molecular, Cellular and Biomedical Sciences Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

Authors

Celine Pesce, Erik Swanson, Stephen Simpson, Krystalynne Morris, W. Kelley Thomas, Louis S. Tisa, and Anita Sellstedt

Short Research Paper

Draft Genome Sequence of the Symbiotic *Frankia* Sp. Strain KB5 Isolated from Root Nodules of *Casuarina equisetifolia*

Céline Pesce¹, Erik Swanson¹, Stephen Simpson¹, Krystalynne Morris¹, W. Kelley Thomas¹, Louis S. Tisa¹✉ and Anita Sellstedt²✉

1. University of New Hampshire, Durham, New Hampshire, USA;
2. UPSC, Department of Plant physiology, Umeå University, S-90187 Umeå, Sweden.

✉ Corresponding authors: Louis S. Tisa **Mailing address:** Department of Molecular, Cellular, and Biomedical Sciences, University of New Hampshire, 46 College Rd., Durham, NH 03824-2617. **Email:** louis.tisa@unh.edu **Telephone:** 1-603-862-2442 **Fax:** 1-603-862-2621 Anita Sellstedt **Mailing address:** UPSC, Department of Plant physiology, Umeå University, S-90187 Umeå, Sweden. **Email:** anita.sellstedt@umu.se

© Ivyspring International Publisher. This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY-NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>). See <http://ivyspring.com/terms> for full terms and conditions.

Received: 2017.05.05; Accepted: 2017.05.29; Published: 2017.06.09

Abstract

Frankia sp. strain KB5 was isolated from *Casuarina equisetifolia* and previous studies have shown both nitrogenase and uptake hydrogenase activities under free-living conditions. Here, we report 5.5-Mbp draft genome sequence with a G+C content of 70.03 %, 4,958 candidate protein-encoding genes, and 2 rRNA operons.

Key words: actinobacteria, actinorhizal symbiosis, hydrogenase, nitrogen fixation, natural products, host microbe interactions, genomes.

Introduction

Biological nitrogen fixation, which is carried out by prokaryotes, converts atmospheric dinitrogen gas into a reduced biologically useful form. One group of these nitrogen-fixing microbes is the genus *Frankia*, which is found in association with actinorhizal plants or free-living in the soil [1, 2]. Based on molecular studies, there are four distinct *Frankia* lineages [3-5]. Three of the Clusters (I, II, and III) are infective on actinorhizal host plants, while members of Cluster IV are considered “atypical” strains, which are unable to re-infect the plants or form ineffective nodules that unable to fix nitrogen.

Frankia sp. strain KB5 was isolated from the root nodules of *Casuarina equisetifolia* ssp. *incana* in Australia [6]. This strain is a member of *Frankia* subcluster Ic, which has its host range limited to *Casuarina* and *Allocasuarina* plants. The physiology of *Frankia* sp. strain KB5 has been well studied especially the close relationship between hydrogenase and

nitrogenase activities [7-9]. Interestingly, the addition of nickel results in increased uptake hydrogenase activity [7]. *Frankia* sp. strain KB5 genome was chosen to be sequenced for several reasons including an interesting physiology. Furthermore, this strain represents an isolate from Australia, the native biogeographical region of *Frankia* subcluster Ic [10]. This database could provide more information on the evolution of this *Frankia* subcluster.

Sequencing of the draft genome of *Frankia* sp. strain KB5 was performed at the Hubbard Center for Genome Studies (University of New Hampshire, Durham, NH) using Illumina technology techniques [11]. A standard Illumina shotgun library was constructed and sequenced using the Illumina HiSeq2500 platform, using a pair-end library with an average size of 600 bp obtaining 10,539,470 reads of 250 bp in length. The Illumina sequence data were trimmed by Trimmomatic version 0.32 [12], assembled

using Spades version 3.5 [12] and ALLPaths-LG version r52488 [13]. The final draft assembly for *Frankia* sp. strain KB5 consisted of 420 contigs with an N₅₀ contig size of 24.2 kb and 236.8X coverage of the genome. The final assembled genome contained a total sequence length of 5,455,564 bp with a G+C content of 70.03%.

The assembled *Frankia* sp. strain KB5 genome was annotated via the NCBI Prokaryotic Genome Annotation Pipeline (PGAP), and resulted in 4,958 candidate protein-encoding genes and 2 rRNA operons. The genome features of *Frankia* sp. strain KB5 fall with the realm of the other *Casuarina* genomes including *Frankia casuarinae* strain (CcI3) [14] the type strain (Table 1).

The genome also contained a *nif* and 2 *hup* operons encoding the nitrogenase and uptake hydrogenase enzymes, respectively. The operons were organized similar to those reported for *Frankia* cluster I genomes [15]. The 2 *hup* operons have been shown to be expressed differently, with *hup* 1 being mainly expressed in free-living conditions and *hup* 2 in symbiotic [16]. Bioinformatic analysis of this genomes by the use of the AntiSMASH program [17, 18] revealed the presence of high numbers of secondary metabolic biosynthetic gene clusters, which is consistent with previous results with other *Frankia* genomes including subcluster Ic [15, 19]. Table 2

shows a comparison of the various profiles of different *Casuarina* isolates for these secondary metabolic biosynthetic gene clusters. Although the majority of these secondary metabolic biosynthetic gene clusters were shared among the *Casuarina* genomes, the *Frankia* sp. strain KB5 genome contained several unique clusters that had homologues in other bacteria or were completely novel. For example, cluster KB-24 (location: *KB15_16785-KB15_16785* genes) involved in terpene biosynthesis had no homology with any of the *Frankia* genomes, but was homologous to the phenalinolactoneA biosynthetic gene cluster of *Streptomyces pactum* KLBMP 5084, including the surrounding gene neighborhood. Cluster KB-26 (location: *KB15_17660-KB15_17705* genes) containing a nonribosomal peptide synthase (NRPS) is only found in the *Frankia* sp. strain KB5 genome and predicted to produce a core structure (Fig. 1). Further bioinformatics analysis revealed that the *Frankia* sp. strain KB5 genome contained 715 unique genes that had no homologues in any of the other *Casuarina* genomes. Although some of these genes like cluster KB-24 genes have predicted functions, the majority of the genes code for hypothetical proteins without a known function. The hypothesis that these genes play a role in the biogeographical distribution of this strain remains to be tested.

Table 1. Genome features of *Frankia* sp. strain KB5 and other *Frankia* strains isolated from *Casuarina* root nodules.

Strain	Source	Location ¹	Size (Mb)	No. of Contig(s)	G+C (%)	No. of CDS	No. of rRNA	No. of tRNA
KB5	This study	Australia	5.46	420	70.0	4,958	6	45
CcI3	[10]	USA	5.43	1	70.1	4,598	6	46
CeD	[20]	Senegal	5.00	120	70.1	4,403	7	45
Allo2	[21]	Uruguay	5.33	110	69.8	4,838	7	46
Thr	[22]	Egypt	5.31	171	70.0	4,805	5	46
BMG5.23	[23]	Tunisia	5.27	167	70.0	4,747	9	47
CcI6	[24]	Egypt	5.39	138	67.6	4,902	9	46
BR	[25]	Brazil	5.23	180	70.0	4,777	5	46

¹The source of the isolate.

Table 2. Biosynthetic gene clusters for natural products found in the genomes from *Casuarina Frankia* strains.

Strain	No. of Biosynthetic gene clusters ¹	NRPS ²	PKS ³	Terpene	Siderophore	Bacteriocin	Lantipeptide
KB5	34	4	9	6	1	1	4
CcI3	29	3	5	4	1	3	6
CeD	30	7	7	4	1	1	4
Allo2	32	7	9	4	1	3	5
Thr	33	6	7	4	1	1	6
BMG5.23	31	8	6	4	1	2	4
CcI6	33	8	8	4	1	3	5
BR	29	5	5	4	1	2	5

¹Biosynthetic gene clusters were identified by the use of the AntiSMASH software [17, 18]

²NRPS: Nonribosomal peptide synthase

³PKS: polyketide synthase including Type I, II, III, Trans-AT, and other types




Figure 1. The predicted *Frankia* chemical structure for secondary metabolic biosynthetic gene cluster KB-24.

In summary, the *Frankia* sp. strain KB5 genome has revealed an interesting potential natural product profile and serves as a representative of *Frankia* subcluster Ic from its native environment. Further analysis of this genome and experimental evidence will be needed to support the predicted natural product profile and to provide insight on the evolution of this *Frankia* subcluster.

Nucleotide sequence accession numbers

This whole-genome shotgun sequence has been deposited at DDBJ/EMBL/GenBank under the accession number MRUJ00000000. The version described in this paper is the first version, MRUJ01000000.

Acknowledgement

Partial funding was provided by the New Hampshire Agricultural Experiment Station. This is Scientific Contribution Number 2725. This work was also supported by the Swedish Energy Agency 38239-1 (AS), USDA National Institute of Food and Agriculture Hatch 022821 (LST), Agriculture and Food Research Initiative Grant 2015-67014-22849 from the USDA National Institute of Food and Agriculture (LST), and the College of Life Science and Agriculture at the University of New Hampshire-Durham. Sequencing was performed on an Illumina HiSeq2500 purchased with an NSF MRI Grant: DBI-1229361 to WK Thomas.

Competing Interests

The authors have declared that no competing interest exists.

References

- Normand P, Benson DR, Berry AM, Tisa LS. Family Frankiaceae. In: Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F, editors. *The Prokaryote – Actinobacteria* Berlin Heidelberg: Springer-Verlag; 2014:339-56
- Chaia EE, Wall LG, Huss-Danell K. Life in soil by the actinorhizal root nodule endophyte *Frankia*. A review. *Symbiosis*. 2010; 51: 201-26.
- Normand P, Orso S, Cournoyer B, Jeannin P, Chapelon C, Dawson J, et al. Molecular phylogeny of the genus *Frankia* and related genera and emendation of the family Frankiaceae. *Int J Syst Bacteriol*. 1996; 46: 1-9.
- Nouioui I, Ghodhbane-Gtari F, Beauchemin NJ, Tisa LS, Gtari M. Phylogeny of members of the *Frankia* genus based on *gyrB*, *nifH* and *glnII* sequences. *Anton Leeuw Int J G*. 2011; 100: 579-87.
- Ghodhbane-Gtari F, Nouioui I, Chair M, Boudabous A, Gtari M. 16S-23S rRNA Intergenic Spacer Region Variability in the Genus *Frankia*. *Microb Ecol*. 2010; 60: 487-95.
- Rosbrook PA, Burggraaf AJP, Reddell P. A Comparison of 2 Methods and Different Media for Isolating *Frankia* from *Casuarina* Root-Nodules. *Plant Soil*. 1989; 120: 187-93.
- Mattsson U, Sellstedt A. Nickel affects activity more than expression of hydrogenase protein in *Frankia*. *Curr Microbiol*. 2002; 44: 88-93.
- Mattsson U, Johansson L, Sandstrom G, Sellstedt A. *Frankia* KB5 possesses a hydrogenase immunologically related to membrane-bound. *Curr Microbiol*. 2001; 42: 438-41.
- Mattsson U, Sellstedt A. Hydrogenase in *Frankia* KB5: Expression of and relation to nitrogenase. *Can J Microbiol*. 2000; 46: 1091-5.
- Normand P, Lapierre P, Tisa LS, Gogarten JP, Alloisio N, Bagnarol E, et al. Genome characteristics of facultatively symbiotic *Frankia* sp strains reflect host range and host plant biogeography. *Genome research*. 2007; 17: 7-15.
- Bennett S. Solexa Ltd. *Pharmacogenomics*. 2004; 5: 433-8.
- Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. 2014; 30: 2114-20.
- Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, et al. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci USA*. 2011; 108: 1513-8.
- Nouioui I, Ghodhbane-Gtari F, Montero-Calasanz MD, Goker M, Meier-Kolthoff JP, Schumann P, et al. Proposal of a type strain for *Frankia alni* (Woronin 1866) Von Tubeuf 1895, emended description of *Frankia alni*, and recognition of *Frankia casuarinae* sp. nov. and *Frankia elaeagni* sp. nov. *Int J Syst Evol Microbiol*. 2016; 66: 5201-10.
- Tisa LS, Oshone R, Sarkar I, Ktari A, Sen A, Gtari M. Genomic approaches toward understanding the actinorhizal symbiosis: an update on the status of the *Frankia* genomes. *Symbiosis*. 2016; 70: 5-16.
- Leul M, Normand P, Sellstedt A. The organization, regulation and phylogeny of uptake hydrogenase genes in *Frankia*. *Physiol Plantarum*. 2007; 130: 464-70.
- Blin K, Medema MH, Kazempour D, Fischbach MA, Breitling R, Takano E, et al. antiSMASH 2.0-a versatile platform for genome mining of secondary metabolite producers. *Nucleic Acids Res*. 2013; 41: W204-W212.
- Weber T, Blin K, Duddela S, Krug D, Kim HU, Brucoleri R, et al. antiSMASH 3.0-a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res*. 2015; 43: W237-W243.
- Udwaray DW, Gontang EA, Jones AC, Jones CS, Schultz AW, Winter JM, et al. Significant Natural Product Biosynthetic Potential of Actinorhizal Symbionts of the Genus *Frankia*, as Revealed by Comparative Genomic and Proteomic Analyses. *Appl Environ Microb*. 2011; 77: 3617-25.
- Ngom M, Oshone R, Hurst SGT, Abebe-Akele F, Simpson S, Morris K, et al. Permanent Draft Genome Sequence for *Frankia* sp. Strain CeD, a Nitrogen-Fixing Actinobacterium Isolated from the Root Nodules of *Casuarina equisetifolia* Grown in Senegal. *Genome announcements*. 2016; 4.
- Oshone R, Ngom M, Abebe-Akele F, Simpson S, Morris K, Sy MO, et al. Permanent Draft Genome Sequence of *Frankia* sp. Strain Allo2, a Salt-Tolerant Nitrogen-Fixing Actinobacterium Isolated from the Root Nodules of *Allocastrum*. *Genome announcements*. 2016; 4.
- Hurst SGT, Oshone R, Ghodhbane-Gtari F, Morris K, Abebe-Akele F, Thomas WK, et al. Draft Genome Sequence of *Frankia* sp. Strain Thr, a Nitrogen-Fixing Actinobacterium Isolated from the Root Nodules of *Casuarina cunninghamiana* Grown in Egypt. *Genome announcements*. 2014; 2: e00493-14.
- Ghodhbane-Gtari F, Hurst SGT, Oshone R, Morris K, Abebe-Akele F, Thomas WK, et al. Draft Genome Sequence of *Frankia* sp. Strain BMG5.23, a Salt-Tolerant Nitrogen-Fixing Actinobacterium Isolated

- from the Root Nodules of *Casuarina glauca* Grown in Tunisia. Genome announcements. 2014; 2: e00520-14.
24. Mansour SR, Oshone R, Hurst SG, Morris K, Thomas WK, Tisa LS. Draft Genome Sequence of *Frankia* sp. Strain Ccl6, a Salt-Tolerant Nitrogen-Fixing Actinobacterium Isolated from the Root Nodule of *Casuarina cunninghamiana*. Genome announcements. 2014; 2.
 25. D'Angelo T, Oshone R, Abebe-Akele F, Simpson S, Morris K, Thomas WK, et al. Permanent Draft Genome Sequence of *Frankia* sp. Strain BR, a Nitrogen-Fixing Actinobacterium Isolated from the Root Nodules of *Casuarina equisetifolia*. Genome announcements. 2016; 4.