

University of New Hampshire

University of New Hampshire Scholars' Repository

New Hampshire Agricultural Experiment Station Publications New Hampshire Agricultural Experiment Station

3-1991

Fragaria vesca Chlorophyll Mutants

Thomas M. Davis

University of New Hampshire, Tom.Davis@unh.edu

James E. Pollard

University of New Hampshire - Main Campus, Jim.Pollard@unh.edu

Follow this and additional works at: <https://scholars.unh.edu/nhaes>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), [Fruit Science Commons](#), [Horticulture Commons](#), and the [Plant Breeding and Genetics Commons](#)

Recommended Citation

Fragaria vesca Chlorophyll Mutants Thomas M. Davis and James E. Pollard HortScience March 1991 26:311

This Article is brought to you for free and open access by the New Hampshire Agricultural Experiment Station at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in New Hampshire Agricultural Experiment Station Publications by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

Fragaria vesca Chlorophyll Mutants

Thomas M. Davis and James E. Pollard

Dept. of Plant Biology, University of New Hampshire, Durham,
NH 03824

Additional index words. strawberry, genetics, induced mutation

Despite the successful history of strawberry breeding (Bringhurst and Voth, 1984), relatively little is known of strawberry genetics (e.g., Arulsekhar et al., 1981; Mok and Evans, 1971). Genetic studies of the commercially cultivated strawberry, *Fragaria* × *ananassa* Duchesne (2N = 8X = 56), are hampered by genetic complexity at the octoploid level (Arulsekhar et al., 1981). Because the AAA'A'BBBB genomic constitution of *F.* × *ananassa* incorporates the AA genomes of diploids *F. vesca* and *F. viridis* (Senanayake and Bringhurst, 1967), genetic analysis of diploid strawberries may provide insight into the genetic constitution and behavior of the cultivated strawberry (Arulsekhar and Bringhurst, 1983; Brown and Waring, 1965).

Only a few morphological (Brown and Waring, 1965; Richardson, 1914, 1918) and isozyme (Arulsekhar et al., 1981) variants of diploid strawberries have been genetically characterized. We mutagenized *F. vesca* 'Baron Solemacher' with the objective of identifying new and useful genetic markers in a diploid strawberry.

Seeds of 'Baron Solemacher' (Johnny's Selected Seeds, Albion, Maine) were imbibed under moist cotton overnight, then exposed to a ¹³⁷Cs source for 60 min, receiving

an estimated γ -irradiation dose of 7.5 krad. M₂ and M₃ generation seed production resulted from natural self-pollination of M₁ and M₂ plants, respectively, under greenhouse conditions.

Among 90 M₁ families grown for evaluation, four contained stable chlorophyll mutants. In families 456, 502, and 648, mutant seedlings (mutants M456, M502, and M648, respectively) had light green, yellow, and white shoots, respectively, and died at the cotyledon or first true leaf stage. Light green seedlings of mutant M538 were viable, grew to sexual maturity, and produced only light green progeny when allowed to self-pollinate in the greenhouse.

Normally pigmented siblings in M₂ families 456, 502, 538, and 648 were all fully viable; a proportion (8/13, 6/15, 4/15, and 7/15, respectively) of these normal M₂ plants produced M₃ families that segregated for the respective chlorophyll abnormalities, showing that each mutant phenotype was inherited in a recessive manner.

When M₁ phenotypic segregation patterns (within segregating M₃ families) were tested for goodness-of-fit to a 3:1 ratio, only mutant M502 [521 normal: 191 yellow (X² = 1.17 n.s.)] fit the monogenic expectation. Observed deviations from 3:1 for M456 [547 normal : 122 light green (X² = 15.8**)], M538 [109 normal : 55 light green (X² = 5.93*)], and M648 [376 normal : 52 white (X² = 37.0**)] may have been due to gametic or zygotic selection against (M456, M648), or in favor of (M538) the mutant type, or perhaps to multiple factor inheritance.

Mutants M456, M502, M538, and M648, which can all be phenotypically classified at seedling emergence, will be useful in linkage and other genetic studies in strawberry. Additionally, the M538 mutation has potential value as a marker for estimation of outcrossing rates. *F. vesca* is regarded as a predominantly self-pollinating species (Arulsekhar and Bringhurst, 1981). However, when seeds were harvested from M538 plants grown in a field among a population of normal 'Baron Solemacher' plants, 240 out of 522 of the resulting seedlings were dark green. Whether this apparent outcrossing rate of 46% was due to specific reproductive characteristics of mutant M538 or was truly representative of normal 'Baron Solemacher' remains to be determined. Mutants M456, M502, M538, and M648 are available to interested researchers upon request to the authors.

Literature Cited

- Arulsekhar, S. and R.S. Bringhurst. 1983. Strawberry, p. 391-400. In: S.D. Tanksley and T.J. Orton (eds.) Isozymes in plant genetics and breeding, part B. Elsevier, Amsterdam.
- Arulsekhar, S., R.S. Bringhurst, and V. Voth. 1981. Inheritance of PGI and LAP isozymes in octoploid cultivated strawberries. *J. Amer. Soc. Hort. Sci.* 16:679-683.
- Bringhurst, R.S. and V. Voth. 1984. Breeding octoploid strawberries. *Iowa State J. of Res.* 58:371-381.
- Brown, T. and P.F. Waring. 1965. The genetical control of the everbearing habit and three other characters of *Fragaria vesca*. *Euphytica* 14:97-112.
- Mok, D.W.S. and W.D. Evans. 1971. Chromosome association at diakinesis in the cultivated strawberry. *Can. J. Genet. Cytol.* 13:231-236.
- Richardson, C.W. 1914. A preliminary note on the genetics of *Fragaria*. *J. Genet.* 3:171-177.
- Richardson, C.W. 1918. A further note on the genetics of *Fragaria*. *J. Genet.* 7:167-170.
- Senanayake, Y.D.A. and R.S. Bringhurst. 1967. Origin of *Fragaria* polyploids. I. Cytological analysis. *Amer. J. Bot.* 54:221-228.

Received for publication 24 Aug. 1989. Scientific contribution no. 1640 from the New Hampshire Agricultural Experiment Station. The cost of publishing this paper was defrayed in part by the payment of page charges. Under postal regulations, this paper therefore must be hereby marked *advertisement* solely to indicate this fact.