

University of New Hampshire

University of New Hampshire Scholars' Repository

New Hampshire Agricultural Experiment Station Publications New Hampshire Agricultural Experiment Station

8-4-2016

Permanent draft genome sequence of *Nocardia* sp. BMG111209, an actinobacterium isolated from nodules of *Casuarina glauca*

Faten Ghodhbane-Gtari
Université Tunis El Manar

Nicholas Beauchemin
University of New Hampshire, Durham

Abdellatif Gueddou
Université Tunis El Manar

Karima Hezbri
Université Tunis El Manar

Moussa Louati
Université Tunis El Manar

Follow this and additional works at: <https://scholars.unh.edu/nhaes>

See next page for additional authors

Comments

This is an article published by American Society for Microbiology in Genome Announcements in 2016, available online: <https://dx.doi.org/10.1128/genomeA.00770-16>

Recommended Citation


Ghodhbane-Gtari F, Beauchemin N, Gueddou A, Hezbri K, Ktari A, Louati M, Nouioui I, Chen A, Huntemann M, Ivanova N, Kyrpides N, Markowitz V, Mavrommatis K, Pagani I, Sen A, Wall L, Woyke T, Gtari M, Tisa LS. 2016. Permanent draft genome sequence of *Nocardia* sp. BMG111209, an actinobacterium isolated from nodules of *Casuarina glauca*. *Genome Announc* 4(4):e00770-16. <https://dx.doi.org/10.1128/genomeA.00770-16>

This Article is brought to you for free and open access by the New Hampshire Agricultural Experiment Station at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in New Hampshire Agricultural Experiment Station Publications by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

Authors

Faten Ghodhbane-Gtari, Nicholas Beauchemin, Abdellatif Gueddou, Karima Hezbri, Moussa Louati, Imen Nouioui, Amy Chen, Marcel Huntemann, Natalia Ivanova, Nikos Kyripides, Victor Markowitz, Kostas Mavrommatis, Ioanna Pagani, Arnab Sen, Luis Wall, Tanja Woyke, Maher Gtari, and Louis S. Tisa

Permanent Draft Genome Sequence of *Nocardia* sp. BMG111209, an Actinobacterium Isolated from Nodules of *Casuarina glauca*

Faten Ghodhbane-Gtari,^a Nicholas Beauchemin,^b Abdellatif Gueddou,^a Karima Hezbri,^a Amir Ktari,^a Moussa Louati,^a Imen Nouioui,^a Amy Chen,^c Marcel Huntemann,^c Natalia Ivanova,^c Nikos Kyrpides,^c Victor Markowitz,^c Kostas Mavrommatis,^c Ioanna Pagani,^c Arnab Sen,^d Luis Wall,^e Tanja Woyke,^c  Maher Gtari,^a Louis S. Tisa^b

University of Tunis-El Manar, Tunis, Tunisia^a; University of New Hampshire, Durham, New Hampshire, USA^b; DOE Joint Genome Institute, Walnut Creek, California, USA^c; University of North Bengal, Siliguri, India^d; University of Quilmes, Quilmes, Argentina^e

This is scientific contribution number 2677.

***Nocardia* sp. strain BMG111209 is a non-*Frankia* actinobacterium isolated from root nodules of *Casuarina glauca* in Tunisia. Here, we report the 9.1-Mbp draft genome sequence of *Nocardia* sp. strain BMG111209 with a G + C content of 69.19% and 8,122 candidate protein-encoding genes.**

Received 8 June 2016 Accepted 10 June 2016 Published 4 August 2016

Citation Ghodhbane-Gtari F, Beauchemin N, Gueddou A, Hezbri K, Ktari A, Louati M, Nouioui I, Chen A, Huntemann M, Ivanova N, Kyrpides N, Markowitz V, Mavrommatis K, Pagani I, Sen A, Wall L, Woyke T, Gtari M, Tisa LS. 2016. Permanent draft genome sequence of *Nocardia* sp. BMG111209, an actinobacterium isolated from nodules of *Casuarina glauca*. *Genome Announc* 4(4):e00770-16. doi:10.1128/genomeA.00770-16.

Copyright © 2016 Ghodhbane-Gtari et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Louis S. Tisa, louis.tisa@unh.edu.

Actinorhizal plants are well known for their symbiotic association with the actinobacteria, *Frankia*, which results in the formation of root nodule structures (1). Besides hosting these bacterial symbionts, root nodules have other occupants or plant endophytes that seem to aid plant growth and health. Many non-*Frankia* actinobacteria have been isolated from actinorhizal root nodules, occupying the same microniche as *Frankia* (2–14). These non-*Frankia* actinobacteria appear to have functional roles including plant growth-promoting activity (4). As early as 1981, *Nocardia* species were found associated with actinorhizal nodules (7, 8). *Nocardia* sp. strain BMG111209 was isolated from the root nodules of *Casuarina glauca* grown in Tunisia (2). This isolate was sequenced to provide a greater understanding of the metabolic potential of this microbe and its interaction with actinorhizal plants.

The draft genome of *Nocardia* sp. strain BMG111209 was generated at the DOE Joint genome Institute (JGI) using a hybrid of Illumina (15) and Pacific Biosciences (PacBio) technologies. An Illumina Std. shotgun library and long insert mate-pair library were constructed and sequenced using the Illumina HiSeq 2000 platform. This procedure generated 51,070,879 reads totaling 7,660.6 Mbp from the standard shotgun library and 47,626,505 reads totaling 4,134.5 Mb from the long insert mate-pair library. A PacBio SMRTbell library was constructed and sequenced on the PacBio RS platform, which generated 250,559 raw PacBio reads yielding 166,907 adapter trimmed and quality filtered subreads totaling 471.4 Mb. All techniques for DNA isolation, library construction, and sequencing were performed according to JGI standards and protocols (<http://www.jgi.doe.gov>). The Illumina and PacBio sequence data were assembled using Allpaths-LG (version r42328) (16). The final draft assembly contained six contigs in five scaffolds.

The total size of the genome is 9.1 Mbp. The final assembly is based on 7,660.9 Mbp of Illumina Standard PE, 4,143.5 Mb of Illumina CLIP and 471.4 Mb of PacBio post filtered data, which provides an average 1297.2× Illumina coverage and 51.8× PacBio coverage of the genome, respectively.

The assembled *Nocardia* BMG111209 genome was annotated using the JGI annotation pipeline (17, 18) and the data are available from the IMG data management system (19). The draft genome of *Nocardia* BMG111209 was resolved into 5 scaffolds consisting of 9,143,142 bp with a G + C content of 69.19%, 8,122 candidate protein-encoding genes, 52 tRNA genes, and 2 rRNA regions. Analysis of this genome for metabolism potential revealed phytohormone (i.e., IAA) biosynthesis pathways and the presence of a *hup* operon encoding an Ni-hydrogenase suggesting possible roles in their functioning as a plant-endophyte.

Nucleotide sequence accession numbers. This whole-genome shotgun sequence has been deposited at DDBJ/EMBL/GenBank under the accession number [ARMU000000000](https://www.ncbi.nlm.nih.gov/nuclink/ARMU000000000). The version described in this paper is the first version, ARMU01000000.

ACKNOWLEDGMENTS

Partial funding by the New Hampshire Agricultural Experiment Station. The work conducted by the U.S. Department of Energy Joint Genome Institute was supported by the Office of Science of the U.S. Department of Energy under contract DE-AC02-05CH11231. This project (L.S.T.) was supported in part by the USDA National Institute of Food and Agriculture, Hatch 022821, and The College of Life Sciences and Agriculture at the University of New Hampshire, Durham, NH. M.G. and F.G.-G. were supported in part by a Visiting Scientist and Postdoctoral Scientist Program administered by the NH AES at the University of New Hampshire.

FUNDING INFORMATION

This work, including the efforts of Louis S. Tisa, was funded by USDA | National Institute of Food and Agriculture (NIFA) (Hatch 022821). This work, including the efforts of Tanja Woyke, was funded by U.S. Department of Energy (DOE) (DE-AC02--05CH11231).

REFERENCES

1. Normand P, Benson DR, Berry AM, Tisa LS. 2014. Family *Frankiaceae*, p. 339–356. In Rosenberg E, DeLong EF, Lory S, Stackebrandt E, Thompson F (ed), *The prokaryote—Actinobacteria*. Springer-Verlag, Berlin, Germany.
2. Ghodhbane-Gtari F, Essoussi I, Chattaoui M, Chouaia B, Jaouani A, Daffonchio D, Boudabous A, Gtari M. 2010. Isolation and characterization of non-*Frankia* *Actinobacteria* from root nodules of *Alnus glutinosa*, *Casuarina glauca* and *Elaeagnus angustifolia*. *Symbiosis* 50:51–57. <http://dx.doi.org/10.1007/s13199-009-0029-7>.
3. Ghodhbane-Gtari F, Nouioui I, Salem K, Ktari A, Montero-Calasanz Mdel C, Tisa LS, Klenk HP, Gtari M. 2014. *Nocardia casuarinae* sp. nov., an actinobacterial endophyte isolated from root nodules of *Casuarina glauca*. *Antonie Van Leeuwenhoek* 105:1099–1106. <http://dx.doi.org/10.1007/s10482-014-0168-6>.
4. Ghodhbane-Gtari F, Tisa LS. 2014. Ecology and physiology of non-*Frankia* *Actinobacteria* from actinorhizal plants, p. 27–42. In Katsey EI (ed), *Plasticity in plant-growth-promoting and phytopathogenic Bacteria*. Springer, New York, NY.
5. Solans M, Vobis G. 2003. Actinomycetes saprofiticos asociados a la rizósfera de *Discaria trinervis*. *Ecol Aust* 13:97–107.
6. Solans M, Vobis G, Cassán F, Luna V, Wall LG. 2011. Production of phytohormones by root-associated saprophytic actinomycetes isolated from the actinorhizal plant *Ochetophila trinervis*. *World J Microbiol Biotechnol* 27:2195–2202. <http://dx.doi.org/10.1007/s11274-011-0685-7>.
7. Malishkaite YB, Nechaeva GA, Kuimova TF, Evtushenko LI, Agre NS. 1987. *Nocardia* strains from actinorhizal plant nodules. *Microbiology* 56: 100–106.
8. Pariiskaya AN, Novik SN, Agre NS, Kalakutskii LV. 1982. *Nocardia* Isolated from nitrogen-fixing nodules on the roots of the gray and black alder. *Microbiology* 51:117–121.
9. Carro L, Pujic P, Trujillo ME, Normand P. 2013. *Micromonospora* is a normal occupant of actinorhizal nodules. *J Biosci* 38:685–693. <http://dx.doi.org/10.1007/s12038-013-9359-y>.
10. Trujillo ME, Kroppenstedt RM, Schumann P, Carro L, Martínez-Molina E. 2006. *Micromonospora coriariae* sp. nov., isolated from root nodules of *Coriaria myrtifolia*. *Int J Syst Evol Microbiol* 56:2381–2385. <http://dx.doi.org/10.1099/ijs.0.64449-0>.
11. Trujillo ME, Riesco R, Benito P, Carro L. 2015. Endophytic Actinobacteria and the interaction of *Micromonospora* and nitrogen fixing plants. *Front Microbiol* 6:1341. <http://dx.doi.org/10.3389/fmicb.2015.01341>.
12. Niner BM, Brandt JP, Villegas M, Marshall CR, Hirsch AM, Valdés M. 1996. Analysis of partial sequences of genes coding for 16S rRNA of actinomycetes isolated from *Casuarina equisetifolia* nodules in Mexico. *Appl Environ Microbiol* 62:3034–3036.
13. Valdés D, Huss-Danell K, Lavire C, Normand P, Wall L. 2006. Further characterization of new symbiotic nitrogen fixing non-*Frankia* actinomycetes isolated from nodules of *Alnus acuminata*. In 14th International Meeting on Frankia and Actinorhizal Plants. Umea University, Umea, Sweden.
14. Valdés M, Pérez NO, Estrada-de Los Santos P, Caballero-Mellado J, Peña-Cabiales JJ, Normand P, Hirsch AM. 2005. Non-*Frankia* actinomycetes isolated from surface-sterilized roots of *Casuarina equisetifolia* fix nitrogen. *Appl Environ Microbiol* 71:460–466. <http://dx.doi.org/10.1128/AEM.71.1.460-466.2005>.
15. Bennett S. 2004. Solexa Ltd. *Pharmacogenomics* 5:433–438. <http://dx.doi.org/10.1517/14622416.5.4.433>.
16. Gnerre S, MacCallum I, Przybylski D, Ribeiro FJ, Burton JN, Walker BJ, Sharpe T, Hall G, Shea TP, Sykes S, Berlin AM, Aird D, Costello M, Daza R, Williams L, Nicol R, Gnirke A, Nusbaum C, Lander ES, Jaffe DB. 2011. High-quality draft assemblies of mammalian genomes from massively parallel sequence data. *Proc Natl Acad Sci U S A* 108:1513–1518. <http://dx.doi.org/10.1073/pnas.1017351108>.
17. Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Palaniappan K, Szeto E, Pillay M, Chen IMA, Pati A, Nielsen T, Markowitz VM, Kyrpides NC. 2015. The standard operating procedure of the DOE-JGI microbial genome annotation pipeline (MGAP v.4). *Stand Genomic Sci* 10:86.
18. Markowitz VM, Mavromatis K, Ivanova NN, Chen IM, Chu K, Kyrpides NC. 2009. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 25:2271–2278. <http://dx.doi.org/10.1093/bioinformatics/btp393>.
19. Markowitz VM, Korzeniewski F, Palaniappan K, Szeto E, Werner G, Padki A, Zhao X, Dubchak I, Hugenholtz P, Anderson I, Lykidis A, Mavromatis K, Ivanova N, Kyrpides NC. 2006. The integrated microbial genomes (IMG) system. *Nucleic Acids Res* 34:D344–D348. <http://dx.doi.org/10.1093/nar/gkj024>.