

3-2015

Scanning tunneling microscopy studies of graphene and hydrogenated graphene on Cu

Shawna M. Hollen

University of New Hampshire - Main Campus

G A. Gambrel

Ohio State University - Main Campus

S.J. Tjung

Ohio State University - Main Campus

N M. Santagata

Ohio State University - Main Campus

Ezekiel Johnston-Halperin

Ohio State University - Main Campus

See next page for additional authors

Follow this and additional works at: https://scholars.unh.edu/physics_facpub

 Part of the [Physics Commons](#)

Recommended Citation

Hollen, S.-M., Gambrel, G., Tjung, S., et al. \ 2015, APS Meeting Abstracts, 17003. <http://meetings.aps.org/link/BAPS.2015.MAR.W17.3>

This Conference Proceeding is brought to you for free and open access by the Physics at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Physics Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact nicole.hentz@unh.edu.

Authors

Shawna M. Hollen, G A. Gambrel, S.J. Tjung, N M. Santagata, Ezekiel Johnston-Halperin, and Jay A. Gupta

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Scanning tunneling microscopy studies of graphene and hydrogenated graphene on Cu(111)¹ SHAWNA M. HOLLEN, GRADY GAMBREL, STEVEN TJUNG, NANCY M. SANTAGATA, EZEKIEL JOHNSTON-HALPERIN, JAY A. GUPTA, The Ohio State University — Because of the innate sensitivity of 2D material surfaces, it is increasingly important to understand and characterize surface functionalization and interactions with environmental elements, such as substrate, metallic contacts, and adatoms. We developed a method for reproducible, epitaxial growth of pristine graphene islands on Cu(111) in UHV and use scanning tunneling microscopy and spectroscopy (STM) to study the interaction of these graphene islands with the Cu substrate. Tunneling spectroscopy measurements of the electronic surface states over the graphene islands indicate a lower local work function, decreased coupling to bulk Cu states, and a decreased electron effective mass. Additionally, we developed a novel field electron dissociation technique to form hydrogen-terminated graphene at low temperatures and in UHV. This method produced what may be the first STM images of crystalline hydrogenated graphene. The pristine graphene island is then recovered by scanning at a high tip-sample bias. The hydrogenation and its reversibility suggest writing lateral 2D devices using the STM tip. Toward this end, we are developing the capability to repeat the hydrogenation on working graphene devices.

¹This research is supported by the Center for Emergent Materials NSF-funded MRSEC (DMR-0820414)

Shawna Hollen
Ohio State Univ - Columbus

Date submitted: 14 Nov 2014

Electronic form version 1.4