

University of New Hampshire

University of New Hampshire Scholars' Repository

Physics Scholarship

Physics

3-2015

Transport in thin insulating films close to the Boson-Fermion Crossover

J C. Joy
Brown University

X Zhang
Brown University

Shawna M. Hollen
University of New Hampshire - Main Campus

C Zhao
Brown University

G. E. Fernandes
Brown University

See next page for additional authors

Follow this and additional works at: https://scholars.unh.edu/physics_facpub



Part of the [Physics Commons](#)

Recommended Citation

<http://meetings.aps.org/link/BAPS.2015.MAR.M25.11>

This Conference Proceeding is brought to you for free and open access by the Physics at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Physics Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

Authors

J C. Joy, X Zhang, Shawna M. Hollen, C Zhao, G. E. Fernandes, J. M. Xu, and J M. Valles Jr.

Abstract Submitted
for the MAR15 Meeting of
The American Physical Society

Transport in thin insulating films close to the Boson-Fermion Crossover J.C. JOY, X. ZHANG, S.M. HOLLEN¹, C. ZHAO², Department of Physics, Brown University, G. FERNANDES, J.M. XU, Division of Engineering, Brown University, J.M. VALLES, JR., Department of Physics, Brown University — In two-dimensional systems, sufficient levels of disorder are known to localize Cooper Pairs into a phase incoherent insulating state. While many theoretical and experimental works have shown this state's existence, its ubiquity close to the disorder tuned Superconductor to Insulator transition is still an open problem. Recent experiments on nanopatterned $\text{Pb}_{0.9}\text{Bi}_{0.1}$ films have suggested a crossover from Bosonic to Fermionic transport deep in the insulating phase, indicating that the Cooper Pair Insulator (CPI) only persists to a finite level of microscopic disorder. The normal state resistance at which this crossover occurs is governed by the extent coupling constant inhomogeneities on the scale of the coherence length, which allow the formation of locally phase coherent superconducting islands in the insulating state. By tuning the scale of these inhomogeneities and examining the extent of the CPI state, we argue that the disorder tuned Superconductor to Insulator transition proceeds via pair breaking and Anderson localization of fermions when the level spacing in the islands approaches the size of the mean field gap. This work was supported by the NSF through grants No. DMR-1307290 and DMR-0907357 and by the AFRL, the ONR, and the AFOSR.

¹Currently at the Center for Emergent Materials, Ohio State University

²Currently at Northwestern Polytechnical University, Xian, China

James Joy
Department of Physics, Brown University

Date submitted: 14 Nov 2014

Electronic form version 1.4