

University of New Hampshire

## University of New Hampshire Scholars' Repository

---

Faculty Publications

---

5-15-2003

### Production of methyl bromide in a temperate forest soil

Ruth K. Varner

*University of New Hampshire, Durham, ruth.varner@unh.edu*

Marguerite L. White

*University of New Hampshire, Durham*

Cindy H. Mosedale

*University of New Hampshire, Durham*

Patrick M. Crill

*Stockholm University*

Follow this and additional works at: [https://scholars.unh.edu/faculty\\_pubs](https://scholars.unh.edu/faculty_pubs)

---

#### Recommended Citation

Varner, R.K., M. White, C.H. Mosedale and P.M. Crill, (2003) Production of methyl bromide in a temperate forest soil, *Geophys. Res. Lett.*, 30, 10.1029/2002GL016592.

This Article is brought to you for free and open access by University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact [Scholarly.Communication@unh.edu](mailto:Scholarly.Communication@unh.edu).

## Production of methyl bromide in a temperate forest soil

Ruth K. Varner, Marguerite L. White, Cindy H. Mosedale, and Patrick M. Crill

Complex Systems Research Center, Institute for the Study of Earth, Oceans and Space, University of New Hampshire, Durham, New Hampshire, USA

Received 11 November 2002; revised 14 February 2003; accepted 2 April 2003; published 23 May 2003.

[1] Field enclosure measurements of a temperate forest soil show net uptake of ambient methyl bromide ( $\text{CH}_3\text{Br}$ ), an important trace gas in both tropospheric and stratospheric ozone cycling. The net flux for 1999 was estimated to be  $-168 \pm 72 \mu\text{g CH}_3\text{Br m}^{-2}$  (negative indicates loss from the atmosphere). Individual enclosure flux measurements ranged from  $-4.0$  to  $+3.3 \mu\text{g CH}_3\text{Br m}^{-2} \text{d}^{-1}$ . Soil consumption of  $\text{CH}_3\text{Br}$  was estimated from laboratory soil incubations. Production of  $\text{CH}_3\text{Br}$  was calculated as the difference between net flux and predicted consumption. Fungi could be responsible for the production of  $\text{CH}_3\text{Br}$  in this temperate forest soil. **INDEX TERMS:** 0315 Atmospheric Composition and Structure: Biosphere/atmosphere interactions; 0322 Atmospheric Composition and Structure: Constituent sources and sinks; 1615 Global Change: Biogeochemical processes (4805); 0330 Atmospheric Composition and Structure: Geochemical cycles; 9350 Information Related to Geographic Region: North America. **Citation:** Varner, R. K., M. L. White, C. H. Mosedale, and P. M. Crill, Production of methyl bromide in a temperate forest soil, *Geophys. Res. Lett.*, 30(10), 1521, doi:10.1029/2002GL016592, 2003.

### 1. Introduction

[2] The tropospheric budget of  $\text{CH}_3\text{Br}$  is out of balance with sinks exceeding sources by  $59 \text{ Gg yr}^{-1}$  [Yvon-Lewis, 2000]. Natural sources and sinks are of particular concern because significant gaps remain in our understanding of ecosystem  $\text{CH}_3\text{Br}$  cycling.

[3] Several terrestrial sources of  $\text{CH}_3\text{Br}$  have been identified [Gan *et al.*, 1998; Varner *et al.*, 1999b; Redeker *et al.*, 2000; Rhew *et al.*, 2000; Dimmer *et al.*, 2001; Rhew *et al.*, 2001]. The production mechanism of  $\text{CH}_3\text{Br}$  in these ecosystems is uncertain. Abiotic production of methyl halides can occur during the oxidation of organic matter [Keppler *et al.*, 2000]. Leaf disc studies of a variety of plants, including *Brassica* [Gan *et al.*, 1998], have shown that enzyme mediated methyl transferase can produce  $\text{CH}_3\text{Br}$  [Attieh *et al.*, 1995; Saini *et al.*, 1995]. Wood rotting fungi and ectomycorrhizal fungi are also potential sources of  $\text{CH}_3\text{Br}$  in these ecosystems [Harper, 1985; Lee-Taylor and Holland, 2000 (L-TH2000); Redeker *et al.*, unpublished (KR2003)].

[4] Consumption and production of  $\text{CH}_3\text{Br}$  may be occurring simultaneously in these ecosystems. Jeffers *et al.* [1998] report a variety of leaves consume elevated levels of  $\text{CH}_3\text{Br}$ . Soil has also been identified as a sink of atmospheric  $\text{CH}_3\text{Br}$  [Shorter *et al.*, 1995; Serça *et al.*, 1998]. Hines *et al.* [1998] determined the process to be aerobic bacterial uptake. Bacteria that consume fumigant

and ambient levels of  $\text{CH}_3\text{Br}$  have been isolated from soil [e.g., Connell Hancock *et al.*, 1998; Miller *et al.*, 1997; Goodwin *et al.*, 2001].

[5] This paper examines production and consumption of  $\text{CH}_3\text{Br}$  in a temperate forest soil in New Hampshire. Field measurements of soil-atmosphere exchange of ambient  $\text{CH}_3\text{Br}$  were completed from May 28 to October 28, 1999. Rates of  $\text{CH}_3\text{Br}$  consumption in the soil were estimated with a model derived from temperature and moisture manipulated soil incubations. We estimated production in the soil as the difference between measured net flux from field measurements and the modeled consumption based on soil incubations.

### 2. Methods

#### 2.1. Field Measurements

[6] College Woods ( $43^\circ 08' \text{N}$ ,  $71^\circ 57' \text{W}$ ), Durham, NH is a mixed deciduous conifer forest abandoned as a woodlot approximately 110 years ago. Soils are well drained, weakly to moderately acidic inceptisols with a thin, variable litter layer. A dark organic rich layer extended from 0 to 5 cm then transitioned into light brown mineral soil below 5 cm. [Crill, 1991].


[7] Field enclosure measurements were made on a near weekly basis from May 28 to September 9 with two additional measurements on September 24 and October 28. Measurements were completed at two aluminum collars previously established at the site in 1989 [Crill, 1991]. One was located on the slope of a small hill while the other collar was in a hollow approximately 5 m away. There was no above-ground vegetation in the collars.

[8] An aluminum enclosure ( $0.152 \text{ m}^3$  volume), with a fan mounted inside to mix the headspace, was placed on the collar and sealed with water. Four headspace samples ( $2.5 \text{ L}$ ) were collected every 5 minutes. The gas samples were collected in stainless-steel electropolished cylinders and analyzed for  $\text{CH}_3\text{Br}$  by GC-ECD as described in Kerwin *et al.* [1996]. Artifacts due to enclosure configuration were below the analytical limit of detection.

[9] Soil samples of litter, 0–5 cm (organic layer) and 5–10 cm (mineral layer) were collected. Soil moisture was calculated as soil weight loss after oven drying at  $75^\circ \text{C}$  for 24 hrs divided by the dry weight of the sample. Air and soil surface, 5 and 10 cm temperatures were measured manually while datalogger recorded hourly-averaged air, 2, 8, and 15 cm soil temperatures from thermistors.

#### 2.2. Laboratory Incubations

[10] Static soil incubations were performed to determine the consumption rate of  $\text{CH}_3\text{Br}$ . Soil samples were collected from College Woods, stored at  $4^\circ \text{C}$  in air tight plastic bags



**Figure 1.** A. Daily total precipitation (bars), hourly average -8 cm soil temperature (solid line) and soil moisture (squares and dotted line) for 1999. B. CH<sub>3</sub>Br flux measurements for College Woods for the two collars. Error bars are the error of the linear regression of the concentration versus time flux data. The cross indicates days when Br<sup>-</sup> ion in the soil was measured.

and were processed within 1 week of collection. For more details on the sampling and analysis method see Kerwin *et al.* [1996]. A reaction rate constant,  $k$  (min<sup>-1</sup>), was determined as the slope of the regression fit of the natural log of nmoles of CH<sub>3</sub>Br versus time. Uptake rate constants were determined for the soil at 5, 15, 25, 35 and 45°C and for moisture contents ranging from 26.3% to 344%.

### 3. Results

#### 3.1. Field Measurements

[11] The mean flux for the site was  $-0.70 \pm 0.31$  μg CH<sub>3</sub>Br m<sup>-2</sup> d<sup>-1</sup> (Figure 1B). Seasonal flux was calculated based on a 240 day growing season as  $-168 \pm 72$  μg CH<sub>3</sub>Br m<sup>-2</sup>.

#### 3.2. Soil Incubations

[12] By applying a Gaussian fit to the soil incubation data (Figure 2), the predictor equation for  $k$  becomes

$$-k = 1.24e^{-0.5 \left[ \left( \frac{x-26.85}{18.57} \right)^2 + \left( \frac{y-194.34}{77.80} \right)^2 \right]} \quad (1)$$


$x$  and  $y$  are soil temperature (°C) and soil moisture, respectively. This fit resulted in an  $r^2 = 0.853$  and an estimate of error for  $k$  of  $\pm 0.21$  min<sup>-1</sup>.

[13] The model was then used to estimate field uptake rate constants using soil moisture and temperature data collected during the 1999 sampling season. Soil consumption of CH<sub>3</sub>Br for each day of sampling was calculated from Varner *et al.* [1999a]. A production estimate of CH<sub>3</sub>Br was calculated for 5 sampling days as the difference between the measured or net flux and the estimated consumption (Table 1).

### 4. Discussion

#### 4.1. CH<sub>3</sub>Br Flux Measurements

[14] These are the first seasonal field measurements of CH<sub>3</sub>Br exchange in a temperate forest. The net measure-




**Figure 2.** College Woods 0–3 cm soil temperature and moisture manipulations. Uptake rate constant versus %soil moisture content at 25°C (■) and temperature at 128.5% soil moisture (●).

ments range from  $-3.0$  to  $+4.0$  μg CH<sub>3</sub>Br m<sup>-2</sup> d<sup>-1</sup> and overlap the range seen in the more arid shrubland environments in Southern California ( $-0.95$  to  $+14.7$  μg m<sup>-2</sup> d<sup>-1</sup>) [Rhew *et al.*, 2001]. High moisture and organic matter content in the soil could account for higher rates of consumption [Hines *et al.*, 1998]. The soil moisture of the Rhew *et al.* [2001] sites ranged from 0.3 to 24%. We measured a minimum moisture content of 50%. Organic matter content data was not reported for the shrubland study. Our site ranged in organic matter from 78.5% (0–5 cm) to 17.4% (5–10 cm).

#### 4.2. Consumption and Production Estimates

[15] Using laboratory-derived rates of consumption in a field setting brings with it many uncertainties. The bulk



**Figure 3.** Modeled versus measured consumption of CH<sub>3</sub>Br for field data from 1994.

**Table 1.** Measured Net Flux, Modeled Consumption and Estimated Production of CH<sub>3</sub>Br in College Woods Soils

| Sampling Date | Surface soil Br <sup>-</sup> (mM) | Measured Net Flux | Modeled Consumption | Field Production | EF high and low (SD)        | WF (Range)      |
|---------------|-----------------------------------|-------------------|---------------------|------------------|-----------------------------|-----------------|
| 05/28         | 0.05                              | 0.5 ± 0.16        | -11.0 ± 5.9         | 11.5 ± 5.9       | 2.6 - 5E-05 (0.7) (1.5E-05) | 0.28 (0.9-0.08) |
| 06/01         | 0.09                              | -4.0 ± 1.1        | -4.0 ± 5.9          | 0.1 ± 6.0        | 3.1 - 5E-05 (0.9) (1.5E-05) | 0.17 (0.5-0.05) |
| 06/30         | 0.09                              | 3.3 ± 0.9         | -15.9 ± 5.9         | 19.2 ± 6.0       | 3.0 - 5E-05 (0.9) (1.5E-05) | 0.53 (1.6-0.14) |
| 07/21         | 0.16                              | -0.3 ± 0.4        | -9.1 ± 5.9          | 8.7 ± 5.9        | 4.0 - 5E-05 (1.2) (1.5E-05) | 0.69 (2.1-0.19) |
| 08/23         | 0.20                              | 0.1 ± 0.66        | -4.2 ± 5.9          | 4.3 ± 5.9        | 4.6 - 5E-05 (1.3) (1.5E-05) | 0.44 (1.3-0.12) |

All numbers reported in  $\mu\text{g m}^{-2} \text{d}^{-1}$ . Italicized data are calculated values. SD = standard deviation.

density, temperature, moisture and biological activity will be different than that encountered in the field. There have been successful attempts to estimate field mechanisms of NO production and consumption from laboratory measurements [Galbally and Johansson, 1989; van Dijk et al., 2002]. Furthermore, we applied our laboratory derived model to field measurement and see a reasonable estimate of field consumption (Figure 3). Finally, our consumption estimates for the 1999 field season (2.6 and 19.2  $\mu\text{g CH}_3\text{Br m}^{-2} \text{d}^{-1}$ ) encompass the rate reported by Serça et al. [1995] for a temperate forest soil in Colorado.

[16] Soil production estimates calculated from the difference between net field measurements and the modeled consumption in the forest soil range from 0.05 to 19.2  $\mu\text{g CH}_3\text{Br m}^{-2} \text{d}^{-1}$ . These overlap with the range of net positive flux measurements reported by Rhew et al. [2001] for Southern California shrubland (0.03–14.7  $\mu\text{g m}^{-2} \text{d}^{-1}$ ) and Dimmer et al. [2001] for a conifer forested peatland in Ireland (0.08 to 18  $\mu\text{g m}^{-2} \text{d}^{-1}$ ).

### 4.3. Sources of CH<sub>3</sub>Br

[17] Production of CH<sub>3</sub>Br in a temperate forest soil could be the result of abiotic [Keppler et al., 2000], fungal [Harper, 1985; L-TH2000, KR2003] or other unidentified processes. Keppler et al. [2001] present an abiotic mechanism for production of halocarbons during the oxidation of Fe<sup>3+</sup> in the presence of organic matter. We do not have the information available to determine if this process occurs at our site.

[18] Fungi could also be responsible for the emission of CH<sub>3</sub>Br from this soil. There was visual evidence throughout the sampling period of fungal mycelium, fruiting bodies and ectomycorrhizal. We predicted production of CH<sub>3</sub>Br by ectomycorrhizal fungi (EF) in the College Woods soils based on observations by KR2003. Their observations for *Cenococcum geophilium* revealed a linear increase in CH<sub>3</sub>Br production rates with halide content in the media. We calculated a simple linear increase in production between 0.02 mM and 20 mM Br<sup>-</sup> in media for both the highest (*Laccaria laccata*) and lowest (*Hebeloma crustuliniforme*) observed production rates and determined the rates in our soil based on the Br<sup>-</sup> content of the surface soil on 5 sampling days. Fungal production by EF ( $\mu\text{g CH}_3\text{Br m}^{-2} \text{d}^{-1}$ ) was estimated using the following equation:

$$EF_{\text{CH}_3\text{Br}} = \frac{\left[ \frac{\mu\text{gCH}_3\text{Br}}{\text{g}_{\text{fungi}}\text{d}} * \text{g}_{\text{fungi}} \right]}{A_c} \quad (2)$$

The mass of fungi (g<sub>fungi</sub>) in the collar was estimated as the fungal biomass in  $\text{g}_{\text{fungi}} \text{kg}_{\text{dry soil}}^{-1}$  multiplied by the grams of dry soil in the collar. Fungal biomass  $50.7 \pm 18.4 \mu\text{g}_{\text{fungi}}$

$\text{g}_{\text{org. matter}}^{-1}$  was an average of the fungal biomass measured in a northern hardwood forest stand of similar age and species composition [Taylor et al., 1999]. Conservatively, we believe half of the total fungal biomass to be EF. Fungal biomass was then calculated as  $0.34 \pm 0.25 \text{ g fungi in the collar area}$ . Organic matter content in the collar was measured as 60.3%. A<sub>c</sub>, collar area, is 0.397 m<sup>2</sup>.

[19] The CH<sub>3</sub>Br produced by wood-rotting fungi was estimated using equation (3) modified from L-TH2000:

$$WF_{\text{CH}_3\text{Br}} = D * [\text{Br}^-] * 10^{12} * k_{\text{fc}} * \left( \frac{m_{\text{CH}_3\text{Br}}}{m_{\text{Br}^-}} \right) \quad (3)$$

D is the annual pre-agricultural decomposition rate ( $\text{kg}_{\text{dry matter}} \text{m}^{-2} \text{yr}^{-1}$ ). Assuming steady state with decomposition equal to production, this value (0.448  $\text{kg m}^{-2} \text{yr}^{-1}$ ) was based on the annual litter production rates calculated from direct measurements by Matthews [1997] for cool-deciduous forests with evergreens. [Br<sup>-</sup>] is the measured bromide concentration in the high organic matter soil below the litter surface and falls within the range reported by L-TH2000 for litter. The net efficiency of fungal conversion of Br<sup>-</sup> to CH<sub>3</sub>Br, k<sub>fc</sub>, was calculated as 0.021 according to parameters for temperate regions given in L-TH2000. m<sub>CH<sub>3</sub>Br</sub> and m<sub>Br<sup>-</sup></sub> are the molar mass of CH<sub>3</sub>Br and Br<sup>-</sup>.

[20] Estimates from EF and WF indicate that they could be responsible for some of the production of CH<sub>3</sub>Br in these soils (Table 1). The error of the Gaussian fit controls the soil production error and is high due to the limited number of temperature and moisture manipulations. The Gaussian fit, a smoothed peak, may overestimate uptake rates when soil moisture is between 75 and 150% and when temperatures are between 10 and 25°C (Figures 2A and 2B). This may account for some of the differences between measured and modeled uptake (Figure 3).

[21] The range of EF production of CH<sub>3</sub>Br in Table 1 is driven by our fungal mass estimate, the Br<sup>-</sup> content of the soil and the high and low estimates reported by KR2003. EF may subsist on leaf or litter tissue which may have a higher Br<sup>-</sup> content [e.g. L-TH2000 and references therein]. We assumed that half of the total fungal biomass in the collar is EF. Total fungal biomass can vary seasonally due to varying substrate availability, soil temperature and soil moisture of the system [Myers et al., 2001]. Fungal biomass can also vary spatially on a local scale based on topography and disturbances such as tree fall [Morris and Boerner, 1999].

[22] Our calculations for the WF production of CH<sub>3</sub>Br are +3 and -0.27 times the production estimate. L-TH2000 believe this is a conservative estimate because 25 to 40% of the global woody decay is not included and their estimates



use production by one species of fungi and therefore a single ratio of Cl/Br emission. KR2003 have measured differing ratios of halide ion production from one species to the next. We feel that our estimate is conservative and could result in a larger range of emission if the above issues were addressed.

#### 4.4. Global Extrapolation

[23] The net consumption rate of  $168 \pm 72 \mu\text{g m}^{-2}$  for the 1999 growing season extrapolated over a global area of  $12.9 \times 10^{12} \text{ m}^2$  for temperate forests [Matthews, 1983] yields an estimate of net uptake of  $2.2 \pm 0.9 \text{ Gg of CH}_3\text{Br yr}^{-1}$ . This estimate is an order of magnitude less than the Shorter *et al.* [1995] and the Serça *et al.* [1998] estimates for temperate forest soil uptake of  $\text{CH}_3\text{Br}$ . Differences in measurement technique, sampling site characteristics or a production mechanism in the soil could all be responsible for the discrepancy between these estimates. Consumption rates change with temperature and moisture and therefore an estimate should take into account seasonal changes in consumption rate. The discrepancy between the estimates may reflect an abiotic or fungal production mechanism in the soil. The two estimates for fungal production from temperate forests: 0.5 to 5.2 Gg  $\text{CH}_3\text{Br yr}^{-1}$  from WF by L-TH2000 and 7 to 65 Gg  $\text{yr}^{-1}$  from EF by KR2003 could account for the difference between the estimates.

#### 5. Conclusions

[24] Soils have a tremendous potential to consume  $\text{CH}_3\text{Br}$  and are currently identified as significant sinks in the tropospheric budget. Production of  $\text{CH}_3\text{Br}$  occurs in soils as well and can exceed consumption resulting in a net efflux of  $\text{CH}_3\text{Br}$  to the atmosphere. An abiotic mechanism during organic matter degradation and/or fungi associated with litter and/or tree roots may be responsible for this production. Both the consumption and production processes are important to our understanding of the natural cycling of  $\text{CH}_3\text{Br}$  and the net  $\text{CH}_3\text{Br}$  exchange with these systems.

[25] **Acknowledgments.** The authors would like to acknowledge Andrew Mosedale, Sarah Pfafflin, Richelle Shaffer, Michael Keller, and Claire McSweeney for their contributions. This project was funded by a National Science Foundation Grant (EAR-9630694).

#### References

- Attieh, J. M., et al., Purification and characterization of a novel methyltransferase responsible for biosynthesis of halomethanes and methanethiol in *Brassica oleracea*, *J. Biol. Chem.*, 270, 9250–9257, 1995.
- Connell Hancock, T. L., et al., Strain IMB-1, a Novel Bacterium for the Removal of Methyl Bromide in Fumigated Agricultural Soils, *Appl. Environ. Microbiol.*, 64(8), 2899–2905, 1998.
- Crill, P. M., Seasonal patterns of methane uptake and carbon dioxide release by a temperature woodland soil, *Global Biogeochem. Cycles*, 5, 319–334, 1991.
- Dimmer, C. H., et al., Biogenic fluxes of halomethanes from Irish peatland ecosystems, *Atmos. Env.*, 35, 321–330, 2001.
- Galbally, I. E., and C. Johansson, A model relating laboratory measurements of rates of nitric oxide production and field measurements of nitric oxide emission from soils, *J. Geophys. Res.*, 94(D5), 6473–6480, 1989.
- Gan, J. S. R., et al., Production of methyl bromide by terrestrial higher plants, *Geophys. Res. Lett.*, 25, 3595–3598, 1998.
- Goodwin, K. D., et al., Consumption of tropospheric levels of methyl bromide by Cl compound-utilizing bacteria and comparison to saturation kinetics, *Appl. Environ. Microbiol.*, 67, 5437–5443, 2001.
- Harper, D. B., Halomethane from halide ion—a highly efficient fungal conversion of environmental significance, *Nature*, 315, 55–57, 1985.
- Hines, M. E., et al., Rapid consumption of low concentrations of methyl bromide by soil bacteria, *Appl. Environ. Microbiol.*, 64, 1864–1870, 1998.
- Jeffers, P. M., N. L. Wolfe, and V. Nzengung, Green plants: A terrestrial sink for atmospheric  $\text{CH}_3\text{Br}$ , *Geophys. Res. Lett.*, 25, 43–46, 1998.
- Keppeler, F., et al., Halocarbons produced by natural oxidation processes during degradation of organic matter, *Nature*, 403, 298–301, 2000.
- Kerwin, R. A., et al., Determination of atmospheric methyl bromide by cryotrapping/gas chromatography and application to soil kinetic studies using a dynamic dilution system, *Anal. Chem.*, 68, 899–903, 1996.
- Lee-Taylor, J. M., and E. A. Holland, Litter decomposition as a potential natural source of methyl bromide, *J. Geophys. Res.*, 105, 8857–8864, 2000.
- Matthews, E., Global vegetation and land use: new high-resolution data bases for climate studies, *J. Clim. Appl. Meteorol.*, 22, 474–487, 1983.
- Matthews, E., Global litter production, pools, and turnover times: Estimates from measurement data and regression models, *J. Geophys. Res.*, 102, 18,771–18,800, 1997.
- Miller, L. G., et al., Bacterial oxidation of methyl bromide in fumigated agricultural soils, *Appl. Environ. Microbiol.*, 63(11), 4346–4354, 1997.
- Morris, S. J., and R. E. J. Boerner, Spatial distribution of fungal and bacterial biomass in southern hardwood forest soils: scale dependency and landscape patterns, *Soil Biol. Biochem.*, 31, 887–902, 1999.
- Myers, R. T., et al., Landscape-level patterns of microbial community composition and substrate use in upland forest ecosystems, *Soil Sci. Soc. Am. J.*, 65, 359–367, 2001.
- Redeker, K. R., et al., Emissions of methyl halides and methane from rice paddies, *Science*, 290, 966–968, 2000.
- Rhew, R. C., et al., Natural methyl bromide and methyl chloride emissions from coastal salt marshes, *Science*, 403, 292–295, 2000.
- Rhew, R. C., et al., Shrubland fluxes of methyl bromide and methyl chloride, *J. Geophys. Res.*, 106, 20,875–20,882, 2001.
- Saini, H. S., J. M. Attieh, and A. D. Hanson, Biosynthesis of halomethanes and methanethiol by higher plants via a novel methyltransferase reaction, *Plant Cell Environ.*, 18, 1027–1033, 1995.
- Serça, D., et al., Methyl bromide deposition to soils, *Atmos. Environ.*, 32, 1581–1586, 1998.
- Shorter, J. H., et al., Rapid degradation of atmospheric methyl bromide in soils, *Nature*, 377, 717–719, 1995.
- Taylor, L. A., et al., Forest floor microbial biomass across a northern hardwood successional sequence, *Soil Biol. Biochem.*, 31, 431–439, 1999.
- UNEP, Report of the Ninth Meeting of the Parties to the Montreal Protocol on Substances that Deplete the Ozone Layer, United Nations Environment Programme UNEP, Montreal, Canada, 1997.
- van Dijk, S. M., et al., Biogenic NO emissions from forest and pasture soils: Relating laboratory studies to field measurements, *J. Geophys. Res.*, 107(D20), 8058, doi:10.1029/2001JD000358, 2002.
- Varnier, R. K., et al., An estimate of the uptake of atmospheric methyl bromide by agricultural soils, *Geophys. Res. Lett.*, 26, 727–730, 1999a.
- Varnier, R. K., P. M. Crill, and R. W. Talbot, Wetlands: a potentially significant source of atmospheric methyl bromide and methyl chloride, *Geophys. Res. Lett.*, 26, 2433–2436, 1999b.
- Yvon-Lewis, S. A., Methyl bromide in the atmosphere and ocean, *IGAC-Activities Newsletter*, 19, 9–12, 2000.

P. M. Crill, C. H. Mosedale, R. K. Varnier, and M. L. White, Complex Systems Research Center, Institute for the Study of Earth, Oceans and Space, University of New Hampshire, Durham, NH 03824, USA. (ruth.varnier@unh.edu)