

4-25-2013

# Controls on dissolved organic carbon quantity and chemical character in temperate rivers of North America

Kevin W. Hanley

*University of New Hampshire, Durham*

Wilfred M. Wollheim

*University of New Hampshire, Durham, wil.wollheim@unh.edu*

Joseph Salisbury

*University of New Hampshire, Durham, Joe.Salisbury@unh.edu*

Thomas Huntington

*United States Geological Survey*

George Aiken

*United States Geological Survey*

Follow this and additional works at: [https://scholars.unh.edu/faculty\\_pubs](https://scholars.unh.edu/faculty_pubs)

---

## Recommended Citation

\* Hanley, K. W., W.M. Wollheim, J. Salisbury, T. Huntington, G. Aiken. 2013. Controls on dissolved organic carbon quantity and chemical character in temperate rivers of North America. *Global Biogeochemical Cycles*. DOI: 10.1002/gbc.20044

This Article is brought to you for free and open access by University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact [nicole.hentz@unh.edu](mailto:nicole.hentz@unh.edu).

## Controls on dissolved organic carbon quantity and chemical character in temperate rivers of North America

Kevin W. Hanley,<sup>1</sup> Wilfred M. Wollheim,<sup>1,2</sup> Joseph Salisbury,<sup>3</sup> Thomas Huntington,<sup>4</sup> and George Aiken<sup>5</sup>

Received 16 October 2012; revised 20 March 2013; accepted 8 April 2013; published 23 May 2013.

[1] Understanding the processes controlling the transfer and chemical composition of dissolved organic carbon (DOC) in freshwater systems is crucial to understanding the carbon cycle and the effects of DOC on water quality. Previous studies have identified watershed-scale controls on bulk DOC flux and concentration among small basins but fewer studies have explored controls among large basins or simultaneously considered the chemical composition of DOC. Because the chemical character of DOC drives riverine biogeochemical processes such as metabolism and photodegradation, accounting for chemical character in watershed-scale studies will improve the way bulk DOC variability in rivers is interpreted. We analyzed DOC quantity and chemical character near the mouths of 17 large North American rivers, primarily between 2008 and 2010, and identified watershed characteristics that controlled variability. We quantified DOC chemical character using both specific ultraviolet absorbance at 254 nm (SUVA<sub>254</sub>) and XAD-resin fractionation. Mean DOC concentration ranged from 2.1 to 47 mg C L<sup>-1</sup> and mean SUVA<sub>254</sub> ranged from 1.3 to 4.7 L mg C<sup>-1</sup> m<sup>-1</sup>. We found a significant positive correlation between basin wetland cover and both bulk DOC concentration ( $R^2 = 0.78$ ;  $p < 0.0001$ ) and SUVA<sub>254</sub> ( $R^2 = 0.91$ ;  $p < 0.0001$ ), while other land use characteristics were not correlated. The strong wetland relationship with bulk DOC concentration is similar to that found by others in small headwater catchments. However, two watersheds with extremely long surface water residence times, the Colorado and St. Lawrence, diverged from this wetland relationship. These results suggest that the role of riverine processes in altering the terrestrial DOC signal at the annual scale was minimal except in river systems with long surface water residence times. However, synoptic DOC sampling of both quantity and character throughout river networks will be needed to more rigorously test this finding. The inclusion of DOC chemical character will be vital to achieving a more complete understanding of bulk DOC dynamics in large river systems.

**Citation:** Hanley, K. W., W. M. Wollheim, J. Salisbury, T. Huntington, and G. Aiken (2013), Controls on dissolved organic carbon quantity and chemical character in temperate rivers of North America, *Global Biogeochem. Cycles*, 27, 492–504, doi:10.1002/gbc.20044.

### 1. Introduction

[2] Dissolved organic carbon (DOC) quantity and chemical character in rivers and streams play key biogeochemical roles influencing drinking water quality, heavy metal

transport, stream ecosystem processes, coastal eutrophication, and the global carbon cycle [Aiken *et al.*, 2011; Buffam *et al.*, 2001; Cole *et al.*, 2007; Frey and Smith, 2005; Gattuso *et al.*, 1998; Lehtoranta *et al.*, 2009; Sholkovitz, 1976; Singer, 1999]. Large rivers are particularly important because they are a major source of material to the coastal ocean and they indicate dynamics across broad regions. However, most previous basin-scale riverine organic carbon studies have focused either on small or individual watersheds, with many finding that bulk DOC variability is related to basin-scale characteristics such as wetland cover and runoff [Buffam *et al.*, 2007; Chorover and Amistadi, 2001; Clair and Ehrman, 1996; Creed *et al.*, 2003; Dalzell *et al.*, 2007; Gergel *et al.*, 1999; Mulholland and Kuenzler, 1979; Raymond and Hopkinson, 2003]. We sought to address whether the processes that appear to control DOC quantity in small basins also scale to large and continental-scale systems. In addition, current global land-to-ocean carbon flux

<sup>1</sup>Water Systems Analysis Group, University of New Hampshire, Durham, New Hampshire, USA.

<sup>2</sup>Department of Natural Resources and Environment, University of New Hampshire, Durham, New Hampshire, USA.

<sup>3</sup>Ocean Process Analysis Laboratory, University of New Hampshire, Durham, New Hampshire, USA.

<sup>4</sup>United States Geological Survey, Augusta, Maine, USA.

<sup>5</sup>United States Geological Survey, Boulder, Colorado, USA.

Corresponding author: K. W. Hanley, Water Systems Analysis Group, University of New Hampshire, Morse Hall, 8 College Rd., Durham, NH 03824, USA. (kwz22@wildcats.unh.edu)

©2013. The Authors.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

0886-6236/13/10.1002/gbc.20044

models continue to rely on DOC concentration data of questionable quality, often collected more than 30 years ago [Alexander et al., 1998; Harrison et al., 2005; Lauerwald et al., 2012; Meybeck and Ragu, 1996; Seitzinger et al., 2005]. Here we present updated estimates of DOC concentration and flux from 17 large rivers in North America that may be used in future modeling efforts.

[3] Interpreting variability in the quantity of DOC and predicting its impact in natural systems are difficult without also taking into account its chemical character. The chemical character of DOC is largely determined by its source material and past biogeochemical transformations [Schlesinger, 1997] and as a result, DOC is made up of thousands of different molecules with a broad spectrum of characteristics and reactivity [Maurice et al., 2002]. Knowledge of DOC chemical character can improve understanding of both the sources and fate of DOC in river systems. In addition, the chemical makeup of DOC in aquatic systems influences the transport and bioavailability of heavy metals [Dittman et al., 2010] and anthropogenic organic compounds [Neale et al., 2011]. The chemical character of DOC also determines how it interacts with natural and engineered nanoparticles [Aiken et al., 2011] and impacts the production of harmful byproducts of chlorine disinfection during drinking water sanitization [Singer, 1999]. Therefore, a more complete understanding of DOC chemical character in rivers and streams will aid in our interpretation of bulk DOC variability and its influences on water quality.

[4] Some previous large and continental-scale studies have explored controls on DOC quantity and chemical character, though each had limitations. Aitkenhead and McDowell [2000] found a strong link between soil C:N and DOC flux at the annual scale among biomes ( $R^2=0.992$ ;  $p < 0.0001$ ). Despite the strength of this relationship, when it is applied to predicting DOC flux from individual watersheds, particularly large ones, its utility is limited by the necessity of geospatially extensive soil C:N data. Frost et al. [2006] characterized DOC concentration and chemical character throughout a single large river network. They found that concentration was related to a range of landscape variables including percent wetland cover and the total drainage area of individual sub-catchments. They also found that the molecular weight of DOC and its aromaticity were related to the percent lake cover and percent wetland cover of individual sub-catchments. Shih et al. [2010] developed a continental-scale total organic carbon flux model based on a variety of watershed parameters using the SPARROW modeling framework [Alexander et al., 2000]. They found that in-stream processes were significant in controlling both the quantity and inferred sources of DOC. However, implicit in their model was the assumption that all organic carbon in a reach, irrespective of chemical character, was remineralized at the same rate. This type of model simplification may be adequate to predict bulk organic carbon quantity, but it does not reflect important complexities in the underlying biogeochemical processes and makes the interpretation of model predictions problematic. By not taking into account the spectrum of organic matter chemical character among different sources, the authors likely overestimated the contribution to basin exports by more easily remineralized autochthonous sources and underestimated the more refractory allochthonous sources

[Benner, 2003; del Giorgio and Davis, 2003; del Giorgio and Pace, 2008]. Unfortunately, few comparative large-basin DOC studies have been conducted that also incorporate chemical character. Here we update large-river DOC flux estimates and improve understanding of the processes underlying DOC variability in freshwater systems by examining DOC quantity together with chemical character among 17 large rivers throughout temperate North America. We address two primary research questions:

[5] 1. Are the biogeochemical processes underlying the observed relationships between watershed-scale characteristics and DOC quantity among small rivers also important among large and continental-scale systems?

[6] 2. Can watershed-scale characteristics explain the variability of DOC chemical character among large river systems?

[7] Answers to these questions will also help to clarify the role of in-stream processes accumulated at network scales in altering the quantity and chemical character of DOC transferred from the continents to the oceans.

## 2. Methods

### 2.1. Study Sites


[8] Our study sites included 17 large watersheds from across a wide range of biomes in North America (Figure 1 and Table 1), 11 of which are monitored by the U.S. Geological Survey's National Stream Quality Accounting Network (NASQAN; <http://water.usgs.gov/nasqan/>). Basins were selected based on two criteria: large drainage areas ( $>1000 \text{ km}^2$ ) and complete daily discharge records available for the sampling period. NASQAN locations in Alaska (Yukon River) were excluded to eliminate the confounding influence of permafrost from the analysis. The primary sampling period (2008–2010) occurred during years of average temperature and only slightly elevated precipitation [NOAA-NCDC, 2012]. Site information, including geospatial coordinates, contributing drainage area, and most discharge data, was acquired through the USGS National Water Information System (NWIS; <http://waterdata.usgs.gov/nwis/>). Discharge data for the Rio Grande were obtained from the International Boundary and Water Commission [IBWC, 2010]. In all cases, runoff was calculated as discharge divided by drainage area.

[9] For several rivers, daily discharge data were available only at a nearby USGS gauging station located on the same main stem. In these cases, discharge ( $Q$ ) was scaled by the percent difference in upstream drainage areas ( $A$ ) for the nearby and the NASQAN stations:

$$Q_{\text{NASQAN}} = Q_{\text{nearby}} (A_{\text{NASQAN}} / A_{\text{nearby}}) \quad (1)$$

[10] Very little scaling was done in any case; the Altamaha, Potomac, Androscoggin, and Mobile Rivers were each scaled by 5% or less. The Penobscot River was scaled by just under 15%, but the suitability of this particular assumption was explicitly addressed by Hodgkins [1999].

[11] We used ESRI ArcGIS 9.3 to extract land cover and mean annual temperature from the National Hydrography Dataset (NHDPlus) [USGS, 2006] and spatially joined our gauging stations with the NHDPlus topological stream network. Each gauging station's corresponding stream reach



**Figure 1.** Map showing the drainage basins and sampling locations for 17 North American rivers.

record was joined with cumulative upstream percent land covers from the National Land Cover Database (NLCD) [Homer *et al.*, 2004], including agricultural, forest, grassland, shrubland, urban, and wetland land cover types. Stream reach records were also joined with cumulative-upstream mean basin air-temperature [Daly and Taylor, 1998]. Both land cover and air-temperature records were available as a part of the NHDPlus. Unfortunately, the NHDPlus topological network did not correctly model the flow path of the St. Lawrence River, and thus land cover data were unavailable. Instead we used a St. Lawrence watershed polygon provided by NASQAN [Butman *et al.*, 2012] to extract land cover data from the Commission for Environmental Cooperation's North American Environmental Atlas [NRCan/CCRS *et al.*, 2010]. We used BasinDelineator (v2.009), a tool provided as a part of the NHDPlus, to build watershed polygons for each of the other gauging stations. These polygons were then used to extract, for each basin, the percent cover of reservoirs, small lakes, and large lakes from the Global Reservoir and Dam database [Lehner *et al.*, 2011] and the Global Lake and Wetland Database [Lehner and Doll, 2004].

## 2.2. Quantity and Chemical Character

[12] Stations were sampled approximately monthly over 2 to 4 year periods between 2002 and 2010 by the U.S. Geological Survey, with most samples taken between 2008 and 2010 (Table 2). Samples were either filtered in the field using 0.45  $\mu\text{m}$  Gelman capsule filters or shipped on ice to the U.S. Geological Survey in Boulder, Colorado, within 24 h of collection and immediately filtered using pre-combusted Whatman GF/F filters of 0.7  $\mu\text{m}$  nominal pore size. DOC concentration was measured utilizing the platinum catalyzed persulfate wet oxidation method on an O.I. Analytical Model 700 TOC Analyzer<sup>TM</sup> using established methods [Aiken *et al.*, 1992]. Reported values are the averages of duplicate analyses. Standard deviation for the DOC measurement was determined to be  $\pm 0.2 \text{ mg C L}^{-1}$ . We also

measured DOC chemical character in terms of specific ultraviolet absorbance ( $\text{SUVA}_{254}$ ), which is defined as a sample's spectral absorbance at 254 nm (UVA) normalized to its DOC concentration. All samples were analyzed for UVA using a Hewlett-Packard photo-diode array spectrophotometer, and  $\text{SUVA}_{254}$  was calculated by dividing UVA by DOC concentration. We chose  $\text{SUVA}_{254}$  as the primary measure of chemical character because it is a good indicator of DOC aromaticity [Weishaar *et al.*, 2003].

[13] The proportion of bulk DOC as hydrophobic organic acids (HPOA) was determined using XAD-resin fractionation analysis following Aiken *et al.* [1992]. In brief, samples were acidified to pH 2 using HCl and passed through a column of XAD-8 resin. The HPOA fraction was retained on the XAD-8 resin and then back eluted with 0.1 M NaOH. The concentration of HPOA was determined by direct measurement of the eluent and is presented here as a fraction of bulk DOC. XAD fractionation is useful because it allows us to directly identify the hydrophobic and generally more aromatic and allochthonous compounds in the bulk DOC pool such as fulvic and humic acids [Aiken *et al.*, 1979; Aiken *et al.*, 1992].

[14] For each station daily values and the flow-weighted overall-mean for the entire sampling period (henceforth simply referred to as "mean") for DOC concentration and  $\text{SUVA}_{254}$  were estimated using LoadRunner, a graphical front-end to the USGS application LOADEST [Booth *et al.*, 2007; Runkel *et al.*, 2004]. LOADEST incorporates daily discharge, seasonality, and measured constituent data to parameterize a multiple-regression model that allows a continuous time series to be estimated from discrete measurements. Root mean square error in  $\text{mg C L}^{-1}$  (RMSE) for each basin was calculated as

$$\text{RMSE} = \sqrt{\frac{\sum ([\text{DOC}]_{\text{modeled}} - [\text{DOC}]_{\text{measured}})^2}{n}} \quad (2)$$

where [DOC] is DOC concentration in  $\text{mg C L}^{-1}$  and  $n$  is the number of observations. Annual discharge-weighted

**Table 1.** Table Showing Station Information, Upstream Drainage Area, Mean Runoff, Mean Temperature, and LOADEST Results for Annual Mean Dissolved Organic Carbon (DOC), Specific Ultraviolet Absorption at 254 nm (SUVA<sub>254</sub>), and Hydrophobic Organic Acids (HPOA)<sup>a</sup>

| Site Number | River Name                      | Location             | Lat   | Long    | Drainage (km <sup>2</sup> ) | Runoff (cm/yr) | Temperature (°C) | DOC Yield (g C yr <sup>-1</sup> m <sup>-2</sup> ) | DOC Load (kg C d <sup>-1</sup> ) | Mean [DOC] (mg C L <sup>-1</sup> ) | Mean SUVA <sub>254</sub> (L mg C <sup>-1</sup> m <sup>-1</sup> ) | Mean HPOA (Fraction [DOC]) | HPOA Load (kg C d <sup>-1</sup> ) |
|-------------|---------------------------------|----------------------|-------|---------|-----------------------------|----------------|------------------|---|----------------------------------|------------------------------------|--|----------------------------|-----------------------------------|
| 01036390    | Penobscot River                 | Eddington, ME        | 44.83 | -68.70  | 19460                       | 80.66          | 4.4              | 7.55  | 403000                           | 9.3                                | 3.8  | ND                         | ND                                |
| 01049265    | Kennebec River                  | North Sidney, ME     | 44.47 | -69.68  | 13990                       | 78.12          | 4.3              | 4.97  | 191000                           | 6.4                                | 3.6  | ND                         | ND                                |
| 01059400    | Androscoggin River              | Brunswick, ME        | 43.92 | -69.97  | 8894                        | 88.00          | 4.9              | 5.31  | 129000                           | 6.0                                | 3.6  | ND                         | ND                                |
| 01578310    | Susquehanna River <sup>b</sup>  | Conowingo, MD        | 39.66 | -76.17  | 70200                       | 52.41          | 8.7              | 1.38  | 266000                           | 2.7                                | 2.3  | 0.39                       | 90700                             |
| 01646580    | Potomac River <sup>b</sup>      | Washington, D.C.     | 38.93 | -77.12  | 29970                       | 39.66          | 11.0             | 1.64  | 134000                           | 4.3                                | 2.6  | 0.36                       | 45000                             |
| 02175000    | Edisto River                    | Givhans, SC          | 33.03 | -80.39  | 7071                        | 16.86          | 17.6             | 1.96  | 37900                            | 11.2                               | 4.0  | 0.66                       | 24000                             |
| 02226160    | Altamaha River <sup>b</sup>     | Everett City, GA     | 31.43 | -81.61  | 36000                       | 28.76          | 17.5             | 2.99  | 297000                           | 10.1                               | 4.2  | 0.44                       | 128000                            |
| 02231000    | St. Mary's River                | MacClenny, FL        | 30.36 | -82.08  | 1800                        | 39.01          | 19.7             | 18.52   | 92000                            | 46.8                               | 4.7  | 0.71                       | 65600                             |
| 02322500    | Santa Fe River                  | Fort White, FL       | 29.85 | -82.72  | 2634                        | 34.88          | 20.1             | 5.54  | 40000                            | 12.9                               | 4.0  | 0.66                       | 29300                             |
| 02470500    | Mobile River <sup>b</sup>       | Mount Vernon, AL     | 31.09 | -87.98  | 111030                      | 25.17          | 16.5             | 1.36  | 414000                           | 5.7                                | 3.4  | 0.52                       | 209000                            |
| 04264331    | St. Lawrence River <sup>b</sup> | Cornwall, ON         | 45.01 | -74.79  | 773900                      | 30.58          | ND               | 1.24  | 1800000                          | 2.8                                | 1.3  | 0.29                       | 529000                            |
| 07374525    | Mississippi River <sup>b</sup>  | Belle Chasse, LA     | 29.86 | -89.98  | 2930000                     | 16.44          | 9.8              | 0.66  | 5260000                          | 4.0                                | 3.0  | 0.43                       | 2050000                           |
| 08475000    | Rio Grande <sup>b</sup>         | Brownsville, TX      | 25.88 | -97.45  | 456700                      | 0.11           | 18.0             | 0.01  | 9200                             | 5.9                                | 2.1  | 0.35                       | 3180                              |
| 09522000    | Colorado River <sup>b</sup>     | Morelos Dam, AZ      | 32.72 | -114.72 | 639000                      | 0.30           | 10.3             | 0.01  | 16700                            | 3.1                                | 1.7  | 0.37                       | 5490                              |
| 11303500    | San Joaquin River <sup>b</sup>  | Vernalis, CA         | 37.68 | -121.27 | 35058                       | 5.16           | 12.0             | 0.19  | 18400                            | 3.6                                | 2.5  | 0.44                       | 5160                              |
| 11447650    | Sacramento River <sup>b</sup>   | Freepoint Bridge, CA | 38.46 | -121.50 | 69457                       | 21.11          | 12.0             | 0.63  | 119000                           | 2.9                                | 2.7  | 0.39                       | 33400                             |
| 14246900    | Columbia River <sup>b</sup>     | Quincy, OR           | 46.18 | -123.18 | 665400                      | 30.59          | 6.5              | 0.65  | 1180000                          | 2.1                                | 2.7  | 0.42                       | 461000                            |

<sup>a</sup>Rivers are sorted clockwise around the North American coastline starting in the Northeastern United States. *ND* indicates no data available.

<sup>b</sup>Denotes NASQAN monitoring.

**Table 2.** Table Showing LOADEST Model Information<sup>a</sup>

| River Name         | <i>n</i> | Year Begin | Year End | <i>R</i> <sup>2</sup> | DOC<br>RMSE<br>(mg C L <sup>-1</sup> ) | NS   | <i>R</i> <sup>2</sup> | SUVA<br>RMSE<br>(L mg C <sup>-1</sup> m <sup>-1</sup> ) | NS   | <i>R</i> <sup>2</sup> | HPOA<br>RMSE<br>(%[DOC]) | NS        |
|--------------------|----------|------------|----------|-----------------------|--|------|-----------------------|---|------|-----------------------|--------------------------|-----------|
| Penobscot River    | 61       | 2004       | 2008     | 0.97                  | 1.8                                    | 0.60 | 0.99                  | 0.2   | 0.39 | <i>ND</i>             | <i>ND</i>                | <i>ND</i> |
| Kennebec River     | 12       | 2006       | 2007     | 0.99                  | 0.5                                    | 0.75 | 0.99                  | 0.1   | 0.43 | <i>ND</i>             | <i>ND</i>                | <i>ND</i> |
| Androscoggin River | 12       | 2006       | 2007     | 0.99                  | 0.4                                    | 0.81 | 0.99                  | 0.1   | 0.58 | <i>ND</i>             | <i>ND</i>                | <i>ND</i> |
| Susquehanna River  | 22       | 2008       | 2010     | 0.94                  | 0.4                                    | 0.39 | 0.96                  | 0.3   | 0.09 | 0.90                  | 0.04                     | 0.18      |
| Potomac River      | 21       | 2008       | 2010     | 0.99                  | 0.3                                    | 0.80 | 0.99                  | 0.1   | 0.81 | 0.99                  | 0.03                     | 0.92      |
| Edisto River       | 18       | 2005       | 2008     | 0.96                  | 1.6                                    | 0.81 | 0.99                  | 0.2   | 0.28 | 0.96                  | 0.04                     | 0.81      |
| Altamaha River     | 19       | 2008       | 2009     | 0.99                  | 1.1                                    | 0.67 | 0.99                  | 0.3   | 0.66 | 0.90                  | 0.11                     | -0.51     |
| St. Mary's River   | 31       | 2002       | 2006     | 0.99                  | 8.5                                    | 0.68 | 0.99                  | 0.3   | 0.17 | 0.98                  | 0.04                     | 0.67      |
| Santa Fe River     | 29       | 2002       | 2004     | 0.93                  | 5.3                                    | 0.78 | 0.98                  | 0.4   | 0.79 | 0.93                  | 0.07                     | 0.62      |
| Mobile River       | 25       | 2008       | 2010     | 0.98                  | 0.6                                    | 0.71 | 0.99                  | 0.2   | 0.72 | 0.98                  | 0.04                     | 0.36      |
| St. Lawrence River | 16       | 2008       | 2009     | 0.93                  | 0.1                                    | 0.80 | 0.74                  | 0.1   | 0.53 | 0.68                  | 0.02                     | 0.36      |
| Mississippi River  | 23       | 2008       | 2010     | 0.94                  | 0.3                                    | 0.50 | 0.99                  | 0.1   | 0.71 | 0.98                  | 0.03                     | 0.80      |
| Rio Grande         | 21       | 2008       | 2009     | 0.99                  | 0.3                                    | 0.44 | 0.98                  | 0.2   | 0.32 | 0.98                  | 0.02                     | 0.30      |
| Colorado River     | 27       | 2008       | 2010     | 0.92                  | 0.4                                    | 0.17 | 0.96                  | 0.1   | 0.54 | 0.99                  | 0.03                     | 0.37      |
| San Joaquin River  | 23       | 2008       | 2010     | 0.90                  | 0.8                                    | 0.48 | 0.98                  | 0.2   | 0.27 | 0.81                  | 0.06                     | 0.21      |
| Sacramento River   | 24       | 2008       | 2010     | 0.95                  | 0.4                                    | 0.80 | 0.97                  | 0.2   | 0.82 | 0.97                  | 0.05                     | 0.95      |
| Columbia River     | 18       | 2009       | 2010     | 0.97                  | 0.2                                    | 0.66 | 0.96                  | 0.3   | 0.61 | 0.94                  | 0.03                     | 0.60      |

<sup>a</sup>*R*<sup>2</sup>, Nash-Sutcliffe coefficient, and root mean square error are shown for dissolved organic carbon (DOC), specific ultraviolet absorption at 254 nm (SUVA<sub>254</sub>) and hydrophobic organic acids (HPOA). *ND* indicates that no data were available.

concentration means were automatically calculated from the daily modeled values, and fluxes were simply the sum of daily concentrations multiplied by daily discharge over the time period of interest. Mean concentration, flux, and yield were calculated by taking the average of the annual means for each basin. In order to compare among watersheds, we divided flux by basin area to obtain DOC yield.

### 3. Results

#### 3.1. Basin Attributes and Mean DOC Characteristics

[15] Basins ranged in drainage area from 1800 km<sup>2</sup> for the St. Mary's River in Florida to 2,930,000 km<sup>2</sup> for the Mississippi River in Louisiana (Table 1). Compared to the Mississippi, the St. Mary's River basin is quite small, but it is still much larger than most basins found in comparative riverine DOC studies. In total, the watersheds for all 17 rivers accounted for more than 70% of the land area of the contiguous United States and 26% of the land area of North

America. The most northerly river was the Columbia in Oregon with a mean watershed latitude of 46.2°N, though the coldest watershed belonged to the Kennebec River in Maine, with a mean annual temperature of 4.3°C. The most southerly and the warmest was the Santa Fe River in Florida, with a mean watershed latitude of 29.9°N and a mean annual temperature of 20.1°C. Mean runoff during each basin's sampling period ranged from 0.11 cm/yr for the Rio Grande in Texas to 88.00 cm/yr for the Androscoggin River in Maine (Table 1).


[16] Wetland cover ranged from 0.1% for the Rio Grande in Texas to 41.6% for the St. Mary's River (Table 3). Agricultural land cover ranged from 1.5% for the Penobscot River in Maine to 42.3% for the Mississippi River. The Rio Grande had the greatest grassland cover, at 34.2%, while several basins possessed 0.0% grassland cover, including nearly all eastern basins. The Rio Grande was the least forested, at 11.3% cover, and was tied with the Colorado River in Arizona for the least urban, at 0.5%. The most forested

**Table 3.** Table Showing Various Land Cover Attributes in Percent Basin Cover<sup>a</sup>

| River Name         | Agricultural (ag) | Forest | Grassland (grass) | Shrubland (shrub) | Urban | Wetland (wet) | Reservoir | Lake |
|--------------------|-------------------|--------|-------------------|-------------------|-------|---------------|-----------|------|
| Penobscot River    | 1.5               | 89.3   | 0.0               | 0.3               | 0.6   | 5.5           | 2.3       | 5.3  |
| Kennebec River     | 6.0               | 85.0   | 0.0               | 0.4               | 1.2   | 3.9           | 3.8       | 5.9  |
| Androscoggin River | 5.6               | 87.7   | 0.0               | 0.2               | 1.7   | 3.5           | 2.3       | 4.1  |
| Susquehanna River  | 28.7              | 67.4   | 0.0               | 0.0               | 2.6   | 0.5           | 0.1       | 0.7  |
| Potomac River      | 34.7              | 61.2   | 0.0               | 0.0               | 2.8   | 0.5           | 0.0       | 0.3  |
| Edisto River       | 32.3              | 45.4   | 0.0               | 0.0               | 1.6   | 15.3          | 0.0       | 0.0  |
| Altamaha River     | 24.0              | 58.4   | 0.0               | 0.0               | 3.5   | 8.3           | 0.4       | 0.6  |
| St. Mary's River   | 2.0               | 46.1   | 1.2               | 0.2               | 1.5   | 41.6          | 0.0       | 0.2  |
| Santa Fe River     | 16.9              | 46.4   | 11.2              | 1.5               | 2.4   | 16.0          | 0.0       | 1.5  |
| Mobile River       | 18.5              | 70.8   | 0.0               | 0.0               | 1.8   | 6.4           | 0.5       | 0.9  |
| St. Lawrence River | 18.2              | 37.7   | 1.9               | 0.4               | 3.1   | 3.9           | 3.5       | 34.3 |
| Mississippi River  | 42.3              | 23.0   | 25.2              | 4.7               | 1.7   | 2.5           | 0.5       | 0.9  |
| Rio Grande         | 1.8               | 11.3   | 34.2              | 50.1              | 0.5   | 0.1           | 0.2       | 0.1  |
| Colorado River     | 2.1               | 23.7   | 12.3              | 57.1              | 0.5   | 0.2           | 0.3       | 0.2  |
| San Joaquin River  | 16.9              | 36.3   | 15.3              | 16.2              | 2.2   | 0.8           | 0.6       | 0.6  |
| Sacramento River   | 9.9               | 54.6   | 15.2              | 14.5              | 1.7   | 0.7           | 0.9       | 0.7  |
| Columbia River     | 12.9              | 42.3   | 10.4              | 29.7              | 0.8   | 0.4           | 0.7       | 0.6  |

<sup>a</sup>Corresponding variables in LOADEST models (Table 4) are shown in parenthesis.





**Figure 2.** Mean dissolved organic carbon (DOC) concentration versus percent wetland cover (a) for all 17 sites, represented by a dashed line and (b) excluding the St. Mary's River, represented by a solid black line. The St. Mary's River and the St. Lawrence River are highlighted in green. Error bars represent root mean square error for the LOADEST model and are smaller than the size of the data point in some cases.

and the most urban were the Penobscot River at 89.3% and the Altamaha River at 3.5%, respectively. Shrubland ranged from 0.0% for the Mobile, Susquehanna, Potomac, Edisto, and Altamaha Rivers to 57.1% for the Colorado River. For reservoirs, the Potomac, Edisto, St. Mary's, and Santa Fe Rivers were least covered at 0.0% and the Kennebec River was the most covered at 3.8%. Finally, lake cover ranged from 0.0% for the Edisto River to 34.3% for the St. Lawrence River.

[17] Mean DOC concentrations from LOADEST ranged from 2.1 mg C L<sup>-1</sup> for the Columbia River in Oregon to 46.8 mg C L<sup>-1</sup> for the St. Mary's River, while DOC load ranged from 9200 kg C d<sup>-1</sup> for the Rio Grande to 5,260,000 kg C d<sup>-1</sup> for the Mississippi River. DOC yield ranged from 0.01 g C yr<sup>-1</sup> m<sup>-2</sup> for the Colorado River in Arizona and the Rio Grande, to 18.5 g C yr<sup>-1</sup> m<sup>-2</sup> for the St. Mary's River. Mean SUVA<sub>254</sub>, ranged from 1.3 L mg C<sup>-1</sup> m<sup>-1</sup> for the St. Lawrence River to 4.7 L mg C<sup>-1</sup> m<sup>-1</sup> for the St. Mary's River. Mean HPOA fraction ranged from 0.29 for the St. Lawrence River to 0.71 for the Santa Fe River in Florida, while HPOA load ranged from 3180 kg C d<sup>-1</sup> for the Rio Grande to 2,050,000 kg C d<sup>-1</sup> for the Mississippi River. Total DOC and HPOA flux for all basins studied was 3.80 Tg C yr<sup>-1</sup> and 1.34 Tg C yr<sup>-1</sup>, respectively (Table 1).

[18] All LOADEST models used to estimate mean DOC concentration, SUVA<sub>254</sub>, and HPOA concentration were significant to the level of  $\alpha=0.01$ , with  $p < 0.0001$  in all cases except the SUVA<sub>254</sub> and HPOA models for the St. Lawrence River where  $p=0.0008$  and 0.002, respectively. For DOC,  $R^2$  ranged from 0.90 to 0.99 and for SUVA<sub>254</sub>, from 0.74 to 0.99. Nash-Sutcliffe coefficients [Nash and Sutcliffe, 1970] ranged from 0.17 to 0.81 for DOC and from 0.09 to 0.82 for SUVA<sub>254</sub>. For HPOA,  $R^2$  ranged from 0.68 to 0.99 and Nash-Sutcliffe coefficients ranged from -0.51 (for the Altamaha River in Georgia) to 0.95. The positive Nash-Sutcliffe values presented here indicate that with the exception of the Altamaha HPOA model, all LOADEST models predicted measured values with more accuracy than a simple mean (Table 2).

### 3.2. DOC Quantity Patterns

[19] We found a strong positive correlation between percent wetland cover ( $L_w$ ) and mean DOC concentration in mg C L<sup>-1</sup> ( $R^2=0.93$ ,  $p < 0.0001$ ; Figure 2a and Table 3). However, the slope of this relationship was highly skewed by the wetland-dominated St. Mary's River. When we excluded the St. Mary's as an outlier, the model better fulfilled the assumptions of a linear regression and the DOC

**Table 4.** Table Showing Relationships Between Both Dissolved Organic Carbon (DOC) Concentration and Specific Ultraviolet Absorption at 254 nm (SUVA<sub>254</sub>) and a Variety of Land Cover Attributes (Table 3)<sup>a</sup>

| Predictor    | Land Cover Versus DOC          | $R^2$ | $p$ -Value | Land Cover Versus SUVA <sub>254</sub>      | $R^2$ | $p$ -Value |
|--------------|--------------------------------|-------|------------|--|-------|------------|
| Wetland      | [DOC]=0.590 × wet+3.278        | 0.78  | <0.0001    | SUVA=0.440 × log(wet)+2.884                | 0.91  | <0.0001    |
| Agricultural | [DOC]=0.004 × ag+5.733         | 0.00  | 0.954      | SUVA=-0.052 × log(ag)+3.187                | 0.00  | 0.812      |
| Forest       | [DOC]=0.033 × forest+4.072     | 0.05  | 0.387      | SUVA=0.821 × log(forest)+-0.085            | 0.23  | 0.054      |
| Grass        | [DOC]=-0.071 × grass+6.364     | 0.05  | 0.405      | SUVA=-0.109 × log(grass)+2.96              | 0.18  | 0.092      |
| Shrub        | [DOC]=-0.064 × shrub+6.504     | 0.13  | 0.177      | SUVA=-0.127 × log(shrub)+2.989             | 0.19  | 0.077      |
| Urban        | [DOC]=0.29 × urban+5.284       | 0.01  | 0.770      | SUVA=0.177 × log(urban)+2.99               | 0.01  | 0.663      |
| Reservoir    | [DOC]=-0.295 × reservoir+6.099 | 0.01  | 0.684      | SUVA=-0.163 × log(reservoir)+2.863         | 0.12  | 0.173      |
| Lake         | [DOC]=-0.074 × lake+6.066      | 0.03  | 0.493      | SUVA=-0.087 × log(lake)+3.042              | 0.03  | 0.518      |
| Predictor    | Land Cover Versus Residual-DOC | $R^2$ | $p$ -Value | Runoff versus Residual SUVA <sub>254</sub> | $R^2$ | $p$ -Value |
| Agricultural | RD=-0.096 × ag+1.551           | 0.21  | 0.062      | RS=-0.12 × log(ag)+0.28                    | 0.06  | 0.359      |
| Forest       | RD=-0.001 × forest+0.071       | 0.00  | 0.964      | RS=0.172 × log(forest)+-0.658              | 0.03  | 0.532      |
| Grass        | RD=0.061 × grass+-0.456        | 0.06  | 0.358      | RS=-0.034 × log(grass)+-0.033              | 0.05  | 0.411      |
| Shrub        | RD=0.05 × shrub+-0.517         | 0.12  | 0.175      | RS=-0.008 × log(shrub)+-0.005              | 0.00  | 0.868      |
| Urban        | RD=-0.98 × urban+1.739         | 0.11  | 0.195      | RS=-0.284 × log(urban)+0.122               | 0.09  | 0.241      |
| Reservoir    | RD=-0.039 × reservoir+0.037    | 0.00  | 0.944      | RS=-0.069 × log(reservoir)+-0.086          | 0.06  | 0.355      |
| Lake         | RD=-0.083 × lake+0.277         | 0.06  | 0.324      | RS=-0.136 × log(lake)+-0.037               | 0.19  | 0.083      |

<sup>a</sup>The second group of formulas represents the relationships between the residuals of the DOC or SUVA<sub>254</sub> versus percent wetland-cover regression (equations (3) and (5)) and the same land cover attributes. For all DOC relationships, the St. Mary's River is excluded as an outlier. For the SUVA<sub>254</sub> versus percent wetland cover relationship, the Colorado River and the St. Lawrence River are excluded as outliers.

**Table 5.** Table Showing Within-Basin Runoff (RO) Relationships for Dissolved Organic Carbon (DOC) Concentration and Specific Ultraviolet Absorption at 254 nm (SUVA<sub>254</sub>)<sup>a</sup>

| River Name         | Runoff-DOC Formula        | R <sup>2</sup> | p-Value | Runoff-SUVA <sub>254</sub> Formula | R <sup>2</sup> | p-Value |
|--------------------|---------------------------|----------------|---------|------------------------------------|----------------|---------|
| Penobscot River    | [DOC] = 0.01 × RO + 8.46  | 0.11           | 0.004   | N/S                                | N/S            | 0.119   |
| Kennebec River     | [DOC] = -0.01 × RO + 7.40 | 0.28           | 0.043   | N/S                                | N/S            | 0.275   |
| Androscoggin River | N/S                       | N/S            | 0.191   | N/S                                | N/S            | 0.915   |
| Susquehanna River  | N/S                       | N/S            | 0.963   | N/S                                | N/S            | 0.227   |
| Potomac River      | N/S                       | N/S            | 0.706   | N/S                                | N/S            | 0.058   |
| Edisto River       | [DOC] = 0.27 × RO + 4.63  | 0.40           | 0.003   | N/S                                | N/S            | 0.055   |
| Altamaha River     | [DOC] = 0.03 × RO + 8.53  | 0.18           | 0.022   | SUVA = 0.01 × RO + 3.89            | 0.27           | 0.006   |
| St. Mary's River   | N/S                       | N/S            | 0.336   | N/S                                | N/S            | 0.101   |
| Santa Fe River     | [DOC] = 0.17 × RO + 2.78  | 0.62           | < 0.001 | SUVA = 0.01 × RO + 3.76            | 0.10           | 0.045   |
| Mobile River       | N/S                       | N/S            | 0.443   | SUVA = 0.01 × RO + 3.06            | 0.35           | 0.001   |
| St. Lawrence River | [DOC] = 0.06 × RO + 0.97  | 0.38           | 0.005   | N/S                                | N/S            | 0.89    |
| Mississippi River  | N/S                       | N/S            | 0.086   | SUVA = 0.02 × RO + 2.57            | 0.27           | 0.007   |
| Rio Grande         | N/S                       | N/S            | 0.06    | SUVA = 2.05 × RO + 1.85            | 0.16           | 0.043   |
| Colorado River     | [DOC] = 1.67 × RO + 2.56  | 0.14           | 0.029   | SUVA = 0.95 × RO + 1.35            | 0.20           | 0.011   |
| San Joaquin River  | N/S                       | N/S            | 0.583   | N/S                                | N/S            | 0.107   |
| Sacramento River   | [DOC] = 0.06 × RO + 1.37  | 0.50           | < 0.001 | SUVA = 0.04 × RO + 1.67            | 0.71           | < 0.001 |
| Columbia River     | [DOC] = 0.02 × RO + 1.47  | 0.47           | 0.001   | SUVA = 0.02 × RO + 1.96            | 0.27           | 0.017   |

<sup>a</sup>Basins without a significant relationship are labeled N/S.

concentration variability was still well explained by percent wetland cover ( $R^2 = 0.78$ ,  $p < 0.0001$ ; Figure 2b and Table 4):

$$[\text{DOC}] = 0.59 \times L_W + 3.28 \quad (3)$$

[20] In this relationship, the St. Lawrence River falls well below the regression line but is still included in the analysis. We found no relationships significant to the level of  $\alpha = 0.01$  between any other land cover attribute (including land cover classes, lake cover, and reservoir cover) and DOC concentration. There were also no correlations between any land cover attribute and the residuals of the DOC concentration versus percent wetland regression (Table 4).

[21] We found no significant relationship at the level of  $\alpha = 0.01$  between mean annual runoff (RO) and mean DOC concentration among basins. However, we did find a significant positive correlation between mean DOC yield ( $\text{g C yr}^{-1} \text{m}^{-2}$ ) and runoff ( $\text{cm/yr}$ ) which strengthened when the St. Mary's River was again excluded as an outlier ( $R^2 = 0.68$ ,  $p < 0.0001$ ):

$$\text{DOC}_{\text{yield}} = 0.07 \times \text{RO} - 0.158 \quad (4)$$

[22] It should be noted that runoff is used within the LOADEST model for estimating mean annual DOC concentration. Within individual systems, around half of basins exhibited a significant positive correlation between daily runoff and discrete DOC concentration. In these basins, runoff explained between 11% and 62% of concentration variability (Table 5). However, we found no correlation between the statistical significance of these relationships and a basin's percent wetland cover or mean annual runoff.


### 3.3. DOC Chemical Character Patterns

[23] Percent wetland cover also appeared to be an important variable in controlling DOC chemical character. We found a strong positive correlation between the logarithm of percent wetland cover and mean SUVA<sub>254</sub> among the rivers in our data set ( $R^2 = 0.60$ ,  $p = 0.0001$ ). In this relationship the St. Mary's River was not found to be an outlier. However, the St. Lawrence and Colorado Rivers exhibited far lower SUVA<sub>254</sub> in  $\text{L mg C}^{-1} \text{m}^{-1}$  than expected based

on their wetland cover, and when they were excluded from the regression as outliers, the relationship between percent wetland cover and SUVA<sub>254</sub> improved ( $R^2 = 0.91$ ,  $p < 0.0001$ ; Figure 3 and Table 4):


$$\text{SUVA}_{254} = 0.441 \times \log(L_W) + 2.884 \quad (5)$$

[24] No significant relationships were found between any other land cover attribute and SUVA<sub>254</sub>, nor between any land cover attribute and the residuals of the SUVA<sub>254</sub> versus percent wetland regression (Table 4).



**Figure 3.** Mean specific ultraviolet absorption at 254 nm (SUVA<sub>254</sub>) versus percent wetland cover for all basins. X axis is logscale and excluded outliers are in green. Final model is in black. Model before the outliers were excluded is shown in dotted grey. Error bars represent root mean square error for the LOADEST model.





**Figure 4.** Mean specific ultraviolet absorption at 254 nm ( $SUVA_{254}$ ) versus mean runoff for all basins. Excluded outliers (St. Lawrence and Colorado) are shown in green.

[25] We also did not find a statistically significant relationship between mean runoff and  $SUVA_{254}$  among basins, with or without outliers (Figure 4), but within individual systems, around half of basins exhibited a significant positive correlation between discrete  $SUVA_{254}$  and daily runoff, with runoff explaining between 10% and 71% of variability (Table 5). Basins with a significant relationship tended to be in the south and the west, whereas northern and eastern rivers did not tend to show significance. As with DOC concentration, we found no correlation between the statistical significance of these relationships and a basin's percent wetland cover.

[26] HPOA as a percentage of bulk DOC was positively correlated with  $SUVA_{254}$  measurements in individual grab samples across 14 basins where both measurements were made ( $R^2 = 0.89$ ,  $p < 0.0001$ ). Within individual basins, the relationship between HPOA and  $SUVA_{254}$  was significant for all but three basins (Mobile, Mississippi, and Colorado) with  $R^2$  ranging from 0.24 for the St. Mary's River to 0.87 for the neighboring Santa Fe River. As a result, HPOA patterns were very similar to  $SUVA_{254}$ .

## 4. Discussion


### 4.1. DOC Quantity

[27] We identified a significant positive relationship between basin wetland cover and mean DOC concentration among large watersheds that was consistent with what has been reported for small basins [Buffam et al., 2007; Creed et al., 2003; Eckhardt and Moore, 1990; Gergel et al., 1999; Gorham et al., 1998; Raymond and Hopkinson, 2003]. Similar observations previously made among small basins have typically been explained by the hypothesis that runoff from a wetland to a stream channel would be less likely to have intersected the mineral soil horizon than runoff from non-wetland systems. These flow paths are

important because DOC builds up in wetlands due to anaerobic conditions, while DOC in subsurface flow intersecting the mineral horizon is more likely to be removed from solution by biotic processing and adsorption [Aitkenhead-Peterson et al., 2003; Buffam et al., 2007; Eckhardt and Moore, 1990; Tipping et al., 1999]. We found that these small-basin patterns also occurred in large river systems, demonstrating that the control exerted by wetlands on riverine DOC continued to be evident in large and continental scale systems. The lack of a significant relationship between any other land attribute and annual mean DOC concentration further highlighted the unique relationship between wetlands and riverine DOC.

[28] Mean runoff appeared to control variation in DOC yield among large basins at the annual scale (equation (4)) without directly influencing DOC concentration [Mulholland and Kuenzler, 1979; Mulholland and Watts, 1982]. The lack of a significant relationship between mean DOC concentration and mean runoff indicated that the controls exerted on mean yield by runoff were unrelated to the factors controlling mean concentration. Rather, when predicting DOC flux at annual scales from large rivers, annual runoff should be considered a vector, rather than an explanatory variable for concentration.

[29] Although percent wetland cover is a powerful explanatory variable, substantial unexplained DOC concentration variability remained among the seven least wetland-dominated watersheds. These watersheds are geographically diverse and include the San Joaquin, Rio Grande, Potomac, Colorado, Columbia, Sacramento, and Susquehanna Rivers, which all have less than 1% wetland cover and possess a mean DOC concentration of  $3.5 \pm 1.3 \text{ mg C L}^{-1}$ . Among these basins, we found that DOC concentration was positively correlated with mean annual watershed temperature ( $R^2 = 0.79$ ,  $p = 0.004$ ; Figure 5) and it was not significantly



**Figure 5.** Mean dissolved organic carbon (DOC) concentration versus mean watershed temperature for systems with less than 1% wetland cover. Error bars represent root mean square error for the LOADEST model.

correlated with any type of land cover. In addition, when temperature was included with wetland cover in a multiple regression across all 17 sites, the prediction did not improve. These results suggested the possibility that climate effects dominate concentration variability among low-wetland systems at the annual scale. Mean annual temperatures may be correlated with DOC concentration because longer growing seasons allow for greater primary production and thus greater production of soil organic material. However, it is clear that wetland controls eclipse climate effects in basins where wetlands are more extensive.

[30] In this study, we sampled across the hydrograph and estimated annual mean concentration and flux using LOADEST, which led to estimates that differed from those made previously. The mean DOC concentration for the Mississippi River reported by *Leenheer* [1982] ( $8.79 \text{ mg C L}^{-1}$ ) was more than double the mean concentration found in this study ( $4.0 \text{ mg C L}^{-1}$ ). The Mississippi is the largest river in North America so even a small percentage error in its concentration can lead to a large absolute error in the estimation of total continental land-sea flux. For example, the DOC flux estimate made by *Leenheer* [1982] for the Mississippi exceeded our estimate by  $1.6 \text{ Tg/yr}$ , nearly as much as the combined annual DOC flux from all other rivers in this study ( $1.9 \text{ Tg/yr}$ ). This estimate was later used by *Ludwig et al.* [1996] in the parameterization of a global continent-to-ocean organic carbon flux model.

[31] Some more recent studies are also partly based on values not updated in more than 30 years, which can include mean DOC concentrations based on only two to four measurements per year [*Alexander et al.*, 1998; *Harrison et al.*, 2005; *Lauerwald et al.*, 2012; *Meybeck and Ragu*, 1996; *Seitzinger et al.*, 2005]. For example, *Seitzinger et al.* [2005] and *Harrison et al.* [2005] use DOC concentration measurements at a Mississippi River gauging station recorded from 1978 to 1984, arriving at a mean concentration of  $6.7 \text{ mg C L}^{-1}$  ( $n = 14$ ). However, we examined the USGS National Water Information System for historic DOC concentration measurements from the same station and found only a single value that exceeded  $5 \text{ mg C L}^{-1}$  in the period since 1997. The particular case of the Mississippi is illustrative because the historical USGS data show a dramatic shift between the 14 samples taken between 1978 and 1984 that led to the mean of  $6.7 \text{ mg C L}^{-1}$  and the far more numerous samples taken after 1997. In the early period, we found a mean concentration of  $5.9 \text{ mg C L}^{-1}$ , with a standard deviation of  $2.6 \text{ mg C L}^{-1}$  and during the later period from 1997 to 2012, the mean was  $3.7 \text{ mg C L}^{-1}$  with a standard deviation of only  $0.5 \text{ mg C L}^{-1}$ . The higher concentration and standard deviation in the early period were driven by a few very high concentration measurements. As previously shown, even a small overestimate of mean DOC concentration for a large river has the potential to result in huge absolute errors in the estimation of flux. By sampling for multiple years across the hydrograph, employing strict QA/QC, and estimating fluxes using the LOADEST model, we present an updated picture of the quantity of DOC recently delivered by these large rivers to estuaries and the coastal ocean.

#### 4.2. DOC Chemical Character

[32] Basin-scale wetland cover was significantly correlated with mean DOC chemical character among large basins. The role of wetlands in controlling  $\text{SUVA}_{254}$  has

been previously demonstrated among small northern basins [*Agren et al.*, 2008] but this is the first time such a clear relationship has been observed in a large and continental scale study. The role of wetlands in controlling  $\text{SUVA}_{254}$  among large basins may be related to the previously discussed hypothesis that subsurface flow through mineral versus organic soil horizons can influence DOC concentration variability. Saturated, anaerobic conditions common in wetland soils can inhibit organic matter remineralization and lead to the persistence of semi-labile aromatic compounds in subsurface flow that would drive up  $\text{SUVA}_{254}$  in the rivers and streams to which it is discharged [*Guillemette and del Giorgio*, 2011]. In the absence of wetlands, extensive biotic processing and the preferential sorption of strongly UV-absorbing, aromatic DOC molecules onto mineral soils, and in some specialized cases, onto sediments and particles within the stream channel, would drive down  $\text{SUVA}_{254}$  [*Chorover and Amistadi*, 2001; *McKnight et al.*, 2002; *McKnight et al.*, 1992; *Meier et al.*, 1999; *Perez et al.*, 2011; *Tipping et al.*, 1999]. Thus, if subsurface flow paths are less likely to intersect mineral horizons in watersheds with extensive wetland cover, less of the aromatic, strongly UV-absorbing DOC would be removed and DOC with a higher  $\text{SUVA}_{254}$  would be more likely to enter river systems. The relationship between percent wetland cover and  $\text{SUVA}_{254}$  in this study is strong, indicating that the wetland processes described here are important in large rivers at the annual scale. The lack of any statistically significant relationships between other land cover attributes and  $\text{SUVA}_{254}$  further strengthens this interpretation. It is worth noting that the wetlands influencing the rivers described here range across a number of biomes and contain a wide range of vegetation types. In our analyses, we did not observe an influence of source (i.e., vegetation) differences on the  $\text{SUVA}_{254}$  versus basin-scale wetland relationship and we found little evidence for source-based differences in DOC chemical character in the literature.

[33] The significant positive correlation between  $\text{SUVA}_{254}$  and the HPOA fraction of bulk DOC was unsurprising because the HPOA fraction consists of aquatic fulvic and humic acids possessing greater aromaticity than other DOC fractions [*Aiken et al.*, 1992]. These results suggest that  $\text{SUVA}_{254}$  may be a useful surrogate for HPOA in organic carbon modeling applications [*Butman et al.*, 2012; *Spencer et al.*, 2012]. However, this relationship broke down within some individual basins (Mobile, Mississippi, and Colorado; Table 6), indicating that non-aromatic hydrophobic acids may constitute HPOA in some cases and that care should be taken when utilizing  $\text{SUVA}_{254}$  as a proxy for HPOA.

#### 4.3. Large River Versus Small Stream DOC Quantity and Chemical Character

[34] Several processes could lead to differences in DOC quantity and chemical character between small streams and large rivers including in-stream transformation of terrestrial headwater sources and additional DOC loads to large rivers not accounted for in small headwater catchment studies. While aquatic transformations may occur in smaller streams, we would expect their ability to influence DOC quantity and chemical character to be minimal at that point due to short residence times. Thus, headwater streams provide a signal of terrestrial influence. By the time water has reached large

**Table 6.** Table Showing Within-Basin Relationships Between Hydrophobic Organic Acid (HPOA) and Specific Ultraviolet Absorbance at 254 nm (SUVA<sub>254</sub>)<sup>a</sup>

| River Name         | HPOA-SUVA <sub>254</sub> formula | R <sup>2</sup> | p-value |
|--------------------|----------------------------------|----------------|---------|
| Penobscot River    | ND                               | ND             | ND      |
| Kennebec River     | ND                               | ND             | ND      |
| Androscoggin River | ND                               | ND             | ND      |
| Susquehanna River  | SUVA = 4.795 × HPOA + 0.346      | 0.64           | < 0.001 |
| Potomac River      | SUVA = 6.584 × HPOA + -0.306     | 0.58           | 0.003   |
| Edisto River       | SUVA = 4.988 × HPOA + 1.061      | 0.54           | 0.001   |
| Altamaha River     | SUVA = 8.267 × HPOA + -0.577     | 0.39           | < 0.001 |
| St. Mary's River   | SUVA = 2.738 × HPOA + 2.826      | 0.24           | 0.003   |
| Santa Fe River     | SUVA = 7.245 × HPOA + -0.19      | 0.87           | < 0.001 |
| Mobile River       | N/S                              | N/S            | 0.201   |
| St. Lawrence River | SUVA = 5.243 × HPOA + -0.18      | 0.64           | < 0.001 |
| Mississippi River  | N/S                              | N/S            | 0.346   |
| Rio Grande         | SUVA = 6.013 × HPOA + -0.102     | 0.48           | 0.001   |
| Colorado River     | N/S                              | N/S            | 0.199   |
| San Joaquin River  | SUVA = 4.807 × HPOA + 0.43       | 0.40           | 0.009   |
| Sacramento River   | SUVA = 8.07 × HPOA + -0.825      | 0.76           | < 0.001 |
| Columbia River     | SUVA = 8.754 × HPOA + -1.143     | 0.56           | 0.003   |

<sup>a</sup>Basins without a significant relationship are labeled N/S. ND indicates that no data were available.


rivers, aquatic processes such as microbial remineralization, sorption to sediments, and photodegradation could accumulate sufficiently to alter DOC [Aufdenkampe et al., 2011; McKnight et al., 2002; McKnight et al., 1992; Perez et al., 2011; Richey et al., 1990; Shih et al., 2010; Weyhenmeyer et al., 2012]. Sources unique to large rivers could also alter large river DOC relative to smaller headwaters, including accumulated aquatic DOC production from primary producers [Shih et al., 2010], different quantity or chemical character from large river floodplains compared to headwater wetlands, poorly understood groundwater inputs, and anthropogenic point sources [Butman et al., 2012] which tend to be located on larger rivers due to the distribution of urban centers.

[35] Given all the potential factors that could alter DOC, it was surprising that the terrestrial wetland signal for both DOC quantity and character was so clear across most of the large watersheds. Percent wetland cover alone explained most of the DOC concentration and SUVA<sub>254</sub> variability among basins at the annual scale. When we compared the concentration versus wetland relationship observed here (equation (3)) to those observed in previous headwater studies [Buffam et al., 2007; Creed et al., 2003; Eckhardt and Moore, 1990; Gergel et al., 1999; Gorham et al., 1998; Raymond and Hopkinson, 2003], we found little obvious difference in absolute concentration, slope, or intercept (Figure 6).

[36] However, the presence of lakes and reservoirs with long residence times appears to alter this pattern, as suggested by the differing DOC quantity and/or chemical character in the Colorado River and St. Lawrence River outliers. Across all basins, we did not find a relationship between either percent reservoir cover or percent lake cover and DOC or SUVA<sub>254</sub>. However, Lakes Mead and Powell on the Colorado River have a combined residence time of approximately 5 years [USBR-LC; USBR-UC], and Lake Ontario alone on the St. Lawrence has a residence time of approximately 6 years [Beltran et al., 1995]. The extremely long residence times in these two outlier rivers might have driven down SUVA<sub>254</sub> by some combination of autochthonous production of less aromatic DOC, biotic processing, and photodegradation [Miller, 2012; Spencer et al., 2012]. Previous studies suggest that in the Colorado River,

artificially flooded canyons like Lake Powell act as a trap for organic material entering from upstream. Over time, settled organic matter can remineralize or adsorb onto precipitating calcite [Reynolds, 1978]. Water discharged from Glen Canyon Dam is left nutrient rich and nearly free of suspended particles. These conditions also facilitate autochthonous production of weakly UV-absorbing DOC in the downstream reach [Henderson et al., 2008; Stanford and Ward, 1990].

[37] Extensive photodegradation and biological processing of terrestrially derived DOC during the long Great Lakes residence time and limited autochthonous production could explain the apparent conundrum in the St. Lawrence River



**Figure 6.** Dissolved organic carbon (DOC) concentration versus percent wetland cover relationships from Figure 2b, which excludes the St. Mary's River (red diamonds) and five previous small-basin studies.

of low DOC concentrations and very low SUVA<sub>254</sub> [Helie and Hillaire-Marcel, 2006]. Photodegradation acts primarily by breaking up strongly UV-absorbing molecules like terrestrial humic and fulvic acids [Moran and Zepp, 1997; Waiser and Robarts, 2004], which drive down SUVA<sub>254</sub> over time. Photodegradation of DOC has also been observed to result in an increase in its lability, leading to greater biotic remineralization [Anesio et al., 2005]. Thus, in addition to driving down SUVA<sub>254</sub> in the St. Lawrence River, photodegradation could lead to increased biotic processing and provide an explanation for its comparatively low DOC concentration (Figure 2b).

[38] SUVA<sub>254</sub> end-members identified in this study were generally similar to those found in the literature [Spencer et al., 2008; Weishaar et al., 2003], further suggesting that larger river DOC is similar to that in smaller headwater streams. Maximum mean SUVA<sub>254</sub> in the wetland dominated St. Mary's ( $4.7 \pm 0.3 \text{ L mg C}^{-1} \text{ m}^{-1}$ ) was similar to that previously reported from specific wetland sites ( $3.2$  to  $5.3 \text{ L mg C}^{-1} \text{ m}^{-1}$ ) [Weishaar et al., 2003]. However, mean SUVA<sub>254</sub> for the six river basins with less than 1% wetland cover (Colorado River excluded) was  $2.5 \pm 0.2 \text{ L mg C}^{-1} \text{ m}^{-1}$ , significantly higher than algal or microbially derived SUVA<sub>254</sub> end-members found by Stets et al. [2010] for closed-basin, seepage lakes which ranged from 0.9 to  $1.2 \text{ L mg C}^{-1} \text{ m}^{-1}$ . If biogeochemical processing played the primary role in driving SUVA<sub>254</sub> by breaking down aromatic allochthonous DOC, we would expect very low wetland systems to exhibit SUVA<sub>254</sub> values similar to previously identified end-members or values from very long residence time systems (St. Lawrence and Colorado Rivers: 1.3 and  $1.7 \text{ L mg C}^{-1} \text{ m}^{-1}$ , respectively). Rather, these results support the hypothesis proposed by del Giorgio and Pace [2008] and Richey et al. [1990] that labile autochthonous material may be rapidly recycled while more refractory, generally allochthonous DOC is delivered to the coastal ocean.

[39] We found that in large systems lacking long residence time in surface waters, DOC concentration and chemical character were well predicted by percent wetland cover. This suggests that for these systems at annual scales, in-stream processes like autochthonous production, biotic remineralization, and photodegradation play a subordinate role in driving DOC quantity and chemical character compared to processes that load allochthonous DOC into the river system. It is likely that any new production is being quickly recycled by the biotic community rather than delivered in large quantities to the coastal ocean [del Giorgio and Pace, 2008; Richey et al., 1990]. However, this finding is based on a relatively weak test that compares a mean annual DOC versus wetland relationship among heterogeneous large basins with snapshots from more homogeneous small basins in a variety of mostly northern regions. In addition, few comparative DOC chemical character studies in small basins have been conducted (but see Jaffe et al. [2008]). Finally, the tendency of smaller basins in this analysis to possess greater wetland coverage complicates the interpretation of our results because it is difficult to divorce the impact of basin size from wetlands. In order to fully evaluate the role of network-scale processing in driving annual DOC quantity and chemical character exported from large basins, more comprehensive, synoptic studies of DOC and DOC chemical character throughout river networks should be conducted.

## 5. Conclusion

[40] We demonstrated that wetlands play an integral role in controlling both DOC concentration and chemical character in 17 large and diverse North American rivers. The relationship between DOC and wetlands is similar to those previously reported for small watersheds. This result suggests that factors that may alter the terrestrial signal in large basins, such as in-stream processes, floodplain dynamics, or major anthropogenic inputs to large rivers, do not alter terrestrial control of carbon quantity and chemical character exports over annual time scales. The exceptions are surface water networks with long residence times, which suggest in-stream sources or transformations overwhelm the terrestrial signal. Our study is strengthened by the consistency of both DOC quantity and chemical character. These results supply valuable insight into the controls on mean DOC quantity and chemical character at broad spatial and temporal scales and provide new annual estimates of DOC flux and chemical character for several of the largest rivers in temperate North America. In order to better understand the source and fate of DOC in the environment, future efforts should include synoptic sampling of DOC quantity and chemical character, from headwaters through large river main stems, which would help unravel the hydrological processes that drive DOC variability within basins, the subsurface biogeochemical reactions that drive baseflow chemistry, and the biological and photolytic riverine processing that drives DOC quantity and chemical character over time.

[41] **Acknowledgments.** We are grateful to S. Glidden, A. Prusevich, and R. Stewart for data processing assistance, to C. Crawford (USGS) and the field teams of the USGS National Stream Quality Accounting Network and National Water Quality Assessment programs for assistance in fieldwork, and to K. Butler for laboratory assistance. This work was funded by National Aeronautics and Space Administration grants NNX09AU89G, NNX04AA62I, and NNX08A1571 as well as the U.S. Geological Survey's NASQAN, NAWQA, Climate and Land Use Change Program, and National Research Programs. This research was completed in partial fulfillment of the requirements for the degree of MS in Earth Science at the University of New Hampshire. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

## References

- Agren, A., I. Buffam, M. Berggren, K. Bishop, M. Jansson, and H. Laudon (2008), Dissolved organic carbon characteristics in boreal streams in a forest-wetland gradient during the transition between winter and summer, *J. Geophys. Res.-Biogeo.*, *113*, G03031, doi:10.1029/2007JG000674.
- Aiken, G. R., E. M. Thurman, R. L. Malcolm, and H. F. Walton (1979), Comparison of XAD macroporous resins for the concentration of fulvic acid from aqueous solution, *Anal. Chem.*, *51*(11), 1799–1803.
- Aiken, G. R., D. M. McKnight, K. A. Thorn, and E. M. Thurman (1992), Isolation of hydrophilic organic-acids from water using nonionic macroporous resins, *Org. Geochem.*, *18*(4), 567–573.
- Aiken, G. R., H. Hsu-Kim, and J. N. Ryan (2011), Influence of dissolved organic matter on the environmental fate of metals, nanoparticles, and colloids, *Environ. Sci. Technol.*, *45*(8), 3196–3201.
- Aitkenhead, J. A., and W. H. McDowell (2000), Soil C:N ratio as a predictor of annual riverine DOC flux at local and global scales, *Global Biogeochem. Cy.*, *14*(1), 127–138.
- Aitkenhead-Peterson, J. A., W. H. McDowell, and J. C. Neff (2003), Sources, production and regulation of allochthonous dissolved organic matter inputs to surface waters, in *Aquatic Ecosystems: Interactivity of Dissolved Organic Matter*, edited by S. E. G. Findlay, and R. L. Sinsabaugh, 519 p. Academic Press, London, UK.
- Alexander, R. B., J. R. Slack, A. S. Ludtke, K. K. Fitzgerald, and T. L. Schertz (1998), Data from selected US Geological Survey National Stream Water Quality Monitoring Networks, *Water Resour. Res.*, *34*(9), 2401–2405.

- Alexander, R. B., R. A. Smith, and G. E. Schwarz (2000), Effect of stream channel size on the delivery of nitrogen to the Gulf of Mexico, *Nature*, 403(6771), 758–761.
- Anesio, A. M., W. Graneli, G. R. Aiken, D. J. Kieber, and K. Mopper (2005), Effect of humic substance photodegradation on bacterial growth and respiration in lake water, *Appl Environ Microb*, 71(10), 6267–6275.
- Aufdenkampe, A. K., E. Mayorga, P. A. Raymond, J. M. Melack, S. C. Doney, S. R. Alin, R. E. Aalto, and K. Yoo (2011), Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere, *Front. Ecol. Environ.*, 9(1), 53–60.
- Beltran, R., L. Botts, P. Brown, T. Clarke, D. Cowell, K. Fuller, and B. Krushelnicki (1995), *The Great Lakes: An Environmental Atlas and Resource Book*, edited by K. Fuller, H. Shear, and J. Wittig, 46 p., Government of Canada and United States Environmental Protection Agency, Chicago, IL.
- Benner, R. (2003), Molecular indicators of the bioavailability of dissolved organic matter, in *Aquatic Ecosystems: Interactivity of Dissolved Organic Matter*, edited by S. E. G. Findlay, and R. L. Sinsabaugh, 519 p., Academic Press, London, UK.
- Booth, G., P. Raymond, and N. H. Oh (2007), LoadRunner, Software and website, Yale University, New Haven, CT, <http://environment.yale.edu/raymond/loadrunner>.
- Buffam, I., J. N. Galloway, L. K. Blum, and K. J. McGlathery (2001), A stormflow/baseflow comparison of dissolved organic matter concentrations and bioavailability in an Appalachian stream, *Biogeochemistry*, 53(3), 269–306.
- Buffam, I., H. Laudon, J. Temnerud, C. M. Morth, and K. Bishop (2007), Landscape-scale variability of acidity and dissolved organic carbon during spring flood in a boreal stream network, *J Geophys Res-Bioge*, 112, G01022, doi:10.1029/2006JG000218.
- Butman, D., P. A. Raymond, K. Butler, and G. Aiken (2012), Relationships between  $\Delta 14C$  and the molecular quality of dissolved organic carbon in rivers draining to the coast from the conterminous United States, *Global Biogeochem Cy*, 26(4), GB4014, doi:10.1029/2012GB004361.
- Chorover, J., and M. K. Amistadi (2001), Reaction of forest floor organic matter at goethite, birnessite and smectite surfaces, *Geochim Cosmochim Acta*, 65(1), 95–109.
- Clair, T. A., and J. M. Ehrman (1996), Variations in discharge and dissolved organic carbon and nitrogen export from terrestrial basins with changes in climate: A neural network approach, *Limnol. Oceanogr.*, 41(5), 921–927.
- Cole, J. J., et al. (2007), Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon budget, *Ecosystems*, 10(1), 171–184.
- Creed, I. F., S. E. Sanford, F. D. Beall, L. A. Molot, and P. J. Dillon (2003), Cryptic wetlands: integrating hidden wetlands in regression models of the export of dissolved organic carbon from forested landscapes, *Hydrol. Process.*, 17(18), 3629–3648.
- Daly, C., and G. Taylor (1998), *United States Average Monthly or Annual Mean Temperature, 1961–90, Spatial Climate Analysis Service at Oregon State University*, Spatial Climate Analysis Service at Oregon State University (SCAS/OSU), Corvallis, OR.
- Dalzell, B. J., T. R. Filley, and J. M. Harbor (2007), The role of hydrology in annual organic carbon loads and terrestrial organic matter export from a midwestern agricultural watershed, *Geochim Cosmochim Acta*, 71(6), 1448–1462.
- Dittman, J. A., J. B. Shanley, C. T. Driscoll, G. R. Aiken, A. T. Chalmers, J. E. Towse, and P. Selvendiran (2010), Mercury dynamics in relation to dissolved organic carbon concentration and quality during high flow events in three northeastern US streams, *Water Resour. Res.*, 46, W07522, doi:10.1029/2009WR008351.
- Eckhardt, B. W., and T. R. Moore (1990), Controls on dissolved organic-carbon concentrations in streams, southern Quebec, *Can. J. Fish. Aquat. Sci.*, 47(8), 1537–1544.
- Frey, K. E., and L. C. Smith (2005), Amplified carbon release from vast West Siberian peatlands by 2100, *Geophys. Res. Lett.*, 32, L09401, doi:10.1029/2004GL020205.
- Frost, P. C., J. H. Larson, C. A. Johnston, K. C. Young, P. A. Maurice, G. A. Lamberti, and S. D. Bridgman (2006), Landscape predictors of stream dissolved organic matter concentration and physicochemistry in a Lake Superior river watershed, *Aquat. Sci.*, 68(1), 40–51.
- Gattuso, J. P., M. Frankignoulle, and R. Wollast (1998), Carbon and carbonate metabolism in coastal aquatic ecosystems, *Annu. Rev. Ecol. Syst.*, 29, 405–434.
- Gergel, S. E., M. G. Turner, and T. K. Kratz (1999), Dissolved organic carbon as an indicator of the scale of watershed influence on lakes and rivers, *Ecol. Appl.*, 9(4), 1377–1390.
- del Giorgio, P. A., and J. Davis (2003), Patterns in dissolved organic matter lability and consumption across aquatic ecosystems, in *Aquatic Ecosystems: Interactivity of Dissolved Organic Matter*, edited by S. E. G. Findlay, and R. L. Sinsabaugh, 519 p., Academic Press, London, UK.
- del Giorgio, P. A., and M. L. Pace (2008), Relative independence of dissolved organic carbon transport and processing in a large temperate river: The Hudson River as both pipe and reactor, *Limnol. Oceanogr.*, 53(1), 185–197.
- Gorham, E., J. K. Underwood, J. A. Janssens, B. Freedman, W. Maass, D. H. Waller, and J. G. Ogden (1998), The chemistry of streams in southwestern and central Nova Scotia, with particular reference to catchment vegetation and the influence of dissolved organic carbon primarily from wetlands, *Wetlands*, 18(1), 115–132.
- Guillemette, F., and P. A. del Giorgio (2011), Reconstructing the various facets of dissolved organic carbon bioavailability in freshwater ecosystems, *Limnol. Oceanogr.*, 56(2), 734–748.
- Harrison, J. A., N. Caraco, and S. P. Seitzinger (2005), Global patterns and sources of dissolved organic matter export to the coastal zone: Results from a spatially explicit, global model, *Global Biogeochem Cy*, 19(4), GB4S04, doi:10.1029/2005GB002480.
- Helie, J. F., and C. Hillaire-Marcel (2006), Sources of particulate and dissolved organic carbon in the St Lawrence River: Isotopic approach, *Hydrol. Process.*, 20(9), 1945–1959.
- Henderson, R. K., A. Baker, S. A. Parsons, and B. Jefferson (2008), Characterisation of algal organic matter extracted from cyanobacteria, green algae and diatoms, *Water Res.*, 42(13), 3435–3445.
- Hodgkins, G. A. (1999), Estimating the magnitude of peak flows for streams in Maine for selected recurrence intervals, USGS *Water-Resources Investigations Report. 99-4008*, 45 pp, U.S. Geological Survey.
- Homer, C., C. Q. Huang, L. M. Yang, B. Wylie, and M. Coan (2004), Development of a 2001 national land-cover database for the United States, *Photogramm Eng Rem S*, 70(7), 829–840.
- IBWC (2010), Rio Grande stream gauge, edited, International Boundary and Water Commission. [http://www.ibwc.state.gov/Water\\_Data/](http://www.ibwc.state.gov/Water_Data/)
- Jaffe, R., D. McKnight, N. Maie, R. Cory, W. H. McDowell, and J. L. Campbell (2008), Spatial and temporal variations in DOM composition in ecosystems: The importance of long-term monitoring of optical properties, *J Geophys Res-Bioge*, 113, G04032, doi:10.1029/2008JG000683.
- Lauerwald, R., J. Hartmann, W. Ludwig, and N. Moosdorf (2012), Assessing the nonconservative fluvial fluxes of dissolved organic carbon in North America, *J. Geophys. Res.*, 117, G01027, doi:10.1029/2011JG001820.
- Leenheer, J. (1982), United States Geological Survey data information service, in *Transport of Carbon and Minerals in Major World Rivers*, edited by E. T. Degens, pp. 355–356, Universitat Hamburg, Hamburg.
- Lehner, B., and P. Doll (2004), Development and validation of a global database of lakes, reservoirs and wetlands, *J. Hydrol.*, 296(1-4), 1–22.
- Lehner, B., et al. (2011), High-resolution mapping of the world's reservoirs and dams for sustainable river-flow management, *Front. Ecol. Environ.*, 9(9), 494–502.
- Lehtoranta, J., P. Ekholm, and H. Pitkanen (2009), Coastal eutrophication thresholds: A matter of sediment microbial processes, *Ambio*, 38(6), 303–308.
- Ludwig, W., J. L. Probst, and S. Kempe (1996), Predicting the oceanic input of organic carbon by continental erosion, *Global Biogeochem Cy*, 10(1), 23–41.
- Maurice, P. A., M. J. Pullin, S. E. Cabaniss, Q. H. Zhou, K. Namjesnik-Dejanovic, and G. R. Aiken (2002), A comparison of surface water natural organic matter in raw filtered water samples, XAD, and reverse osmosis isolates, *Water Res.*, 36(9), 2357–2371.
- McKnight, D. M., K. E. Bencala, G. W. Zellweger, G. R. Aiken, G. L. Feder, and K. A. Thorn (1992), Sorption of dissolved organic carbon by hydrous aluminum and iron oxides occurring at the confluence of Deer Creek with the Snake River, Summit County, Colorado, *Environ. Sci. Technol.*, 26(7), 1388–1396.
- McKnight, D. M., G. M. Hornberger, K. E. Bencala, and E. W. Boyer (2002), In-stream sorption of fulvic acid in an acidic stream: A stream-scale transport experiment, *Water Resour. Res.*, 38(1) doi:10.1029/2001WR000269.
- Meier, M., K. Namjesnik-Dejanovic, P. A. Maurice, Y. P. Chin, and G. R. Aiken (1999), Fractionation of aquatic natural organic matter upon sorption to goethite and kaolinite, *Chem. Geol.*, 157(3-4), 275–284.
- Meybeck, M., and A. Ragu (1996), River discharges to the oceans: An assessment of suspended solids, major ions, and nutrients, in *Environment and Assessment*, edited by U. E. Programme, pp. 1–245, Nairobi, Kenya.
- Miller, M. P. (2012), The influence of reservoirs, climate, land use and hydrologic conditions on loads and chemical quality of dissolved organic carbon in the Colorado River, *Water Resour. Res.*, 48, W00M02.
- Moran, M. A., and R. G. Zepp (1997), Role of photoreactions in the formation of biologically labile compounds from dissolved organic matter, *Limnol. Oceanogr.*, 42(6), 1307–1316.
- Mulholland, P. J., and E. J. Kuenzler (1979), Organic-carbon export from upland and forested wetland watersheds, *Limnol. Oceanogr.*, 24(5), 960–966.

- Mulholland, P. J., and J. A. Watts (1982), Transport of organic carbon to the oceans by rivers of North America—A synthesis of existing data, *Tellus*, 34(2), 176–186.
- Nash, J. E., and J. V. Sutcliffe (1970), River flow forecasting through conceptual models part I—A discussion of principles, *J. Hydrol.*, 10(3), 282–290.
- Neale, P. A., A. Antony, W. Gernjak, G. Leslie, and B. I. Escher (2011), Natural versus wastewater derived dissolved organic carbon: Implications for the environmental fate of organic micropollutants, *Water Res.*, 45(14), 4227–4237.
- NOAA-NCDC (2012), Climate at a Glance (National), *National Oceanic and Atmospheric Administration, National Climatic Data Center*. <http://www.ncdc.noaa.gov/oa/climate/research/cag3/na.html>
- NRCan/CCRS, USGS, INEGI, CONABIO, and CONAFOR (2010), 2005 Land Cover of North America at 250 meters, Natural Resources Canada/Canadian Centre for Remote Sensing (NRCan/CCRS), United States Geological Survey (USGS); Instituto Nacional de Estadística y Geografía (INEGI), Comisión Nacional para el Conocimiento y Uso de la Biodiversidad (CONABIO) and Comisión Nacional Forestal (CONAFOR), Sioux Falls, SD. <http://www.ccc.org/naatlas/>
- Perez, M. A. P., P. Moreira-Turcq, H. Gallard, T. Allard, and M. F. Benedetti (2011), Dissolved organic matter dynamic in the Amazon basin: Sorption by mineral surfaces, *Chem. Geol.*, 286(3–4), 158–168.
- Raymond, P. A., and C. S. Hopkins (2003), Ecosystem modulation of dissolved carbon age in a temperate marsh-dominated estuary, *Ecosystems*, 6(7), 694–705.
- Reynolds, R. C. (1978), Polyphenol inhibition of calcite precipitation in Lake Powell, *Limnol. Oceanogr.*, 23(4), 585–597.
- Richey, J. E., J. I. Hedges, A. H. Devol, P. D. Quay, R. Victoria, L. Martinelli, and B. R. Forsberg (1990), Biogeochemistry of carbon in the Amazon River, *Limnol. Oceanogr.*, 35(2), 352–371.
- Runkel, R. L., C. G. Crawford, and T. A. Cohn (2004), Load Estimator (LOADEST): A FORTRAN program for estimating constituent loads in streams and rivers, in *United States Geological Survey Techniques and Methods*, edited, 69 pp.
- Schlesinger, W. H. (1997), *Biogeochemistry: An Analysis of Global Change*, 2nd edn., xiii, 588 p., Academic Press, San Diego, Calif.
- Seitzinger, S. P., J. A. Harrison, E. Dumont, A. H. W. Beusen, and A. F. Bouwman (2005), Sources and delivery of carbon, nitrogen, and phosphorus to the coastal zone: An overview of Global Nutrient Export from Watersheds (NEWS) models and their application, *Global Biogeochem. Cy.*, 19(4), GB4S01, doi:10.1029/2005GB002606.
- Shih, J. S., R. B. Alexander, R. A. Smith, E. W. Boyer, G. E. Schwarz, and S. Chung (2010), An initial SPARROW model of land use and in-stream controls on total organic carbon in streams of the conterminous United States, *USGS Open-File Report*(2010-1276), 22 p.
- Sholkovitz, E. R. (1976), Flocculation of dissolved organic and inorganic matter during mixing of river water and seawater, *Geochim Cosmochim. Acta*, 40(7), 831–845.
- Singer, P. C. (1999), Humic substances as precursors for potentially harmful disinfection by-products, *Water Sci. Technol.*, 40(9), 25–30.
- Spencer, R. G. M., G. R. Aiken, K. P. Wickland, R. G. Striegl, and P. J. Hernes (2008), Seasonal and spatial variability in dissolved organic matter quantity and composition from the Yukon River basin, Alaska, *Global Biogeochem. Cy.*, 22(4), GB4002, doi:10.1029/2008GB003231.
- Spencer, R. G. M., K. D. Butler, and G. R. Aiken (2012), Dissolved organic carbon and chromophoric dissolved organic matter properties of rivers in the USA, *J. Geophys. Res.-Biogeo.*, 117, G03001, doi:10.1029/2011JG001928.
- Stanford, J. A., and J. V. Ward (1990), *Limnology of Lake Powell and the chemistry of the Colorado River Rep.*, 279 pp, Washington, D.C.
- Stets, E. G., R. G. Striegl, and G. R. Aiken (2010), Dissolved organic carbon export and internal cycling in small, headwater lakes, *Global Biogeochem. Cy.*, 24, GB4008, doi:10.1029/2010GB003815.
- Tipping, E., et al. (1999), Climatic influences on the leaching of dissolved organic matter from upland UK Moorland soils, investigated by a field manipulation experiment, *Environ. Int.*, 25(1), 83–95.
- USBR-LC (2012), Water Resources Group-Lower Colorado Region, edited by United States Bureau of Reclamation. <http://www.usbr.gov/lc/>
- USBR-UC (2012), Water Resources Group-Upper Colorado Region, edited by United States Bureau of Reclamation. <http://www.usbr.gov/uc/water/>
- USGS (2006), National Hydrography Dataset (NHDPlus). <http://www.horizon-systems.com/nhdplus/>
- Waiser, M. J., and R. D. Robarts (2004), Photodegradation of DOC in a shallow prairie wetland: Evidence from seasonal changes in DOC optical properties and chemical characteristics, *Biogeochemistry*, 69(2), 263–284.
- Weishaar, J. L., G. R. Aiken, B. A. Bergamaschi, M. S. Fram, R. Fujii, and K. Mopper (2003), Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition and reactivity of dissolved organic carbon, *Environ. Sci. Technol.*, 37(20), 4702–4708.
- Weyhenmeyer, G. A., M. Froberg, E. Karlun, M. Khalili, D. Kothawala, J. Temnerud, and L. J. Tranvik (2012), Selective decay of terrestrial organic carbon during transport from land to sea, *Global Change Biol.*, 18(1), 349–355.