

University of New Hampshire

University of New Hampshire Scholars' Repository

NHAES Bulletin

New Hampshire Agricultural Experiment Station

4-1-1985

Aquatic vascular plants of New England, Station Bulletin, no.528

Crow, G. E.

Hellquist, C. B.

New Hampshire Agricultural Experiment Station

Follow this and additional works at: <https://scholars.unh.edu/agbulletin>

Recommended Citation

Crow, G. E.; Hellquist, C. B.; and New Hampshire Agricultural Experiment Station, "Aquatic vascular plants of New England, Station Bulletin, no.528" (1985). *NHAES Bulletin*. 489.

<https://scholars.unh.edu/agbulletin/489>

This Text is brought to you for free and open access by the New Hampshire Agricultural Experiment Station at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in NHAES Bulletin by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

310Sci
3
89
E21
no. 528

MON BULLETIN 528


BIO SCI
LIBRARY

April, 1985

**Aquatic Vascular Plants of New England:
Part 8. Lentibulariaceae**

by

G. E. Crow and C. B. Hellquist



**NEW HAMPSHIRE
AGRICULTURAL EXPERIMENT STATION
UNIVERSITY OF NEW HAMPSHIRE
DURHAM, NEW HAMPSHIRE 03824**

UNIVERSITY OF NEW HAMPSHIRE
LIBRARY

ISSN: 0077-8338

noSci
2
9
E21
no. 528

ION BULLETIN 528


BIO SCI
LIBRARY

April, 1985

**Aquatic Vascular Plants of New England:
Part 8. Lentibulariaceae**

by

G. E. Crow and C. B. Hellquist



**NEW HAMPSHIRE
AGRICULTURAL EXPERIMENT STATION
UNIVERSITY OF NEW HAMPSHIRE
DURHAM, NEW HAMPSHIRE 03824**

UNIVERSITY OF NEW HAMPSHIRE
LIBRARY

ISSN: 0077-8338

ACKNOWLEDGEMENTS

We wish to thank Drs. Robert K. Godfrey and George B. Rossbach for their helpful comments on the manuscript. We are also grateful to the curators of the following herbaria for use of their collections: BRU, CONN, CUW, GH, NHN, KIRI, MASS, MAINE, NASC, NCBS, NHA, NEBC, VT, YU. A special thanks is extended to Pamela Bruns Brayton who prepared the illustrations. Gratitude is expressed to the University of Wisconsin Press for permission to use or redraw several figures from Fassett's *Manual of Aquatic Plants*.

This work is a result of research sponsored by the New Hampshire Agricultural Experiment Station. The NHAES reserves the right to reproduce, publish or otherwise use, and to authorize others to use, the work for Government purposes notwithstanding notice of copyright.

Copyright 1985 by the University of New Hampshire. No part of this work may be reproduced in any manner without permission from the authors and the University of New Hampshire.

Programs of the New Hampshire Agricultural Experiment Station are open to all persons without regard to race, color, national origin or sex. The University of New Hampshire is an Affirmative Action/Equal Opportunity Employer.



ABSTRACT

This paper is the eighth in a series of reports on the aquatic and wetland flora of New England. It treats all species of the Lentibulariaceae occurring in New England and includes keys, comments on taxonomy and nomenclature, habitat and distributional information, water chemistry data, illustrations and dot maps. Those species regarded as rare and endangered in the New England Region or in one or more of the six New England States are also noted.

KEY WORDS: Aquatic Plants, New England Flora, Taxonomy, Lentibulariaceae, *Utricularia*, Bladderwort.

1210 521
S
89
.221
no 528

TABLE OF CONTENTS

	Page
INTRODUCTION	1
LENTIBULARIACEAE	2
<i>Utricularia</i>	2
Key to Species	2
<i>Utricularia fibrosa</i>	4
<i>Utricularia intermedia</i>	4
<i>Utricularia radiata</i>	4
<i>Utricularia purpurea</i>	4
<i>Utricularia minor</i>	10
<i>Utricularia geminiscapa</i>	10
<i>Utricularia vulgaris</i>	10
<i>Utricularia gibba</i>	13
<i>Utricularia biflora</i>	13
<i>Utricularia subulata</i>	13
<i>Utricularia cornuta</i>	13
<i>Utricularia resupinata</i>	17
Literature Cited and Selected References	19

Aquatic Vascular Plants of New England: Part 8. Lentibulariaceae

by

G. E. Crow and C.B. Hellquist

INTRODUCTION

This is the eighth in a series of reports on the aquatic and wetland flora of New England. These reports are intended to aid conservationists, fish and game personnel, consultants, botanists, and students in the identification of aquatic plants. The coverage is strictly New England but is of value throughout the northeast. Data have been gathered from herbaria in New England and from personal field work.

Chemical data presented represent samples from many waters throughout New England. The alkalinity readings are total alkalinity, expressed as milligrams per liter (mg/l) CaCO_3 . The number of observations is given in parentheses following alkalinity and pH values. Since pH and alkalinity vary greatly during the day, the values are only indicative of the water quality.

The rare and endangered plant lists referred to are those prepared for each of the six New England states by the New England Botanical Club in cooperation with the United States Fish and Wildlife Service, Office of Endangered Species, Newton Corner, MA (RI — Church and Champlin, 1978; MA — Coddington and Field, 1978; VT — Countryman, 1978; ME — Eastman, 1978; CT — Mehrhoff, 1978; NH — Storks and Crow, 1978). Taxa indicated as rare, threatened or endangered for the entire New England Region are also noted (Crow *et al.*, 1981, *Rhodora* 83: 259-299.)

We invite comments and/or criticisms on this treatment. Information on any species omitted or any additional localities will be welcomed. If anyone is interested in specific localities of any of the species indicated on the dot maps, please contact us.

Dr. Garrett E. Crow, Department of Botany and Plant Pathology, Nesmith Hall, University of New Hampshire, Durham, NH 03824.

Dr. C. Barre Hellquist, Department of Biology, North Adams State College, North Adams, MA 01247.

LENTIBULARIACEAE

Utricularia (Bladderwort)

Perennial aquatic herbs; plants consisting of stem systems, roots absent; lateral leaf-like branches green, usually bearing bladder-like carnivorous traps, bladders sometimes borne on separate, non-photosynthetic branches; flowers borne on scapes elevated above water surface, solitary or in racemes; fruit a 2-valved capsule; perennating by winter buds (turions).

Key to Species

1. Plants floating in the water or creeping over surface of substrate (sometimes appearing anchored).
 2. Plants with two distinct types of branches, green leaf-like branches lacking bladders, and bladder-bearing branches lacking chlorophyll, sometimes appearing anchored in substrate (figs. 1A,2A).
 3. Segments of leaf-like branches fine, becoming narrower with each dichotomy, ultimate segments thread-like, not toothed (fig. 1B); corolla pale-yellow to greenish-yellow, lower and upper lips nearly equal (fig. 1A).
..... 1. *U. fibrosa*
 3. Segments of leaf-like branches distinctly flat, segments remaining approximately the same width, divisions minutely toothed (fig. 2B); corolla bright yellow, lower lip nearly twice as long as upper lip (fig. 2A).
..... 2. *U. intermedia*
 2. Plants with branches all alike, leaf-like, bearing bladders.
 4. Flowering scapes subtended by a whorl of inflated branches, floats holding the flowers above the water surface (fig. 3A).
..... 3. *U. radiata*
 4. Flowering scapes not subtended by inflated branches (floats).
 5. Leaf-like branches all whorled (fig. 4F); bladders borne at tips of branches (fig. 4F); flowers purple (rarely albino).
..... 4. *U. purpurea*
 5. Leaf-like branches chiefly alternate; bladders borne scattered on branches; flowers yellow.
 6. Leaf-like segments flat; spur of corolla very short, saccate, almost absent.
..... 5. *U. minor*
 6. Leaf-like segments terete; spur of corolla, 3 mm or more long.
 7. Leaf-like branches divided 6 or more times; free-floating; plants large, stout, 30-100 cm long.

8. Scapes with 2-5 flowers; bracts subtending flowers unlobed; submersed, bladder-bearing branches often with small, apetalous cleistogamous flowers (fig. 4D); branches finer, segments entire, except for small spicules at tips (magnification needed).
 6. *U. geminiscapa*
8. Scapes with 6-12 flowers; bracts subtending flowers with basal lobes; cleistogamous flowers absent; branches coarser, segments with spicules along margins (fig. 4B) (magnification needed).
 7. *U. vulgaris*
7. Leaf-like branches divided 1-3 times; creeping over wet mud or in shallow water, or in tangled masses floating in shallow water; plants small, slender, less than 30 cm long.
9. Flower spur much shorter than lower lip of corolla, thick, obtuse, 3.0-3.5 mm long (fig. 1F); lower lip of corolla 5-6 mm long; leaf-like branches with mostly 2 segments.
 8. *U. gibba*
9. Flower spur somewhat shorter to slightly longer than lower lip of corolla, narrow, conic, usually curved, more than 4 mm long; lower lip of corolla 8-10(-15) mm long; leaf-like branches with 3 or more segments (usually forked twice).
 9. *U. biflora*
1. Plants anchored in substrate, apparent only when flowering, appearing terrestrial (vegetative grass-like branches sometimes emergent, or sometimes floating in densely tangled mats due to wave action (fig. 5I).
10. Flowers yellow (rarely purplish or white); bracts along scape alternate (fig. 5A,D); flowers (1-)2-12.
11. Flowers clearly pedicellate; bract at base of pedicel peltate (fig. 5B); spur of corolla appressed to lower lip (fig. 5C).
 10. *U. subulata*
11. Flowers sessile; bract at base of flower attached basally (fig. 5E); spur of corolla divergent (fig. 5D).
 11. *U. cornuta*
10. Flowers purple; bracts along scape paired, fused (fig. 5F,G); flowers 1.
 12. *U. resupinata*

1. *Utricularia fibrosa* Walt. Fig. 1, Map 1

Rare in pools, wet peat, and long sandy shores of southeastern Massachusetts. This species is easily confused with *U. biflora*, but is more robust and has two distinct branch types, one with bladders and one without. Range extends from southeastern Massachusetts and central Connecticut south along the Coastal Plain to Florida and along the Gulf Coast to eastern Oklahoma and eastern Texas; California.

Rare and endangered plant lists: New England, Massachusetts, Connecticut

2. *Utricularia intermedia* Hayne Fig. 2, Map 2

Common in shallow water, peaty soils, and damp sands throughout New England. Although more abundant in acidic waters this species is often found in moderately alkaline waters. The green, leaf-like branches above the substrate lack bladders, hence the plant may not be readily recognized as a bladderwort until the plant is removed from the substrate and the bladders observed. Range extends from Greenland, Newfoundland, and Nova Scotia west to Alaska, south to New England, Long Island, New York, northern Delaware, Pennsylvania, Ohio, northern Indiana, northern Illinois, northwestern Iowa, and California.

alkalinity: mean 9.7 mg/l; range 3.5-52.5 mg/l (17)

pH: mean 6.7; range 5.9-7.9 (17)

3. *Utricularia radiata* Small Fig. 3, Map 3

Common in acidic waters of southern New England, widely scattered in northeastern New England. This taxon is treated by some authors as *Utricularia inflata* Walt. var. *minor* Chapm., but Reinert and Godfrey (1962) regard it as a distinct species from the more robust *Utricularia inflata* Walt. of the southeast. This taxon is easily identified in the flowering state, but sterile, submersed material is occasionally difficult to identify. Range extends from Nova Scotia and New England south along the Coastal Plain to southern Florida and westward along the Gulf Coast to eastern Texas; westward inland to Pennsylvania, northwestern Indiana, West Virginia, and Tennessee.

Rare and endangered plant list: Vermont

alkalinity: mean 7.8 mg/l; range 1.0-33.0 mg/l (26)

pH: mean 6.6; range 5.6-7.3 (27)

4. *Utricularia purpurea* Walt. Fig. 4, Map 4

Common, free-floating in acidic waters of eastern and southern New England, rarely in moderately alkaline waters. This species often flowers sporadically in many populations while flowering regu-




Figure 1

Utricularia fibrosa: A. habit $\times 1$. B. leaf-like branch $\times 4$.


C. spur of corolla $\times 2$, redrawn from Fassett.

Utricularia biflora: D. habit $\times 1$. E. spur of corolla $\times 2$, redrawn from Fassett.

Utricularia gibba: F. habit $\times 1$.



Map 2
Utricularia intermedia



Map 1
Utricularia fibrosa




Figure 2

Utricularia intermedia: A. habit $\times 1$. B. portion of leaf-like branch lacking bladders $\times 8$. C. bract, side and back views, $\times 4$. D. winter bud $\times 1$. All from Fassett.
Utricularia minor: E. habit $\times 1$. F. portion of leaf-like branch bearing bladder $\times 10$. G. winter bud $\times 1$, from Fassett.






Figure 3
Utricularia radiata: A. habit $\times 1$, redrawn from Fassett.



Map 4
Utricularia purpurea



Map 3
Utricularia radiata

larly and abundantly in others. The flower color ranges from purple to deep pink and rarely white (forma *alba* Hellquist). Range extends from Newfoundland, and Nova Scotia, west to southern Ontario, northern Indiana, Michigan, and Wisconsin, south along the Coastal Plain to Florida, Louisiana, and Texas.

Rare and endangered plant list: Vermont

alkalinity: mean 7.8 mg/l; range 2.5-33.0 mg/l (39)

ph: mean 6.6; range 5.6-7.3 (36)

5. *Utricularia minor* L. Fig. 2, Map 5

Widely scattered in acidic, alkaline, and slightly saline waters, wet meadows, bogs, and shores throughout New England. This species is rarely found in flower, but is easily identified by its flattened stems. Range extends from Greenland, Labrador, and Newfoundland west to British Columbia, south to Nova Scotia, New England, Long Island, New York, northern New Jersey, Pennsylvania, northern Ohio, northern Indiana, northern Illinois, northwestern Iowa, Nebraska, Colorado, and California.

Rare and endangered plant list: Vermont

alkalinity: mean 9.6 mg/l; range 3.0-33.5 mg/l (7)

pH: mean 6.9; range 6.0-8.1 (7)a

6. *Utricularia geminiscapa* Benj. Fig. 4, Map 6

Widely scattered in bog waters, acidic or moderately alkaline waters throughout New England. This species may be mistaken for a diminutive *Utricularia vulgaris*, thus flowering specimens are needed for accurate identification. Range extends from Newfoundland and New Brunswick west to Michigan and Wisconsin, south to New England, New York, eastern Pennsylvania, Delaware, and eastern Virginia.

Rare and endangered plant list: Vermont

alkalinity: mean 24.6 mg/l; range 5.4-69.5 mg/l (5)

pH: mean 6.7; range 3.5-8.6 (5)

7. *Utricularia vulgaris* L. Fig. 4, Map 7

Common and abundant throughout New England in acidic and alkaline waters. This is the most robust and common bladderwort in New England. Some stems range up to one meter in length. Range extends from southern Labrador west to Alaska, south to Virginia, North Carolina, Ohio, Indiana, Missouri, Oklahoma, and northern Texas; California and Mexico.

alkalinity: mean 18.6 mg/l; range 2.5-109.0 mg/l (75)

pH: mean 6.7; range 5.7-8.6 (70)





Figure 4


Utricularia vulgaris: A. habit $\times 1$. B portion of leaf-like branch bearing spicules $\times 6$. C. winter bud $\times 1$, from Fassett.

Utricularia geminiscapa: D. habit (cleistogamous flower at arrow) $\times 1$. E. winter bud $\times 1$, from Fassett.

Utricularia purpurea: F. habit $\times 1$. G. winter bud $\times 1$.



Map 6
Utricularia geminisca



Map 5
Utricularia minor

8. *Utricularia gibba* L. Fig. 1, Map 8

Common in shallow water, bogs, and along sandy shores of southern New England, widely scattered in northern New England. In New England this species is much more abundant in acidic sites, while westward it is often found in wet marly sites of higher alkalinity. Range extends from Nova Scotia, southern Quebec, and New England west to New York, southern Ontario, Michigan, Wisconsin, and Minnesota, south to Florida, Oklahoma, and Texas; California and Mexico.

alkalinity: mean 15.6 mg/l; range 3.0-69.5 mg/l (16)

pH: mean 6.7; range 5.8-8.6 (14)

9. *Utricularia biflora* Lam. Fig. 1, Map 9

Rare in shallow pools, wet peat, and along sandy shores of southern New England. This species is easily confused with the more common *U. gibba* and the rarer *U. fibrosa*, and is somewhat intermediate between them. Range extends from the Coastal Plain of eastern Massachusetts, southern Rhode Island, and southern Connecticut south to Florida and west along the Gulf Coast to eastern Oklahoma and southeastern Texas.

Rare and endangered plant list: New England, Massachusetts, Connecticut

10. *Utricularia subulata* L. Fig. 5, Map 10

Rare in wet peats and sands in southeastern New England. Presently this species is known in New England from only two extant sites, one in southeastern Massachusetts and one on Nantucket Island. Range extends along the Coastal Plain from southwestern Nova Scotia, southeastern Massachusetts, Rhode Island, and Long Island, New York, to Florida and along the Gulf Coast to Texas; inland to Tennessee, Arkansas, and eastern Oklahoma.

Rare and endangered plant list: New England

11. *Utricularia cornuta* Michx. Fig. 5, Map 11

Common in sphagnum bogs, wet sands, and along sandy or peaty shores in New England. More abundant in acidic regions of New England, but also found in the alkaline regions of northeastern Maine and northern Vermont. This species is inconspicuous, except when in flower. The bladder bearing, subterranean vegetative material may be found floating in densely tangled mats after being washed free by wave action. Range extends from Newfoundland and Quebec west to northern Ontario, Wisconsin, and Minnesota, south to Dela-




Figure 5


Utricularia subulata: A. habit showing subterranean bladder-bearing branches and vegetative grass-like branches $\times 1$. B. peltate bract $\times 8$. C. corolla showing spur $\times 2$.

Utricularia cornuta: D. habit showing subterranean bladder-bearing branches and vegetative grass-like branches $\times 1$. E. bract $\times 4$. Both from Fassett.


Utricularia resupinata: F. habit showing subterranean bladder-bearing branches and vegetative grass-like branches $\times 1$. G. paired bracts $\times 4$, from Fassett. H. corolla showing divergent spur $\times 2$, from Fassett. I. mat of floating, densely tangled branches.




Map 8
Utricularia gibba



Map 7
Utricularia vulgaris



Map 10
Utricularia subulata



Map 9
Utricularia biflora

ware, Pennsylvania, Ohio, northern Indiana, and northern Illinois; along the Coastal Plain to Florida and eastern Texas.


alkalinity: mean 3.7 mg/l; range 1.0-7.0 mg/l (6)

pH: mean 6.4; range 5.6-6.9 (6)


12. *Utricularia resupinata* B. D. Greene Fig. 5, Map 12

Uncommon in acidic regions of southern and central New England. This species, like *U. subulata* and *U. cornuta*, is difficult to identify in the sterile state and is usually overlooked. Many of the historical sites have been revisited to search for this species, but to no avail. This species seems to flower more often when the water recedes during dry seasons. Range extends from Nova Scotia west to Quebec, northern Michigan, and northwestern Wisconsin south to New York, northern Pennsylvania, northern Indiana, and northeastern Illinois; along the Coastal Plain to Georgia, Florida, and Alabama.

Rare and endangered plant list: Vermont



Map 12
Utricularia resupinata




Map 11
Utricularia cornuta

Literature Cited and
Selected References

- Hellquist, C. B. 1974. A white-flowered form of *Utricularia purpurea* from New Hampshire. *Rhodora* 76: 805.
- Lloyd, F. E. 1933. The structure and behavior of *Utricularia purpurea*. *Canad J. Res.* 8: 234-252.
- Lloyd, F. E. 1935. The traps of *Utricularia*. *Proc. 6th Bot. Congr.* 1: 54-73.
- Lloyd, F. E. 1935a. *Utricularia*. *Biol. Rev.* 10: 72-100.
- Reinert, G. W. and R. K. Godfrey. 1962. Reappraisal of *Utricularia inflata* and *U. radiata* (Lentibulariaceae). *Amer. J. Bot.* 49: 213-220.
- Roszbach, G. B. 1939. Aquatic Utricularias. *Rhodora* 41: 114-128.
- Uttal, L. J. 1960. Notes on *Utricularia biflora* and *U. fibrosa*. *Rhodora* 58: 41.

Station Bulletins of Botanical Interest

- Grasses of New Hampshire. I. Tribes Poeae (Festuceae) and Triticeae (Hordeae). A. R. Hodgdon, G. E. Crow, and F. L. Steele. Bull. No. 512. 1979.
- The Flora of Plum Island, Essex County, Massachusetts. M. J. McDonnell. Bull. No. 513. 1979.
- Aquatic Vascular Plants of New England: Part 1. Zosteraceae, Potamogetonaceae, Zannichelliaceae, Najadaceae. C. B. Hellquist and G. E. Crow. Bull. No. 515. 1980.
- Aquatic Vascular Plants of New England: Part 2. Typhaceae and Sparganiaceae. G. E. Crow and C. B. Hellquist. Bull. No. 517. 1981.
- Aquatic Vascular Plants of New England: Part 3. Alismataceae. C. B. Hellquist and G. E. Crow. Bull. No. 518. 1981.
- Aquatic Vascular Plants of New England: Part 4. Juncaginaceae, Scheuchzeriaceae, Butomaceae, Hydrocharitaceae. G. E. Crow and C. B. Hellquist. Bull. No. 520. 1982.
- Aquatic Vascular Plants of New England: Part 5. Araceae, Lemnaceae, Xyridaceae, Eriocaulaceae, and Pontederiaceae. C. B. Hellquist and G. E. Crow. Bull. No. 523. 1982.
- Aquatic Vascular Plants of New England: Part 6. Trapaceae, Haloragaceae, Hippuridaceae. G. E. Crow and C. B. Hellquist. Bull. No. 524. 1983.
- Aquatic Vascular Plants of New England: Part 7. Cambombaceae, Nymphaeaceae, Nelumbonaceae, and Ceratophyllaceae. C. B. Hellquist and G. E. Crow. Bull. No. 527.
- Hiker Traffic On and Near the Habitat of Robbins Cinquefoil, an Endangered Plant Species. R. E. Graber and G. E. Crow. Bull. No. 522. 1982.



Aroostook

Piscataquis

Somerset

Penobscot

Grand Isle

Franklin

Orleans

Essex

Coos

Franklin

Washington

Lamoille

Chittenden

Caledonia

Washington

Oxford

Kennebec

Waldo

Hancock

Addison

Orange

Gratton

Carroll

Cumberland

Lincoln

Sagadahoc

Androscoggin

Rutland

Windsor

Belknap

York

Sullivan

Merrimack

Strafford

Bennington

Windham

Rockingham

Cheshire

Hillsboro

Middlesex

Essex

Berkshire

Franklin

Hampshire

Worcester

Suffolk

Hampden

Norfolk

Plymouth

Litchfield

Hartford

Barnstable

Fairfield

New Haven

New London

Kent

Nantucket

Tolland

Dukes

Bristol, MA

Bristol, RI

Middlesex

Newport

Providence

Washington

Windham

