

University of New Hampshire

## University of New Hampshire Scholars' Repository

---

Earth Sciences Scholarship

Earth Sciences

---

2013

### Quantifying and mapping China's crop yield gains from sustainable and unsustainable irrigation water use 1981-2000

Danielle S. Grogan

*University of New Hampshire - Main Campus, [danielle.grogan@unh.edu](mailto:danielle.grogan@unh.edu)*

Fan Zhang

*Xi'an Jiaotong University*

Stanley Glidden

*University of New Hampshire - Main Campus, [stanley.glidden@unh.edu](mailto:stanley.glidden@unh.edu)*

Dominik Wisser

*Universitat Bonn*

Alexander Prusevich

*University of New Hampshire - Main Campus*

*See next page for additional authors*

Follow this and additional works at: [https://scholars.unh.edu/earthsci\\_facpub](https://scholars.unh.edu/earthsci_facpub)

---

#### Recommended Citation

Grogan D, Zhang F, Glidden S, Prusevich A, Lammers R, Wisser D, Li C, Froking S. 2013. Quantifying and mapping China's crop yield gains from sustainable and unsustainable irrigation water use 1981-2000, Global Water Systems Program (GWSP) 2013 - Water in the Anthropocene, Bonn Germany, May 2013.

This Conference Proceeding is brought to you for free and open access by the Earth Sciences at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Earth Sciences Scholarship by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact [Scholarly.Communication@unh.edu](mailto:Scholarly.Communication@unh.edu).

---

**Authors**

Danielle S. Grogan, Fan Zhang, Stanley Glidden, Dominik Wisser, Alexander Prusevich, Changsheng Li, Richard B. Lammers, and Steve Froking

# Quantifying and mapping China's crop yield gains from sustainable and unsustainable irrigation water use 1981-2000

Danielle S. Grogan<sup>1\*</sup>, Fan Zhang<sup>1,2</sup>, Stanley Glidden<sup>1</sup>, Dominik Wissler<sup>3</sup>, Alex Prusevich<sup>1</sup>, Changsheng Li<sup>1</sup>, Richard B. Lammers<sup>1</sup>, Steve Froliking<sup>1</sup>

<sup>1</sup>University of New Hampshire, Durham, NH; <sup>2</sup>Xi'an Jiaotong University, Xi'an, China; <sup>3</sup>Utrecht University, Utrecht, Netherlands

\*Danielle Grogan@wildcats.unh.edu  
Earth Systems Research Center  
University of New Hampshire  
208 Morse Hall, 8 College Road  
Durham, New Hampshire 03824 (USA)



## I. Research Questions


How much unsustainable water is used for irrigation in China (c.2000)?  
How will crop yields change when the unsustainable water runs out?

## II. Background

- Chinese agriculture depends greatly on irrigation water. ~40% of China's cropland is irrigated.
- It has been estimated that ~15% of China's irrigation water comes from unsustainable sources<sup>1</sup>
- Regions that rely on unsustainable irrigation water could face water shortages in the future, and may already be experiencing water stress today<sup>1</sup>.
- Identifying crops, regions, and total crop yields that will be impacted by the eventual loss of unsustainable water can help plan for future water management.


## III. Methods

- Use 2 models: DNDC<sup>2</sup> for irrigated and rainfed crop yields per area, WBM for sustainable water availability and crop water requirements. Input 20 years of climate variability.
- Model two scenarios:
  - Only use sustainable water for irrigation
  - Allow unsustainable water for irrigation
- Scale the MIRCA2000 dataset of irrigated and rainfed cropland areas to reduce irrigation demand to sustainable water supply. Algorithm:
  - $I = \text{MIRCA2000 irrigated areas}$
  - $R = \text{MIRCA2000 rainfed areas}$
  - $W_s = \text{sustainable water as a fraction of irrigation demand}$
  - $I_r = \text{rescaled irrigated areas} = W_s * I$
  - $R_r = \text{rescaled rainfed areas} = R + (1 - W_s) * I$
- Calculate total crop yield for MIRCA2000 areas ( $I, R$ ) and for rescaled areas ( $I_r, R_r$ ). The difference is the crop yield reduction.




## IV. Models

### 1. Water Balance Model



### 2. Denitrification-Decomposition (DNDC) Agro-ecosystem model<sup>2</sup>




### Model Inputs:

Model input type	Water Balance Model (WBM)	DNDC
Climate	MERRA <sup>3</sup>	MERRA <sup>3</sup>
Crop distribution	MIRCA2000 <sup>3</sup>	CAAS
Irrigated areas	MIRCA2000 <sup>3</sup>	all crop/soil/climate conditions simulated with and without irrigation
Cropping Intensity	AQUASTAT (2008) <sup>4</sup> , MIRCA2000 <sup>3</sup>	China statistical yearbook <sup>4</sup>
Crop categories	26 crops (31 including subcrops)	17 individual crops & 28 multi-cropping systems
Soil properties	UNESCO/FAO soil map of the world <sup>5</sup>	Third National Soil Survey
Spatial Resolution	30 arc minute grids	China counties (~30 arc min polygons) Resampled to 30 min grids
Temporal Resolution	Daily Aggregated to monthly & annual totals	Daily Aggregated to annual total


References:  
 1) Wissler D., Froliking S., Douglas E.M., Fekete B.M., Schumann A.H., Vorosmarty C.J. 2010. Blue and green water: The significance of local water resources captured in small reservoirs for crop production. *J. Hydrology*, 384:264-275.  
 2) Li, C., Froliking, S., Froliking, T.A., 1992. A model of nitrous oxide evolution from soil driven by rainfall events. I. Model structure and sensitivity. *J. Geophys. Res.*, 97, 9739-9776.  
 3) Mitchell, T.D., Jones, P.D., 2005. An improved method of constructing a database of monthly climate observations and associated high-resolution grids. *International Journal of Climatology*, 25(6), 693-712.  
 4) Portmann, F.T., Siebert, S., D.P., 2010. MIRCA2000 - Global monthly irrigated and rainfed crop areas around the year 2000: A new high-resolution data set for agricultural and hydrological modeling. *Global Biogeochemical Cycles*, 24, GB1011, doi:10.1029/2008GB003435.  
 5) FAO/UNESCO, 2003. *Digital Soil Map of the World and Derived Soil Properties Version 3.6*. CD ROM.  
 6) Allen, S.E., Pereira, L.S., Baer, G., Smith, M., 1988. Crop evapotranspiration: guidelines for computing crop water requirements, Food and Agricultural Organization of the United Nations (FAO).

## V. Results


### a. Annual irrigation water demand (mm) for an average climate year.




### b. Unsustainable water as a fraction of total irrigation water demand in a dry year (1986).




### c. Annual irrigation water withdrawals from streams & reservoirs, rechargeable groundwater, and unsustainable sources. Annual precipitation is shown as the top.



### d. Annual reduction in irrigated crop yields due to loss of unsustainable water in a dry year (1986). Unit: fraction



### e. Annual reduction in rice (top), wheat (bottom left) and maize (bottom right) crop yields due to loss of unsustainable water in a dry year (1986). Unit: metric tons



Sustainable crop yield from areas equipped for irrigation is 10% (wet year) to 15% (dry year) lower than fully irrigated crop yield.

### Impact on food: a quick calculation

10% of irrigated yield = ~50 million metric ton C

1 metric ton Rice = ~1.3 x 10<sup>6</sup> Calories = one year of food for 2 people (assume 2000 Calories per year)

50 million metric tons x one year of food for 2 people = one year of food for 100 million people (7% China's pop.)