

University of New Hampshire

University of New Hampshire Scholars' Repository

Center for Coastal and Ocean Mapping

Center for Coastal and Ocean Mapping

2001

Martha's Vineyard Survey: Data collected by Bill Schwab and Bill Danforth - USGS

Larry A. Mayer

University of New Hampshire, larry.mayer@unh.edu

Follow this and additional works at: <https://scholars.unh.edu/ccom>



Part of the [Oceanography and Atmospheric Sciences and Meteorology Commons](#)

Recommended Citation

Mayer, Larry A., "Martha's Vineyard Survey: Data collected by Bill Schwab and Bill Danforth - USGS" (2001). *Center for Coastal and Ocean Mapping*. 1149.
<https://scholars.unh.edu/ccom/1149>


This Poster is brought to you for free and open access by the Center for Coastal and Ocean Mapping at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Center for Coastal and Ocean Mapping by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact Scholarly.Communication@unh.edu.

MARTHA'S VINEYARD SURVEY


Data collected by Bill Schwab and Bill Danforth - USGS

The Martha's Vineyard field area has been the subject of several preliminary surveys conducted by USGS and WHOI scientists in support of establishing an observatory node (Feb. 2001) as well as in direct support of the ONR Mine Burial Program (Sept - Nov 2001). Data sets collected include:


- Knudsen 3.5 chirp sonar -- Bill Schwab and Bill Danforth USGS
 - Geopulse Boomer -- Bill Schwab and Bill Danforth USGS
 - DF 1000 towed sidescan sonar (100 and 500 kHz) Feb and Sept/Nov - USGS
 - Submetrix interferometric bathymetry and backscatter (234 kHz) -- USGS
- Also rumored are:
- A WHOI core database
 - Tripod data
 - Edgetech Subscan data (Neal Driscoll, Wayne Spencer)
 - EM and GPR data (Rob Evans)




3-D perspective view of proposed site. Bathymetry is based on NOAA's Coastal Relief Model. Higher resolution Submetrix interferometric bathymetry is still being processed




Interpretation (by Schwab and Danforth) of thickness of sand layer as revealed in Knudsen 3.5 chirp profiles (below). Thickness ranges from 0 to 1.3m




Initial processing of recently collected Submetrix interferometric bathymetry superimposed on NOAA Coastal Relief Model. Bathymetric cross-section is along blue line in eastern survey area. Eastern margin is node site.




The proposed study area as depicted on NOAA chart with Feb 2001 100 kHz Df1000 sidescan survey superimposed. High backscatter is light -- note what appear to be several major sand bodies running perpendicular to shore




3-D perspective view of NOAA bathymetry with Feb Df1000 sidescan data superimposed




Examples of Geopulse boomer crossing of filled channel and Knudsen 3.5 kHz profiler imaging surficial sand layer.




Feb Df1000 sidescan sonar data superimposed on Submetrix interferometric bathymetry. Bathymetric cross-section is along blue line in eastern survey area. Eastern margin is node site.




Df1000 100kHz sidescan data collected in Sept-Nov. 2001. High backscatter is light -- a number of shore parallel features are clear. Survey is 13 km wide (E-W) and about 3 km high (N-S).



3-D perspective view of NOAA bathymetry with Sept-Nov Df1000 sidescan data superimposed



Depth of filled channel imaged by Geopulse boomer. Interpretation by Bill Schwab and Bill Danforth, USGS



Sept-Nov Df1000 sidescan sonar data superimposed on Submetrix interferometric bathymetry. Bathymetric cross-section is along blue line in eastern survey area. Eastern margin is node site.

Feb Df1000 sidescan sonar data superimposed on Sept-Nov Df1000 sidescan sonar data. The large high-backscatter targets appear to have not changed!