

University of New Hampshire

## University of New Hampshire Scholars' Repository

---

Faculty Publications

---

5-11-2021

### Fungal community response to long-term soil warming with potential implications for soil carbon dynamics

Gregory J. Pec

*University of New Hampshire, Durham*

Linda T. A. van Diepen

*University of Wyoming*

Melissa A. Knorr

*University of New Hampshire, Durham*

A. Stuart Grandy

*University of New Hampshire, Durham*

Jerry M. Melillo

*Marine Biological Laboratory, Woods Hole*

Follow this and additional works at: [https://scholars.unh.edu/faculty\\_pubs](https://scholars.unh.edu/faculty_pubs)

See next page for additional authors



Part of the [Biogeochemistry Commons](#)

#### Comments

This is an open access article published by ESA in Ecosphere in 2021, available online: <https://dx.doi.org/10.1002/ecs2.3460>

---

#### Recommended Citation

Pec, G. J., L. T. A. van Diepen, M. Knorr, A. S. Grandy, J. M. Melillo, K. M. DeAngelis, J. L. Blanchard, and S. D. Frey. 2021. Fungal community response to long-term soil warming with potential implications for soil carbon dynamics. *Ecosphere* 12(5):e03460. 10.1002/ecs2.3460

This Article is brought to you for free and open access by University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Faculty Publications by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact [Scholarly.Communication@unh.edu](mailto:Scholarly.Communication@unh.edu).

---

**Authors**

Gregory J. Pec, Linda T. A. van Diepen, Melissa A. Knorr, A. Stuart Grandy, Jerry M. Melillo, Kristen M. DeAngelis, Jeffrey L. Blanchard, and Serita D. Frey

# Fungal community response to long-term soil warming with potential implications for soil carbon dynamics

GREGORY J. PEC<sup>1,†</sup>, LINDA T. A. VAN DIEPEN,<sup>2</sup> MELISSA KNORR,<sup>1</sup> A. STUART GRANDY,<sup>1</sup> JERRY M. MELILLO,<sup>3</sup> KRISTEN M. DEANGELIS,<sup>4</sup> JEFFREY L. BLANCHARD,<sup>5</sup> AND SERITA D. FREY<sup>1</sup>

<sup>1</sup>Department of Natural Resources and the Environment, University of New Hampshire, Durham, New Hampshire 03824 USA

<sup>2</sup>Department of Ecosystem Science and Management, University of Wyoming, Laramie, Wyoming 82071 USA

<sup>3</sup>The Ecosystems Center, Marine Biological Laboratory, Woods Hole, Massachusetts 02543 USA

<sup>4</sup>Department of Microbiology, University of Massachusetts, Amherst, Massachusetts 01003 USA

<sup>5</sup>Biology Department, University of Massachusetts, Amherst, Massachusetts 01003 USA

**Citation:** Pec, G. J., L. T. A. van Diepen, M. Knorr, A. S. Grandy, J. M. Melillo, K. M. DeAngelis, J. L. Blanchard, and S. D. Frey. 2021. Fungal community response to long-term soil warming with potential implications for soil carbon dynamics. *Ecosphere* 12(5):e03460. 10.1002/ecs2.3460

**Abstract.** The direction and magnitude of climate warming effects on ecosystem processes such as carbon cycling remain uncertain. Soil fungi are central to these processes due to their roles as decomposers of soil organic matter, as mycorrhizal symbionts, and as determinants of plant diversity. Yet despite their importance to ecosystem functioning, we lack a clear understanding of the long-term response of soil fungal communities to warming. Toward this goal, we characterized soil fungal communities in two replicated soil warming experiments at the Harvard Forest (Petersham, Massachusetts, USA) which had experienced 5°C above ambient soil temperatures for 5 and 20 yr at the time of sampling. We assessed fungal diversity and community composition by sequencing the ITS2 region of rDNA using Illumina technology, along with soil C concentrations and chemistry. Three main findings emerged: (1) long-, but not short-term warming resulted in compositional shifts in the soil fungal community, particularly in the saprotrophic and unknown components of the community; (2) soil C concentrations and the total C stored in the organic horizon declined in response to both short- (5 yr) and long-term (20 yr) warming; and (3) following long-term warming, shifts in fungal guild relative abundances were associated with substantial changes in soil organic matter chemistry, particularly the relative abundance of lignin. Taken together, our results suggest that shifts with warming in the relative abundance of fungal functional groups and dominant fungal taxa are related to observed losses in total soil C.

**Key words:** ectomycorrhizal fungi; Illumina; organic matter chemistry; saprotrophic fungi; soil carbon.

**Received** 29 May 2020; revised 2 November 2020; accepted 12 November 2020; final version received 28 January 2021. Corresponding Editor: Debra P. C. Peters.

**Copyright:** © 2021 The Authors. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

† **E-mail:** pecg@unk.edu

## INTRODUCTION

Global warming poses major challenges to the health and functioning of forests worldwide (Peters 1990, Van der Putten 2012, Langley and Hungate 2014, Classen et al. 2015), ecosystems that hold a substantial proportion of carbon (C) in both their vegetation and soil pools (Lal 2004,

Bonan 2008, Gill and Finzi 2016). A number of soil warming studies have examined ecosystem-level effects, with an emphasis on soil respiration and N cycling dynamics (Bradford et al. 2008, Carey et al. 2016, Dacal et al. 2019). Long-term (>10 yr) studies have revealed that chronic soil warming results in a cyclic pattern of soil organic matter (SOM) decomposition and carbon dioxide

(CO<sub>2</sub>) loss to the atmosphere, with phases of substantial soil C loss alternating with phases of no detectable loss (Melillo et al. 2002, 2017). Several mechanisms have been proposed to explain this observation including depletion of microbially accessible carbon pools, reductions in microbial biomass, a shift in microbial carbon use efficiency, and changes in microbial community composition (Oechel et al. 2000, Allison and Martiny 2008, Frey et al. 2008, Bradford 2013, Melillo et al. 2017). Of these, the hypothesis that the loss in soil C from long-term warming is driven by changes in microbial community structure is an important one (Schimel and Schaeffer 2012), since warming may selectively impact microbial taxa with specific functional capacities that influence SOM decomposition and, ultimately, rates of soil C cycling.

Soil fungi, integral components of forest ecosystems, mediate key ecosystem processes including C and nutrient cycling (Zifčáková et al. 2016, Fraç et al. 2018). They form diverse and spatially variable communities as mycorrhizal symbionts, saprotrophs, and pathotrophs (Tedersoo et al. 2014). Fungal community responses to soil warming differ among ecosystems (Frey et al. 2008, Geml et al. 2015, Semanova et al. 2015, Treseder et al. 2016); however, a general finding is that the abundance of soil fungi often declines in response to long-term warming (Rinnan et al. 2007, Frey et al. 2008, DeAngelis et al. 2015, Liang et al. 2015) with shifts in total fungal community composition often observed over the same time frame (Deslippe et al. 2012). It is less clear, however, whether certain functional groups (i.e., ectomycorrhizal vs. saprotrophs) are differentially sensitive to shifts in temperature. For example, the response of ectomycorrhizal fungi to warming has varied from positive (Treseder et al. 2016, Solly et al. 2017) to negative (Geml et al. 2015), whereas saprotrophs have been observed to increase in their relative abundance to warming (Treseder et al. 2016, Romero-Olivares et al. 2017). Shifts in abundance of key fungal functional groups (e.g., mycorrhizal fungi vs. saprotrophs; or cellulose vs. lignin-degraders) are expected to significantly impact soil C cycling (Trumbore et al. 1996, Treseder et al. 2016), highlighting the importance of resolving fungal responses to warming.

Here, we build on previous research in a temperate deciduous forest (Frey et al. 2008, DeAngelis et al. 2015, Melillo et al. 2017) to assess changes in the diversity and community composition of soil fungi following soil warming. We not only wanted to investigate overall community changes, but also evaluate how different taxa and key functional groups vary in their sensitivities to warming. Since microbial responses to warming have been previously shown to be associated with changes in soil chemistry (Pold et al. 2016, 2017), a secondary objective was to assess key soil chemical properties following short- vs. long-term warming as well as examine relationships between the fungal community and soil properties.

## METHODS

### *Site description, experimental design, and sample collection*

This research was performed at two adjacent, replicated soil warming experiments at the Harvard Forest Long-term Ecological Research (LTER) site in Petersham, Massachusetts (MA), USA (42°50' N, 72°18' W). The experimental plots were established in an area of forest abandoned from agriculture in the late 1800s (Peterjohn et al. 1994). Secondary forest regrowth is dominated by paper and black birch (*Betula papyrifera* and *lenta*), red maple (*Acer rubrum*), black and red oak (*Quercus velutina* and *rubra*), and American beech (*Fagus grandifolia*). Soils are classified as coarse-loamy Inceptisols with no significant variation in pH of the organic soil horizon ( $-x = 3.9 \pm 0.1$  standard error) among experimental plots. Mean annual temperature at Harvard Forest ranges from 19°C to -5°C, with precipitation (annual average ~120 cm) distributed evenly throughout the year. One of the longest running soil warming experiments in the world, the Prospect Hill Soil Warming Study, was started in 1991 and consists of six 6 × 6 m control and warmed plots (Melillo et al. 2002). A second experiment, the Soil Warming × Nitrogen Addition Study (SWaN), was established in 2006 adjacent to the Prospect Hill experiment and has six 3 × 3 m control and warmed plots (Contosta et al. 2015). Although there is a nitrogen (N) fertility component at this second experiment, only control and warmed plots were

evaluated here. Soils in the warmed plots at both sites are heated continuously to 5°C above ambient temperatures observed in the control plots by the use of resistance heating cables buried 10 cm deep and spaced 20 cm apart (Melillo et al. 2017).

Soil samples were collected from the organic horizon of four replicate control and warmed plots following five (SWaN) and 20 (Prospect Hill) years of warming for a total of 16 samples (2 sites × 2 warming treatments × 4 replicates). We focused on the organic horizon because previous work at this site has shown that most of the fungal biomass (96%) and microbial activity based on substrate-induced respiration (88%) are located in this horizon (Frey et al. 2008), with a significant warming-induced decline in the soil C pool (−31%) being observed for this horizon as well (Melillo et al. 2017). Samples were collected from experimental plots as intact 20 × 20 cm blocks to the depth of the mineral soil. Samples were sieved (<2 mm), and roots, rocks, and coarse woody debris (>2 mm) were removed. Subsamples were flash-frozen using dry-ice immediately in the field and stored at −80°C until analysis for fungal community characterization. Remaining soils were stored at 4°C until analysis for soil chemical properties.

#### *Fungal community characterization*

To characterize fungal diversity and community composition, genomic DNA was isolated on subsamples (0.25 g) using the PowerSoil DNA Isolation Kit (MoBio Laboratories, Carlsbad, California, USA) following manufacturer protocols. Library preparation and sequencing were done at the Department of Energy Joint Genome Institute (Walnut Creek, California, USA). Briefly, polymerase chain reaction (PCR) was performed to amplify the internal transcribed spacer (ITS) 2 region of nuclear rDNA using primers fITS7 (GTGARTCATCGAATCTTTG; Ihrmark et al. 2012) and ITS4 (TCCTCCGCTTATTGATATGC; White et al. 1990). PCRs contained 13 µL of PCR-grade water, 10 µL of Five Prime Hot Master Mix (5 PRIME, Gaithersburg, Maryland, USA), 0.5 µL of each forward and reverse primer (10 µmol/L final concentration), and 1 µL of template DNA. Thermocycling conditions were as follows: one cycle at 94°C for 3 min; then 35 cycles at 94°C for 45 s, 50°C for 60 s, and 72°C for 90 s; and ending

with one cycle of 72°C at 10 min. Successful amplification was confirmed using gel electrophoresis (1.5% agarose gel). PCR products were cleaned using the AxyPrep MAG PCR clean-up kit (Coming, Tewksbury, Massachusetts, USA), quantified fluorescently on a Qubit fluorometer (ThermoFisher Scientific, Grand Island, New York, USA), and pooled into equimolar concentrations. Amplicon library sequencing was performed on an Illumina MiSeq v3 (2 × 300 bp). Raw sequences are available at the NCBI database under BioProject number PRJNA523054.

#### *Bioinformatic analysis*

Illumina MiSeq data were first demultiplexed and quality-filtered using Trimmomatic v0.36 (Bolger et al. 2014), removing Illumina adapters, sequences with <200 bp, and bases with minimum quality scores <2. All remaining forward and reverse reads were merged with the fastq-join method using the join\_paired\_ends.py script in Qiime v1.9 (Caporaso et al. 2010) followed by ITS2 region extraction using ITSx v1.0.11 (Bengtsson-Palme et al. 2013). Sequences were clustered into operational taxonomic units (OTUs) at 97% sequence similarity while simultaneously removing chimeric sequences using the UPARSE-OTU algorithm (Edgar 2013) with the cluster\_otus command in USEARCH (v9.2.64; Edgar 2010). Additionally, we excluded all global singletons to reduce artificially inflating richness due to sequencing error. Sequence data were first rarefied to a depth of 257,520 sequences per sample to account for uneven sequence depths across samples. OTUs were taxonomically identified by searching representative sequences of each OTU against the UNITE fungal ITS database (Nilsson et al. 2019) using the BLAST option in the assign\_taxonomy.py script in Qiime v1.9. Non-fungal OTUs were excluded from further analyses. Fungal OTUs were initially assigned to three groups (ectomycorrhizal, saprotrophic, and pathotrophic) based on their genus affiliation, trophic mode, and functional guild as previously described (Branco et al. 2013, Tedersoo and Smith 2013, Tedersoo et al. 2014) and using the FUNGuild database (Nguyen et al. 2016). OTUs were placed into groupings only if assignments were deemed as highly probable or probable based on default parameters in the FunGuild

database (<http://stbates.org/guild/app.php>). We further divided ectomycorrhizal fungal OTUs into ectomycorrhizal fungal exploration types (i.e., contact to short distance or medium to long distance) according to (Agerer 2001), while saprotrophic fungal OTUs were also further divided into yeasts, filamentous saprotrophs, and white rot fungi according to the FUNGuild database (Nguyen et al. 2016). Here, white rot fungi were included as a separate group because of our interest in lignin-degraders. Fungal OTUs were designated as “taxonomy insufficient” if they were assigned a genus affiliation but their functional grouping was uncertain. Fungal OTUs that either had no known identification in available databases or were identified only to the kingdom level were designated as “unknown taxonomy.”

### Soil chemical properties

Soil bulk density, determined for each sample, was used to calculate volumetric soil C stocks ( $\text{g C/m}^3$ ). Soil samples were analyzed for total organic C and N on air-dried finely ground samples by dry combustion on a Perkin Elmer 2400 Series II CHN elemental analyzer (Waltham, Massachusetts, USA). Organic matter chemistry was assessed by determining the relative abundances of chemical classes using pyrolysis-GC/MS (Wickings et al. 2011). In brief, samples were pyrolyzed at  $600^\circ\text{C}$ , products transferred to a GC where compounds were separated and transferred to an ion trap mass spectrometer, ionized, detected via electron multiplier, and identified using a compound library from the National Institute of Standards and Technology (NIST; Grandy et al. 2007, 2009).

### Statistical analyses

All statistical analyses were carried out using R v3.3.2 (R Core Team 2019). Due to the close physical proximity of the experimental plots for the two warming experiments (Prospect Hill and SWaN), all control plots were grouped together for downstream analyses as no significant differences were detected across control plots for any measured soil property, including fungal community structure (Appendix S1: Table S1). We ran linear models to examine whether there were differences in measured soil chemical properties in response to short- (5 yr) or long-term (20 yr)

warming. Generalized linear models with a Poisson distribution were run to examine how fungal OTU richness responded to warming, while linear models were run to examine whether there were differences in the relative abundance of soil fungal taxa or functional groups to warming. Post hoc pairwise comparisons were performed using the `glht` function in the package `multcomp` (Hothorn et al. 2008). All model assumptions were checked by visual inspection of residual diagnostic plots (Zuur et al. 2009).

We used multivariate generalized linear models (M-GLM) to test whether fungal OTU relative abundance distributions were affected by warming with the `manyglm` function in the `mvabund` package (Wang et al. 2012), using a negative binomial model, likelihood ratio statistic, and  $P$  values assigned following 9999 iterations via probability integral transform-trap sampling to account for correlation in testing using the `anova` function. We used this approach instead of a distance-based metric such as `perMANOVA`, as `mvabund` directly accounts for the strong mean–variance relationships present in most multivariate community relative abundance analyses (Warton et al. 2015). To visualize differences in soil fungal communities across warming treatments, we used non-metric multidimensional scaling (NMDS) ordination based on Bray–Curtis dissimilarity using the `metaMDS` function in the `vegan` package (Oksanen et al. 2014). To identify fungal OTUs common or representative of either control or warmed plots, we used the multinomial species classification method in the `vegan` package (Oksanen et al. 2014) with supermajority rule and all other parameters set as default (i.e., CLAM test) to statistically classify fungal OTUs into the following categories: fungi shared across control and warmed plots, fungi preferentially associated with either control or warmed plots, and fungi that were too rare (based on low relative abundance) to be assigned association specificity (Chazdon et al. 2011).

Partial least-square regression (Wold et al. 2001) was used to identify covariates (soil chemistry) most strongly correlated with fungal guild richness and relative abundance using the `pls()` function in the `pls` package (Mevik and Wehrens 2007). Models were refined to the most important predictor variable(s) and analyzed independently using linear regression. All model

assumptions were checked by visual inspection of residual diagnostic plots (Zuur et al. 2009).

## RESULTS

### *Fungal diversity and community composition*

In total, 6,988,799 sequences were obtained across all samples. After quality filtering, 6,165,723 sequences (88%) representing 2552 fungal OTUs were assessed for taxonomic affiliation. At the phylum level, there were 1094 OTUs (43%) assigned to Basidiomycota, 498 OTUs (19%) assigned to Ascomycota, and 107 OTUs (4%) assigned to basal clades, while Glomeromycota made up <1% of identified sequences (Appendix S1: Table S2). Since ITS primers are biased against glomeromycetes (arbuscular mycorrhizal fungi; Stockinger et al. 2010), this group was not included in subsequent analyses. At the functional group level, there were 732 OTUs assigned to ectomycorrhizal fungi (28%), 339 OTUs assigned to saprotrophic fungi (13%), and 36 OTUs assigned to pathotrophic fungi (1%; Table 1). In addition, there were 843 fungal OTUs (33%) assigned as unknown taxonomy, with most not being assigned to a phylum. A final group of 593 (23%) OTUs were assigned a taxonomic affiliation; however, there was insufficient ecological information with which to reliably assign them to a functional group.

Overall, long- (20 yr) but not short-term (5 yr) warming resulted in significantly different soil fungal communities compared to the control, and long-term plots were also significantly different from short-term warming plots (Fig. 1; Appendix S1: Table S3). Fungal OTU richness tended to increase with warming compared to control plots; however, this increase was only significant for saprotrophic fungi (Table 2). Specifically, the richness of filamentous saprotrophs increased from 61 OTUs in the control treatment to 80 OTUs following 20 yr of warming. In contrast, saprotrophs declined in relative abundance from 33.7% in control plots to 11.6% in plots warmed for 20 yr. This decline was largely due to a significant negative response to warming of filamentous saprotrophs, which made up the majority of saprotrophic fungi (~74%). As the relative abundance of saprotrophic taxa declined with long-term warming, that of unknown fungi significantly increased,

Table 1. Relative sequence abundance and OTU richness of morphological and functional fungal groups.

Morphological/functional group	Relative sequence abundance (%)	Number of OTUs
<b>Symbiotrophs</b>	<b>29.1 (66.4)</b>	<b>741 (66)</b>
Ectomycorrhizal fungi	28.7 (65.5)	732 (65)
Contact to short types	17.9 (40.9)	459 (41)
Medium to long types	2.9 (6.6)	74 (6)
Endophytic fungi	0.4 (0.9)	9 (0.8)
<b>Saprotrophs</b>	<b>13.3 (30.3)</b>	<b>339 (30)</b>
Filamentous saprotrophs	9.8 (22.3)	250 (22)
White rot fungi	1.7 (3.8)	43 (4)
Yeasts	1.4 (3.3)	37 (3)
Brown rot fungi	0.4 (0.9)	9 (0.8)
<b>Pathotrophs</b>	<b>1.4 (3.3)</b>	<b>36 (3)</b>
Plant pathogens	0.6 (1.5)	16 (1)
Animal pathogens	0.6 (1.5)	15 (1)
Mycoparasites	0.2 (0.3)	5 (0.5)
<b>Total assigned</b>	<b>43.8 (100)</b>	<b>1116 (100)</b>
<b>Total unassigned</b>	<b>56.2</b>	<b>1436</b>
Taxonomy insufficient†	23.2	593
Unknown taxonomy‡	33.0	843

*Notes:* Values represent the total of all OTUs across experimental treatments. Values in parentheses are the percentages represented by each category based on total assigned OTUs. OTUs that could not be identified to the family level were assigned a function of unassigned. General categories are bolded for clarity.

† Represents fungal OTUs that were assigned a taxonomic affiliation but for which their functional guild was uncertain or unknown.

‡ Includes OTUs that were identified to the kingdom level (i.e., fungi), but which have no known taxonomic identification in available databases.

going from 28.7% in the control treatment to 41.5% in warmed soils. Ectomycorrhizal and pathotrophic fungal communities did not differ compositionally or in their relative abundances in control vs. warmed plots (Fig. 1, Table 2). Although fungal biomass was not measured as part of this study, previous research at our site has shown that fungal biomass is not significantly affected by short-term warming (<5 yr; Contosta et al. 2015), while fungal biomass tends to decline in response to warming after 12–20 yr (Frey et al. 2008, DeAngelis et al. 2015).

Since most significant differences were observed in response to long-term warming, we further investigated whether there were representative and/or unique taxa in soils warmed for 20 yr. Overall, about a third of fungal OTUs were present in both control and warmed (20 yr) soils (Appendix S1: Fig. S1a), whereas about half of all OTUs were found primarily in either one or the




Fig. 1. Non-metric multidimensional scaling (NMDS) ordinations depicting overall fungal community composition (A) and the ectomycorrhizal (B), saprotrophic (C), and unknown (D) components of the fungal community in response soil warming at Harvard Forest.

Table 2. Relative sequence abundance and OTU richness of soil fungi in response to short- (5 yr) vs. long-term (20 yr) warming at Harvard Forest.

Morphological/functional group	Relative abundance (%)			Number of OTUs		
	Control	Short-term warming (5 yr)	Long-term warming (20 yr)	Control	Short-term warming (5 yr)	Long-term warming (20 yr)
Overall				620 ± 29	648 ± 38	673 ± 63
Ectomycorrhizal fungi	23.1 ± 2.5	26.8 ± 1.8	27.1 ± 3.7	149 ± 7	151 ± 8	155 ± 18
Contact to short types	17.7 ± 2.1	20.1 ± 2.3	20.8 ± 4.7	91 ± 4	94 ± 7	100 ± 12
Medium to long types	1.3 ± 0.7	1.1 ± 0.6	1.7 ± 0.6	12 ± 1	12 ± 1	11 ± 3
Saprotrophic fungi	<b>32.7 ± 3.4<sup>a</sup></b>	<b>30.1 ± 5.1<sup>a</sup></b>	<b>11.6 ± 4.1<sup>b</sup></b>	<b>66 ± 4<sup>a</sup></b>	<b>67 ± 8<sup>a</sup></b>	<b>87 ± 3<sup>b</sup></b>
Filamentous saprotrophs	<b>28.5 ± 3.5<sup>a</sup></b>	<b>27.1 ± 5.0<sup>a</sup></b>	<b>9.5 ± 4.1<sup>b</sup></b>	<b>61 ± 4<sup>a</sup></b>	<b>60 ± 7<sup>a</sup></b>	<b>80 ± 3<sup>b</sup></b>
White rot	2.5 ± 2.3	1.8 ± 1.4	1.3 ± 0.6	8 ± 1	8 ± 1	8 ± 1
Yeasts	1.7 ± 0.7	1.2 ± 0.4	0.8 ± 0.2	7 ± 1	8 ± 1	7 ± 1
Pathotrophic fungi	0.4 ± 0.1	0.3 ± 0.1	0.7 ± 0.4	10 ± 1	10 ± 1	10 ± 1
Taxonomy insufficient†	<b>15.1 ± 2.5<sup>a</sup></b>	<b>11.7 ± 1.1<sup>b</sup></b>	<b>19.1 ± 3.4<sup>a</sup></b>	122 ± 7	133 ± 3	126 ± 7
Unknown taxonomy‡	<b>28.7 ± 3.8<sup>a</sup></b>	<b>31.1 ± 3.9<sup>a</sup></b>	<b>41.5 ± 4.9<sup>b</sup></b>	169 ± 9	177 ± 15	190 ± 21

Note: Values represent means and standard errors, with different lowercase superscript letters in bold indicating significant difference ( $P < 0.05$ ).

† Represents fungal OTUs that were assigned a taxonomic affiliation but for which their functional guild was uncertain or unknown.

‡ Includes OTUs that were identified to the kingdom level (i.e., fungi), but which have no known taxonomic identification in available databases.

other treatment, with an equal proportion observed to be unique in control (340 OTUs) compared to warmed (345 OTUs) soils. In terms of specific functional groups, a third of ectomycorrhizal fungal OTUs were present in both

treatments (Appendix S1: Fig. S1b), while saprotrophic fungi and unknown fungi were found primarily in warmed (saprotrophic, 31%, 65 OTUs; unknown, 29%, 140 OTUs) compared to control soils (saprotrophic, 15%, 31 OTUs;



unknown, 18%, 87 OTUs; Appendix S1: Fig. S1c, d). In particular, the most abundant taxa (>1% relative abundance) differed between control and warmed plots (Table 3). The dominant fungal taxa observed in control plots declined in relative abundance with long-term warming (42% vs. 10%), while the dominant taxa observed in warmed soils were found in much lower relative abundance in control plots (31% vs. 4%). Specifically, the saprotroph *Megacollybia* decreased from 19% in control plots to 5% in warmed soils, while *Oidiodendron*, a dark septate endophyte, increased from <1% in control plots to 8% with warming. Of particular interest, half of the most abundant taxa in control and warmed plots were of unknown taxonomy, representing ~15% relative abundance combined. The most abundant unknown taxa in control plots (OTUs 57, 394, 666, 751, 1034) declined from 15% to 3.6% with

warming, whereas the most abundant unknown taxa in warmed soils (OTUs 103, 201, 783, 1079, 1273, 1333) increased by sixfold from control plots (Table 3).

### Soil chemical properties

Warming resulted in a significant decline in soil C concentrations ( $F = 3.72$ ,  $P = 0.04$ ) and in the total C stored in the organic horizon ( $F = 5.46$ ,  $P = 0.01$ ) over short- (5 yr) and long-term (20 yr) warming (Table 4). Soil organic matter chemistry was also altered, but only following long-term (20 yr) warming. More specifically, two decades of warming induced a significant reduction (–37%) in the relative abundance of lignin. In contrast, the relative abundance of polysaccharides and unknown origin chemical classes were significantly higher over the same warming period.

Table 3. Shifts in the relative abundance of the most abundant soil fungal taxa found in response to long-term (20 yr) warming.

Taxon	Fungal functional group	Relative abundance (%)	
		Control	Warmed (20 yr)
Most abundant in control plots			
<i>Megacollybia</i>	Saprotroph	19.5 ± 5.2	5.0 ± 1.3
<i>Fungi_751</i>	Unknown taxonomy†	8.4 ± 1.7	2.3 ± 0.8
<i>Trechispora_14</i>	Saprotrophic	2.6 ± 0.8	0.9 ± 0.1
<i>Fungi_394</i>	Unknown taxonomy	2.5 ± 0.6	1.0 ± 0.3
<i>Russula_122</i>	Ectomycorrhizal	1.8 ± 0.5	0.4 ± 0.1
<i>Fungi_57</i>	Unknown taxonomy	1.6 ± 0.4	0.2 ± 0.1
<i>Russula_336</i>	Ectomycorrhizal	1.5 ± 0.4	0.01 ± 0.01
<i>Fungi_666</i>	Unknown taxonomy	1.4 ± 0.4	0.2 ± 0.1
<i>Fungi_1034</i>	Unknown taxonomy	1.3 ± 0.3	0.0 ± 0.0
<i>Herpotrichiellaceae_5</i>	Taxonomy insufficient‡	1.2 ± 0.3	0.3 ± 0.1
Total		41.8%	10.3%
Most abundant in warmed plots			
<i>Oidiodendron</i>	Dark septate endophyte	0.01 ± 0.01	8.1 ± 2.1
<i>Russula_134</i>	Ectomycorrhizal	1.3 ± 0.2	4.3 ± 1.5
<i>Fungi_783</i>	Unknown taxonomy	1.0 ± 0.1	3.5 ± 1.2
<i>Fungi_103</i>	Unknown taxonomy	0.7 ± 0.1	3.2 ± 1.2
<i>Russula_20</i>	Ectomycorrhizal	0.01 ± 0.01	3.1 ± 1.1
<i>Fungi_1273</i>	Unknown taxonomy	0.01 ± 0.01	2.3 ± 0.9
<i>Fungi_1079</i>	Unknown taxonomy	0.01 ± 0.01	1.9 ± 0.9
<i>Fungi_201</i>	Unknown taxonomy	0.4 ± 0.1	1.8 ± 0.8
<i>Fungi_1333</i>	Unknown taxonomy	0.3 ± 0.1	1.7 ± 0.8
<i>Russulaceae_75</i>	Ectomycorrhizal	0.01 ± 0.01	1.4 ± 0.8
Total		3.2%	31.3%

† Includes OTUs that were identified to the kingdom level (i.e., fungi), but which have no known taxonomic identification in available databases.

‡ Represents fungal OTUs that were assigned a taxonomic affiliation but for which their functional guild was uncertain or unknown.

Table 4. Response of key soil chemical properties to short- (5 yr) vs. long-term (20yr) warming at Harvard Forest.

Soil property	Control ( <i>n</i> = 8)	Short-term warming (5 yr; <i>n</i> = 4)	Long-term warming (20 yr; <i>n</i> = 4)
Total organic C (%)	<b>32.9 ± 2.5<sup>a</sup></b>	<b>25.9 ± 3.5<sup>b</sup></b>	<b>23.2 ± 2.1<sup>b</sup></b>
Soil C stock (g C/m <sup>2</sup> )	<b>2939 ± 252<sup>a</sup></b>	<b>2324 ± 284<sup>a</sup></b>	<b>1765 ± 168<sup>b</sup></b>
C:N ratio	24.9 ± 0.4	25.1 ± 0.6	24.0 ± 0.6
SOM chemistry (% relative abundance)			
Lignin	<b>34.9 ± 3.8<sup>a</sup></b>	<b>29.9 ± 4.3<sup>a</sup></b>	<b>22.1 ± 3.1<sup>b</sup></b>
Lipids	12.9 ± 2.6	11.7 ± 2.4	13.4 ± 2.2
N-bearing compounds	4.4 ± 0.3	5.1 ± 0.8	4.6 ± 0.1
Phenols	4.8 ± 1.2	10.0 ± 4.6	4.5 ± 3.1
Polysaccharides	<b>16.3 ± 1.4<sup>a</sup></b>	<b>17.1 ± 1.9<sup>a</sup></b>	<b>23.0 ± 1.1<sup>b</sup></b>
Proteins	3.6 ± 0.2	4.2 ± 0.7	4.1 ± 0.6
Unknown origin	<b>22.8 ± 1.9<sup>a</sup></b>	<b>21.7 ± 4.5<sup>a</sup></b>	<b>28.0 ± 1.3<sup>b</sup></b>

Note: Values represent means and standard errors, with different lowercase superscript letters in bold indicating significant differences ( $P < 0.05$ ).

### Relationships between fungi and soil chemistry

Overall, there were no significant soil chemical predictors of fungal guild richness. However, the relative abundance of lignin was found to be significantly correlated to fungal guild relative abundance (Fig. 2). The relative abundance of saprotrophic fungi as a whole and the most abundant saprotroph, *Megacollybia*, were positively correlated with lignin (saprotrophic fungi,  $F = 15.95$ ,  $P = 0.001$ ; *Megacollybia*,  $F = 13.92$ ,  $P = 0.002$ ). The relative abundances of ectomycorrhizal and unknown fungi were negatively correlated with lignin (ectomycorrhizal:  $F = 9.26$ ,  $P = 0.009$ ; unknown:  $F = 3.49$ ,  $P = 0.05$ ). In contrast, there were no significant correlations with pathotrophic relative abundance.

## DISCUSSION

We assessed soil fungal community diversity and composition, soil C stocks, and key soil chemical properties following short- and long-term warming. Three main findings emerged: (1) long-, but not short-term warming resulted in compositional shifts in the soil fungal community; (2) soil C concentrations in the organic horizon significantly declined in response to short- (5 yr) and long-term (20 yr) warming; and (3) following long-term warming, shifts in fungal guild relative abundances were associated with substantial changes in soil organic matter chemistry, particularly a decline in the relative abundance of lignin.

We found that the composition of the overall soil fungal community was significantly different in warmed vs. control plots, but only following long-term (20 yr) warming. This community shift appears to be attributable to certain fungal functional guilds being differentially sensitive to warming. For example, saprotrophic fungi have been shown to generally increase in relative abundance with warming (Treseder et al. 2016, Romero-Olivares et al. 2017), while the responses of ectomycorrhizal fungi are varied, being dependent on their hosts' growth and photosynthetic rates (Clemmensen et al. 2006, Fernandez et al. 2017). Our contrasting findings of decreased relative abundance of saprotrophic fungi with soil warming are potentially due to (1) competition with ectomycorrhizal fungi (Averill et al. 2014) as supported by an increased ratio of ectomycorrhizal to saprotrophic fungi in long-term warmed plots ( $2.33 \pm 0.2$ ) compared to control plots ( $0.70 \pm 0.5$ ), and/or (2) lower C concentrations and lignin relative abundance. There were shifts in the most abundant soil fungal taxa found in response to 20 yr of soil warming. For example, *Megacollybia*, a common saprotroph in hardwood and mixed forests of northeastern North America, is a highly competitive basidiomycete shown to dominate decomposition (Boddy and Watkinson 1995). It is a white rot fungus with lignin-degrading capability (Worrall et al. 1997). In our study, the relative abundance of *Megacollybia* was significantly correlated with lignin, declining threefold in




Fig. 2. Correlations between soil lignin and fungal functional group relative abundances.

warmed vs. control plots (Fig. 2D). We hypothesize that as the relative abundance of lignin declines in response to warming, so does that of *Megacollybia*. While the relative abundance of *Megacollybia* declined, that of *Oidiodendron*, a dark septate endophyte with a saprotrophic lifestyle (Nilsson 1973) increased in response to warming. *Oidiodendron* is capable of producing a range of enzymes necessary for cellulose and lignin decomposition (Haselwandter et al. 1990, Rice and Currah 2005) and has been observed previously to respond positively to warming (Solly et al. 2017).

Surprisingly, we found ectomycorrhizal fungi to be invariant to soil warming. Though global warming can alter the composition of ectomycorrhizal fungi by pushing them and/or their host plants outside their natural range of physiological tolerances (Kipfer et al. 2010, Fernandez et al. 2017), several other studies examining ectomycorrhizal response to increased temperatures have found small and/or absent effects (Parrent et al. 2006, Tu et al. 2015, Fernandez et al. 2017, Mucha et al. 2018). For example, in the boreal zone,

ectomycorrhizal colonized tree species exposed to warming have shown a reduction in growth and photosynthetic rate (Reich et al. 2015, Fernandez et al. 2017), while warming of tree species in temperate zones has not resulted in similar reductions (Reich and Oleksyn 2008, Wheeler et al. 2017). This suggests that in regions where host plant performance declines with increased temperature, trees may allocate less C to ectomycorrhizal fungi, resulting in declines in ectomycorrhizal fungal richness and relative abundance.

Fungal taxa of unknown identity showed a strong positive response to long-term warming, significantly increasing in their richness and relative abundance compared to the control treatment. Unknown fungi also showed a significant negative correlation with soil chemistry, where lignin relative abundance correlated with lower relative abundance of unknown taxa. In addition, a number of taxa with unknown identity were unique indicator species of either the control treatment or plots warmed for 20 yr. Identifying and understanding the specific

contributions of the dominant unknown fungal taxa described here are of high priority as they show a strong response to long-term warming and may play a critical role in soil C dynamics. The proportion of unidentified taxa (33% of OTUs) is consistent with other fungal community analyses employing high-throughput sequencing (Lücking and Hawksworth 2018 and references therein). The poor characterization of fungi is due to a lack of reference sequences and mis-annotated or misidentified reference sequences (Nilsson et al. 2011, 2016, Lücking and Hawksworth 2018). Initiatives such as the UNITE database for the molecular identification of fungi (Koljal et al. 2013) and alternative classification and identification methods (Hibbett et al. 2016, Irannia and Chen 2016, Abarenkov et al. 2018, Hongsanan et al. 2018) have been proposed to mitigate these issues.

Though this current study focused on fungi, the importance of bacteria in our experimental system at Harvard Forest has been separately studied. Interestingly, bacteria showed a similar response to warming, where only the long-term warming (20 yr) altered bacterial community composition (DeAngelis et al. 2015). The fungal-to-bacterial ratio of the whole microbial community was also reduced in the warming plots caused by a reduction in fungi, not bacteria (DeAngelis et al. 2015), and warming also led to an enrichment of putative lignin-using bacterial taxa (Pold et al. 2015). This suggests that the decline in relative abundance of lignin may be facilitated by microbial community shifts toward bacterial taxa with the capacity to enzymatically access and use lignin, leading to increased competition between bacteria- and fungal lignin-degrading taxa (e.g., *Megacollybia*).

In summary, long-term warming induced changes in fungal diversity and community composition, as well as the relative abundance of fungal groups with different trophic strategies. Our results also highlight that long-term warming has rather large net effects on soil C storage and soil organic matter chemistry, particularly the relative abundance of lignin. Potential reductions in soil C storage and organic matter quality and quantity (Kammer et al. 2009, Harte et al. 2015) could, in turn, effect future ecosystem C fluxes and storage. Taken together, we propose that shifts with warming

in the relative abundance of key fungal functional groups and dominant fungal taxa, along with lower total fungal biomass (Frey et al. 2008, DeAngelis et al. 2015), are related to losses in soil C concentrations and the total C stored in the organic horizon.

## ACKNOWLEDGMENTS

Thanks to Brian Godbois and Chris Cook at the University of New Hampshire for soil sample collection. Sequencing was done at the Joint Genome Institute as part of a Community Sequencing Program (under contract no. DE-AC02-05CH11231 CSP-1058) award. Thanks to Toni Westbrook at UNH for assistance with bioinformatics. The soil warming plots at Harvard Forest are maintained through support from the NSF Long-term Ecological Research (DEB 1237491) and Long-term Research in Environmental Biology (DEB 1456528) Programs.

## LITERATURE CITED

- Abarenkov, K., P. Somervuo, R. H. Nilsson, P. M. Kirk, T. Huotari, N. Abrego, and O. Ovaskainen. 2018. Protax-fungi: a web-based tool for probabilistic taxonomic placement of fungal internal transcribed spacer sequences. *New Phytologist* 220:517–525.
- Agerer, R. 2001. Exploration types of ectomycorrhizae. *Mycorrhiza* 11:107–114.
- Allison, S. D., and J. B. H. Martiny. 2008. Resistance, resilience, and redundancy in microbial communities. *Proceedings of the National Academy of Sciences USA* 105:11512–11519.
- Averill, C., B. L. Turner, and A. C. Finzi. 2014. Mycorrhiza-mediated competition between plants and decomposers drives soil carbon storage. *Nature* 505:543–545.
- Bengtsson-Palme, J., et al. 2013. Improved software detection and extraction of ITS1 and ITS2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. *Methods in Ecology and Evolution* 4:914–919.
- Boddy, L., and S. C. Watkinson. 1995. Wood decomposition, higher fungi, and their role in nutrient redistribution. *Canadian Journal of Botany* 73:1377–1383.
- Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120.
- Bonan, G. B. 2008. Forests and climate change: forcings, feedbacks, and the climate benefits of forests. *Science* 320:1444–1449.

- Bradford, M. A. 2013. Thermal adaptation of decomposer communities in warming soils. *Frontiers in Microbiology* 4:333.
- Bradford, M. A., C. A. Davies, S. D. Frey, T. R. Madrox, J. M. Melillo, J. E. Mohan, J. F. Reynolds, K. K. Treseder, and M. D. Wallenstein. 2008. Thermal adaptation of soil microbial respiration to elevated temperature. *Ecology Letters* 11: 1316–1327.
- Branco, S., T. D. Bruns, and I. Singleton. 2013. Fungi at a small scale: spatial zonation of fungal assemblages around single trees. *PLOS ONE* 8:e78295.
- Caporaso, J. G., et al. 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods* 7:335–336.
- Carey, J. C., et al. 2016. Temperature response of soil respiration largely unaltered with experimental warming. *Proceedings of the National Academy of Sciences USA* 113:13797–13802.
- Chazdon, R., A. Chao, R. Colwell, S.-Y. Lin, N. Norden, S. Letcher, D. B. Clark, B. Finegan, and P. Arroyo. 2011. A novel statistical method for classifying habitat generalists and specialists. *Ecology* 92:1332–1343.
- Classen, A. T., M. K. Sundqvist, J. A. Henning, G. S. Newman, J. A. M. Moore, M. A. Cregger, L. C. Moorhead, and C. M. Patterson. 2015. Direct and indirect effects of climate change on soil microbial and soil microbial-plant interactions: What lies ahead? *Ecosphere* 6:art130.
- Clemmensen, K. E., A. Michelsen, S. Jonasson, and G. R. Shaver. 2006. Increased ectomycorrhizal fungal abundance after long-term fertilization and warming of two arctic tundra ecosystems. *New Phytologist* 171:391–404.
- Contosta, A. R., S. D. Frey, and A. B. Cooper. 2015. Soil microbial communities vary as much over time as with chronic warming and nitrogen additions. *Soil Biology and Biochemistry* 88:19–24.
- Dacal, M., M. A. Bradford, C. Plaza, F. T. Maestre, and P. García-Palacios. 2019. Soil microbial respiration adapts to ambient temperature in global drylands. *Nature Ecology & Evolution* 3:232.
- DeAngelis, K. M., G. Pold, B. D. Topçuoğlu, L. T. A. van Diepen, R. M. Varney, J. L. Blanchard, J. Melillo, and S. D. Frey. 2015. Long-term forest soil warming alters microbial communities in temperate forest soils. *Frontiers in Microbiology* 6:104.
- Deslippe, J. R., M. Hartmann, S. W. Simard, and W. W. Mohn. 2012. Long-term warming alters the composition of Arctic soil microbial communities. *FEMS Microbiology Ecology* 82:303–315.
- Edgar, R. C. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26:2460–2461.
- Edgar, R. C. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. *Nature Methods* 10:996–998.
- Fernandez, C. W., N. H. Nguyen, A. Stefanski, Y. Han, S. E. Hobbie, R. A. Montgomery, P. B. Reich, and P. G. Kennedy. 2017. Ectomycorrhizal fungal response to warming is linked to poor host performance at the boreal-temperate ecotone. *Global Change Biology* 23:1598–1609.
- Fraç, M., S. E. Hannula, M. Bełka, and M. Jędrzycka. 2018. Fungal biodiversity and their role in soil health. *Frontiers in Microbiology* 9:707.
- Frey, S. D., R. Drijber, H. Smith, and J. Melillo. 2008. Microbial biomass, functional capacity, and community structure after 12 years of soil warming. *Soil Biology and Biochemistry* 40:2904–2907.
- Geml, J., L. N. Morgado, T. A. Semenova, J. M. Welker, M. D. Walker, and E. Smets. 2015. Long-term warming alters richness and composition of taxonomic and functional groups of arctic fungi. *FEMS Microbiology Ecology* 91:fiv095.
- Gill, A. L., and A. C. Finzi. 2016. Belowground carbon flux links biogeochemical cycles and resource-use efficiency at the global scale. *Ecology Letters* 19:1419–1428.
- Grandy, A. S., J. C. Neff, and M. N. Weintraub. 2007. Carbon structure and enzyme activities in alpine and forest ecosystems. *Soil Biology and Biochemistry* 39:2701–2711.
- Grandy, A. S., M. S. Strickland, C. L. Lauber, M. A. Bradford, and N. Fierer. 2009. The influence of microbial communities, management and soil texture on soil organic matter chemistry. *Geoderma* 150:278–286.
- Harte, J., S. R. Saleska, and C. Levy. 2015. Convergent ecosystem responses to 23-year ambient and manipulated warming link advancing snowmelt and shrub encroachment to transient and long-term climate–soil carbon feedback. *Global Change Biology* 21:2349–2356.
- Haselwandter, K., O. Bobleter, and D. J. Read. 1990. Degradation of <sup>14</sup>C-labelled lignin and dehydropolymer of coniferyl alcohol by ericoid and ectomycorrhizal fungi. *Archives of Microbiology* 153:352–354.
- Hibbett, D., et al. 2016. Sequence-based classification and identification of Fungi. *Mycologia* 108:1049–1068.
- Hongsanan, S., et al. 2018. Can we use environmental DNA as holotypes? *Fungal Diversity* 92:1–30.
- Hothorn, T., F. Bretz, and P. Westfall. 2008. Simultaneous inference in general parametric models. *Biometrical Journal* 50:346–363.

- Ihrmark, K., et al. 2012. New primers to amplify the fungal ITS2 region—evaluation by 454-sequencing of artificial and natural communities. *FEMS Microbiology Ecology* 82:666–677.
- Irannia, Z. B., and T. Chen. 2016. TACO: taxonomic prediction of unknown OTUs through OTU co-abundance networks. *Quantitative Biology* 4:149–158.
- Kammer, A., F. Hagedorn, I. Shevchenko, J. Leifeld, G. Guggenberger, T. Goryacheva, A. Rigling, and P. Moiseev. 2009. Treeline shifts in the Ural mountains affect soil organic matter dynamics. *Global Change Biology* 15:1570–1583.
- Kipfer, T., S. Egli, J. Ghazoul, B. Moser, and T. Wohlgemuth. 2010. Susceptibility of ectomycorrhizal fungi to soil heating. *Fungal Biology* 114:467–472.
- Koljal, U., et al. 2013. Towards a unified paradigm for sequence-based identification of fungi. *Molecular Ecology* 22:5271–5277.
- Lal, R. 2004. Soil carbon sequestration to mitigate climate change. *Geoderma* 123:1–22.
- Langley, J. A., and B. A. Hungate. 2014. Plant community feedbacks and long-term ecosystem responses to multi-factored global change. *AoB Plants* 6: plu035.
- Liang, C., J. Gutknecht, and T. Balsler. 2015. Microbial lipid and amino sugar responses to long-term simulated global environmental changes in a California annual grassland. *Frontiers in Microbiology* 6:385.
- Lücking, R., and D. L. Hawksworth. 2018. Formal description of sequence-based voucherless Fungi: promises and pitfalls, and how to resolve them. *IMA Fungus* 9:143–166.
- Melillo, J. M., S. D. Frey, K. M. DeAngelis, W. J. Werner, M. J. Bernard, F. P. Bowles, G. Pold, M. A. Knorr, and A. S. Grandy. 2017. Long-term pattern and magnitude of soil carbon feedback to the climate system in a warming world. *Science* 358:101–105.
- Melillo, J. M., P. A. Steudler, J. D. Aber, K. Newkirk, H. Lux, F. P. Bowles, C. Catricala, A. Magill, T. Ahrens, and S. Morrisseau. 2002. Soil warming and carbon-cycle feedbacks to the climate system. *Science* 298:2173.
- Mevik, B. H., and R. Wehrens. 2007. The pls package: principal component and partial least squares regression in R. *Journal of Statistical Software* 18:1–24.
- Mucha, J., K. G. Peay, D. P. Smith, P. B. Reich, A. Stefański, and S. E. Hobbie. 2018. Effect of simulated climate warming on the ectomycorrhizal fungal community of boreal and temperate host species growing near their shared ecotonal range limits. *Microbial Ecology* 75:348–363.
- Nguyen, N. H., Z. Song, S. T. Bates, S. Branco, L. Tedersoo, J. Menke, J. S. Schilling, and P. G. Kennedy. 2016. FUNGuild: an open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecology* 20:241–248.
- Nilsson, T. 1973. Studies on wood degradation and cellulolytic activity of microfungi. *Studia forestalia Suecica* 104:1–40.
- Nilsson, R. H., et al. 2011. Towards standardization of the description and publication of next-generation sequencing datasets of fungal communities. *New Phytologist* 191:314–318.
- Nilsson, R. H., et al. 2016. Top 50 most wanted fungi. *MycKeys* 12:29–40.
- Nilsson, R. H., et al. 2019. The UNITE database for molecular identification of fungi: handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Research* 47:D259–D264.
- Oechel, W. C., G. L. Vourlitis, S. J. Hastings, R. C. Zulueta, L. Hinzman, and D. Kane. 2000. Acclimation of ecosystem CO<sub>2</sub> exchange in the Alaskan Arctic in response to decadal climate warming. *Nature* 406:978–981.
- Oksanen, J., F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'Hara, G. L. Simpson, P. Solyomos, M. H. H. Stevens, and H. Wagner. 2014. *Vegan: community ecology package*. R Package Version 2.2-0. <http://CRAN.Rproject.org/package=vegan>
- Parrent, J. L., W. F. Morris, and R. Vilgalys. 2006. CO<sub>2</sub>-enrichment and nutrient availability alter ectomycorrhizal fungal communities. *Ecology* 87:2278–2287.
- Peterjohn, W. T., J. M. Melillo, P. A. Steudler, K. M. Newkirk, F. P. Bowles, and J. D. Aber. 1994. Responses of trace gas fluxes and N availability to experimentally elevated soil temperatures. *Ecological Applications* 4:617–625.
- Peters, R. L. 1990. Effects of global warming on forests. *Forest Ecology and Management* 35:13–33.
- Pold, G., A. F. Billings, J. L. Blanchard, D. B. Burkhardt, S. D. Frey, J. M. Melillo, Ju. Schnabel, L. T. A. van Diepen, and K. M. DeAngelis. 2016. Long-term warming alters carbohydrate degradation potential in temperate forest soils. *Applied and Environmental Microbiology* 82:6518–6530.
- Pold, G., A. S. Grandy, J. M. Melillo, and K. M. DeAngelis. 2017. Changes in substrate availability drive carbon cycle response to chronic warming. *Soil Biology and Biochemistry* 110:68–78.
- Pold, G., J. M. Melillo, and K. M. DeAngelis. 2015. Two decades of warming increases diversity of a potentially lignolytic bacterial community. *Frontiers in Microbiology* 6:480.
- R Core Team. 2019. *R: a language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria.

- Reich, P. B., and J. Oleksyn. 2008. Climate warming will reduce growth and survival of Scots pine except in the far north. *Ecology Letters* 11:588–597.
- Reich, P. B., K. M. Sendall, K. Rice, R. L. Rich, A. Stefanski, S. E. Hobbie, and R. A. Montgomery. 2015. Geographic range predicts photosynthetic and growth response to warming in co-occurring tree species. *Nature Climate Change* 5:148–152.
- Rice, A. V., and R. S. Currah. 2005. *Oidiendron*: a survey of the named species and related anamorphs of *Myxotrichum*. *Studies in Mycology* 53:83–120.
- Rinnan, R., A. Michelsen, E. Bååth, and S. Jonasson. 2007. Fifteen years of climate change manipulations alter soil microbial communities in a subarctic heath ecosystem. *Global Change Biology* 13:28–39.
- Romero-Olivares, A. L., S. D. Allison, and K. K. Treseder. 2017. Soil microbes and their response to experimental warming over time: a meta-analysis of field studies. *Soil Biology and Biochemistry* 107:32–40.
- Schimel, J., and S. M. Schaeffer. 2012. Microbial control over carbon cycling in soil. *Frontiers in Microbiology* 3:348.
- Semenova, T. A., L. N. Morgado, J. M. Welker, M. D. Walker, E. Smets, and J. Geml. 2015. Long-term experimental warming alters community composition of ascomycetes in Alaskan moist and dry arctic tundra. *Molecular Ecology* 24:424–437.
- Solly, E. F., B. D. Lindahl, M. A. Dawes, M. Peter, R. C. Souza, C. Rixen, and F. Hagedorn. 2017. Experimental soil warming shifts the fungal community composition at the alpine treeline. *New Phytologist* 215:766–778.
- Stockinger, H., M. Krüger, and A. Schüßler. 2010. DNA barcoding of arbuscular mycorrhizal fungi. *New Phytologist* 187:461–474.
- Tedersoo, L., et al. 2014. Global diversity and geography of soil fungi. *Science* 346:1256688.
- Tedersoo, L., and M. E. Smith. 2013. Lineages of ectomycorrhizal fungi revisited: foraging strategies and novel lineages revealed by sequences from belowground. *Fungal Biology Reviews* 27:83–99.
- Treseder, K. K., Y. Marusenko, A. L. Romero-Olivares, and M. R. Maltz. 2016. Experimental warming alters potential function of the fungal community in boreal forest. *Global Change Biology* 22:3395–3404.
- Trumbore, S. E., O. A. Chadwick, and R. Amundson. 1996. Rapid exchange between soil carbon and atmospheric carbon dioxide driven by temperature change. *Science* 272:393–396.
- Tu, Q., M. Yuan, Z. He, Y. Deng, K. Xue, L. Wu, S. E. Hobbie, P. B. Reich, and J. Zhou. 2015. Fungal communities respond to long-term CO<sub>2</sub> elevation by community reassembly. *Applied and Environmental Microbiology* 81:2445–2454.
- Van der Putten, W. H. 2012. Climate change, above-ground-belowground interactions, and species' range shifts. *Annual Review of Ecology, Evolution, and Systematics* 43:365–383.
- Wang, Y., U. Naumann, S. T. Wright, and D. I. Warton. 2012. mvabund – an R package for model-based analysis of multivariate abundance data. *Methods in Ecology and Evolution* 3:471–474.
- Warton, D. I., F. G. Blanchet, R. B. O'Hara, O. Ovaskainen, S. Taskinen, S. C. Walker, and F. K. C. Hui. 2015. So many variables: joint modeling in community ecology. *Trends in Ecology & Evolution* 30:766–779.
- Wheeler, J. A., S. D. Frey, and K. A. Stinson. 2017. Tree seedling responses to multiple environmental stresses: interactive effects of soil warming, nitrogen fertilization, and plant invasion. *Forest Ecology and Management* 403:44–51.
- White, T. J., T. Bruns, S. Lee, and J. Taylor. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to Methods and Applications* 18:315–322.
- Wickings, K., A. Stuart Grandy, S. Reed, and C. Cleveland. 2011. Management intensity alters decomposition via biological pathways. *Biogeochemistry* 104:365–379.
- Wold, S., M. Sjöström, and L. Eriksson. 2001. PLS-regression: a basic tool of chemometrics. *Chemometrics and Intelligent Laboratory Systems* 58:109–130.
- Worrall, J. J., S. E. Anagnost, and R. A. Zabel. 1997. Comparison of wood decay among diverse lignicolous fungi. *Mycologia* 89:199–219.
- Žifčáková, L., T. Větrovský, A. Howe, and P. Baldrian. 2016. Microbial activity in forest soil reflects the changes in ecosystem properties between summer and winter. *Environmental Microbiology* 18:288–301.
- Zuur, A., E. N. Ieno, N. Walker, A. A. Saveliev, and G. M. Smith. 2009. *Mixed effects models and extensions in ecology with R*. Springer, New York, New York, USA.

## SUPPORTING INFORMATION

Additional Supporting Information may be found online at: <http://onlinelibrary.wiley.com/doi/10.1002/ecs2.3460/full>